



Basic Research in Computer Science

BRICS RS-00-25 Z. Ésik: Continuous Additive Algebras and Injective Simulations of Synchronization Trees

Continuous Additive Algebras and Injective Simulations of Synchronization Trees

Zoltán Ésik

BRICS Report Series

RS-00-25

ISSN 0909-0878

September 2000

**Copyright © 2000, Zoltán Ésik.
BRICS, Department of Computer Science
University of Aarhus. All rights reserved.**

**Reproduction of all or part of this work
is permitted for educational or research use
on condition that this copyright notice is
included in any copy.**

**See back inner page for a list of recent BRICS Report Series publications.
Copies may be obtained by contacting:**

**BRICS
Department of Computer Science
University of Aarhus
Ny Munkegade, building 540
DK-8000 Aarhus C
Denmark
Telephone: +45 8942 3360
Telefax: +45 8942 3255
Internet: BRICS@brics.dk**

**BRICS publications are in general accessible through the World Wide
Web and anonymous FTP through these URLs:**

`http://www.brics.dk`
`ftp://ftp.brics.dk`
This document in subdirectory RS/00/25/

Continuous additive algebras and injective simulations of synchronization trees

Z. Ésik^{*†}

Dept. of Computer Science
University of Szeged
P.O.B. 652
6701 Szeged, Hungary
`esik@inf.u-szeged.hu`

September 2000

Abstract

The (in)equational properties of the least fixed point operation on (ω) -continuous functions on (ω) -complete partially ordered sets are captured by the axioms of (ordered) iteration algebras, or iteration theories. We show that the inequational laws of the sum operation in conjunction with the least fixed point operation in continuous additive algebras have a finite axiomatization over the inequations of ordered iteration algebras. As a byproduct of this relative axiomatizability result, we obtain complete infinite inequational and finite implicational axiomatizations. Along the way of proving these results, we give a concrete description of the free algebras in the corresponding variety of ordered iteration algebras. This description uses injective simulations

^{*}Partially supported by grant no. FKFP 247/1999 from the Ministry of Education of Hungary and grant no. T30511 from the National Foundation of Hungary for Scientific Research.

[†]The work reported in the paper was partly carried out while visiting BRICS

of regular synchronization trees. Thus, our axioms are also sound and complete for the injective simulation (resource bounded simulation) of (regular) processes.

Keywords: equational logic, fixed points, synchronization trees, simulation.

1 Introduction

Consider the language of μ -terms given by the syntax

$$T ::= x \mid \sigma(\overbrace{T, \dots, T}^{n\text{-times}}) \mid T + T \mid 0 \mid \mu x.T,$$

where x ranges over a countably infinite set of variables, and for each $n \geq 0$, σ ranges over a set of n -ary function symbols of a signature Σ . Such terms may be interpreted as (ω -)continuous functions on (ω -)continuous additive algebras equipped with both an additive structure and a (ω -)complete partial order such that addition and the functions induced by the letters of Σ are ω -continuous, the additive neutral element 0 is also the least element with respect to the partial order, and where terms of the form $\mu x.t$ denote least (pre-)fixed points. We show that under these interpretations the valid (in)equations between μ -terms possess a finite axiomatization over the axioms of ordered iteration algebras (or iteration theories) [9, 19]. In fact, we show that the following set of (in)equations is relatively complete.

$$x + (y + z) = (x + y) + z \tag{1}$$

$$x + y = y + x \tag{2}$$

$$x + 0 = x \tag{3}$$

$$\mu x.x = 0 \tag{4}$$

$$0 \leq x \tag{5}$$

$$\mu x.\mu y.x + y + z = \mu x.x + z \tag{6}$$

(As usual, the scope of the prefix μx extends to the right as far as possible.) Thus, using known completeness results on iteration algebras, see [9, 17, 19], we obtain complete infinite equational and finite implicational axiomatizations. In fact, it follows that the system consisting of the above (in)equations,

the two *Conway identities*

$$\mu x.t[t'/x] = t[\mu x.t'[t/x]/x] \quad (7)$$

$$\mu x.t[x/y] = \mu x.\mu y.t \quad (8)$$

and an equation associated with each finite (simple) group is complete (in the ordered setting). Moreover, the implicational system consisting of (1) – (4), (6), the *fixed point equation*

$$\mu x.t = t[\mu x.t/x] \quad (9)$$

and the *least pre-fixed point rule* (or *fixed point induction*) [3, 32]

$$t[y/x] \leq y \Rightarrow \mu x.t \leq y \quad (10)$$

is also complete. We also show that there is no finite equational axiomatization. Along the way of proving these results, we give a concrete description of the free algebras in the corresponding variety of iteration algebras. This description uses injective simulations [33] of regular synchronization trees [30]. Our results complement those proved in [10] and [20] to the effect that isomorphism classes of regular synchronization trees form the free iteration algebras satisfying equations (1) – (4) and (6) (see also [14] for a recent related result), bisimulation equivalence classes of regular synchronization trees form the free ordered iteration algebras satisfying equations (1) – (4) and

$$\mu x.x + y = y, \quad (11)$$

and that simulation equivalence classes of regular synchronization trees form the free algebras in the variety of iteration algebras satisfying (1) – (5) and (11). As shown in [20], this latter variety is the same as that generated by the “countinuous semilattice algebras” where $+$ is the operation of taking binary sups and 0 is the least element, and where the letters of Σ are interpreted as continuous functions.

The paper is organized as follows. In Section 2 we define the various models used in the sequel including ω -continuous additive algebras and (ordered) iteration algebras. Section 3 contains the main completeness result Theorem 3.1 and its corollaries. The proof of Theorem 3.1 given here relies on the material proved in the subsequent sections, in particular on the description

of the free algebras (Theorem 7.6) and the embedding theorem of Section 8. In Section 4 we define the category of synchronization trees and (functional) simulations, and establish a useful property of injective functional simulations. In Section 5 we recall a general way of constructing iteration algebras from initial algebras over categories. Applied to trees and functional simulations, this construction provides an iteration algebra of synchronization trees for each signature Σ and set A . In Section 6 we show that finite trees equipped with the partial order defined by injective simulations form the free algebras in a finitely axiomatizable variety of ordered algebras. Section 7 contains the corresponding result for regular trees. It is shown here that for each signature Σ , injective simulation equivalence classes of regular trees, equipped with the simulation order, form the free algebras in the variety of ordered iteration algebras satisfying (1) – (6). In Section 8, we show that every ordered iteration algebra of injective simulation equivalence classes of synchronization trees can be embedded in an ω -continuous additive algebra. Finally, in Section 9, we prove that for nontrivial signatures Σ , the variety of ordered iteration algebras generated by the ω -continuous Σ -algebras and axiomatized by the axioms of ordered iteration algebras and the equations (1) – (6) cannot be defined by a finite number of equation schemes.

2 The models

2.1 Countably complete posets and continuous functions

A poset (P, \leq) is called ω -complete if P has a least element \perp_P and sups $\bigvee D$ of all countable directed sets $D \subseteq P$ (or sups of all ω -chains). It is clear that the direct product $\prod_{i \in I} P_i$ of any number of ω -complete posets P_i , $i \in I$, equipped with the pointwise order, is ω -complete. In particular, any direct power P^I of an ω -complete poset is ω -complete. For ω -complete posets P and Q , a function $f : P \rightarrow Q$ is called ω -continuous if it preserves the sup of any countable directed set, or equivalently, the sup of any ω -chain. It follows that any ω -continuous function is monotonic. Clearly, the composite of ω -continuous functions is ω -continuous. When P_i , $i \in I$ are ω -complete,

the projections $\text{pr}_j : \prod_{i \in I} P_i \rightarrow P_j$ are ω -continuous, as is the target tupling of ω -continuous functions.

Given a poset P and a monotonic function $f : P \rightarrow P$, a *pre-fixed point* of f is any element $a \in P$ with $f(a) \leq a$. If $f(a) = a$, then a is called a *fixed point*. The least pre-fixed point of f , when exists, is unique, and necessarily a fixed point. The following fact is well-known.

PROPOSITION 2.1 *Suppose that P and Q are ω -complete posets and f is an ω -continuous function $P \rightarrow Q$. Then for each $b \in Q$ the ω -continuous function $f_b = f(-, b) : P \rightarrow P$ has a least pre-fixed point denoted $f^\dagger(b)$. Moreover, f^\dagger is an ω -continuous function $Q \rightarrow P$.*

In fact, $f^\dagger(b)$ can be constructed as the sup of the ω -chain $(f_b^n(\perp_P))_{n \geq 0}$.

2.2 Additive algebras

A *signature* is a set Σ equipped with a *rank function* assigning a nonnegative integer rank to each function symbol in Σ . For each $n \geq 0$, we will write Σ_n for the collection of all symbols of rank n . A Σ -*algebra* is a set A equipped with an operation $\sigma_A : A^n \rightarrow A$, for each $n \geq 0$. We will write (A, Σ) or just A to denote a Σ -algebra, and we will sometimes omit the subscript A on the operations. Morphisms of Σ -algebras are defined in the usual way.

When Σ is a signature, let $\Sigma_{+,0}$ denote the signature that results by adding to Σ the symbol $+$ of rank 2 and the symbol 0 of rank 0. An *additive Σ -algebra* $A = (A, \Sigma, +, 0)$ is both a Σ -algebra and a commutative monoid $(A, +, 0)$. An *ordered additive Σ -algebra* is an additive Σ -algebra A equipped with a partial order \leq such that the operations, including the sum operation, are monotonic, and such that the inequation (5) holds, so that 0 is the least element of A . Since the sum operation is monotonic, it follows that the inequation

$$x \leq x + y \tag{12}$$

holds in all ordered additive Σ -algebras. Note that additive Σ -algebras form a *variety of $\Sigma_{+,0}$ -algebras*, and ordered additive algebras form a *variety of*

ordered $\Sigma_{+,0}$ -algebras. A morphism of (ordered) additive algebras is a (monotonic) Σ -algebra morphism which also preserves the sum operation and the constant 0. (For varieties of algebras see any standard text on universal algebra such as [23]. For varieties of ordered algebras see [8].)

Any ordered additive Σ -algebra A may also be equipped with the *sum order* [29] defined by

$$a \sqsubseteq b \Leftrightarrow \exists c \ a + c = b.$$

By (12), it follows that $a \leq b$ whenever $a \sqsubseteq b$. The $+$ operation is monotonic with respect to the sum order. However, the other operations may not be monotonic with respect to \sqsubseteq .

2.3 ω -continuous additive algebras

An ω -continuous additive Σ -algebra A has three structures:

1. A Σ -algebra structure (A, Σ) .
2. A commutative monoid structure $(A, +, 0_A)$.
3. An ω -complete partial order structure (A, \leq, \perp_A) .

Moreover, it is required that $\perp_A = 0_A$ and that the operations σ_A as well as the operation $+$ be ω -continuous. Since $\perp_A = 0_A$ and \perp_A is least, (5) holds. A morphism of ω -continuous additive algebras is an ω -continuous additive Σ -algebra morphism.

In any ω -continuous additive Σ -algebra A , we may define all countable sums. Indeed, if $a_i \in A$ for all $i \in I$, where I is countable, define

$$\sum_{i \in I} a_i = \bigvee_F \sum_{i \in F} a_i,$$

where the sup is taken over all finite sets $F \subseteq I$. It is easy to see that for all countable sets I, J such that I is the disjoint union of sets I_j , for $j \in J$, and for all families $a_i \in A$, $i \in I$,

$$\sum_{i \in I} a_i = \sum_{j \in J} \sum_{i \in I_j} a_i.$$

Thus, any ω -continuous additive Σ -algebra has an underlying countably complete monoid [29, 27]. Moreover, any morphism of ω -continuous additive Σ -algebras preserves all countable sums. Note that any ω -continuous additive Σ -algebra is an ordered additive Σ -algebra. For general facts on continuous algebras see [22, 24].

2.4 Preiteration algebras

Terms, or μ -*terms* over a signature Σ are defined by the syntax

$$T ::= x \mid \sigma \left(\overbrace{T, \dots, T}^{n\text{-times}} \right) \mid \mu x. T,$$

where x ranges over a countably infinite set X of variables, and for each $n \geq 0$, σ ranges over Σ_n . Thus, the terms defined in the Introduction are in fact terms over the signature $\Sigma_{+,0}$. The sets of free and bound variables of a term are defined as usual. We identify any two μ -terms that differ only in the names of the bound variables. Moreover, for any μ -terms t, t_1, \dots, t_n and distinct variables x_1, \dots, x_n , we write $t[t_1/x_1, \dots, t_n/x_n]$ or $t[(t_1, \dots, t_n)/(x_1, \dots, x_n)]$ for the term obtained by simultaneously substituting t_i for x_i , for each $i \in [n] = \{1, \dots, n\}$. Since we may assume that the bound variables in t are different from the variables that have a free occurrence in the terms t_i , no free variable in any t_i may become bound as the result of the substitution.

Terms over $\Sigma_{+,0}$ can be interpreted in any ω -continuous additive Σ -algebra A . For each term t , we define an ω -continuous map $t_A : A^X \rightarrow A$ by structural induction.

1. If t is a variable x , then $t_A = \text{pr}_x$ is the corresponding projection $A^X \rightarrow A$.
2. If $t = \sigma(t_1, \dots, t_n)$, where $\sigma \in \Sigma_n$, $n \geq 0$, and t_1, \dots, t_n are μ -terms, then $t_A = \sigma_A \circ \langle (t_1)_A, \dots, (t_n)_A \rangle$, the composite of σ_A with the target tupling of the functions $(t_i)_A$.
3. If $t = t_1 + t_2$, then $t_A = + \circ \langle (t_1)_A, (t_2)_A \rangle$.

4. If $t = 0$, then t_A is the constant function $A^X \rightarrow A$ with value 0_A .
5. If $t = \mu x.t'$, for some variable x and μ -term t' , then let $Y = X - \{x\}$, and let pr_x and pr_Y denote the corresponding projections $A^X \rightarrow A$ and $A^X \rightarrow A^Y$. We define $t_A = f^\dagger \circ \text{pr}_Y$, where $f : A \times A^Y \rightarrow A$ is the continuous function $t'_A \circ \langle \text{pr}_x, \text{pr}_Y \rangle^{-1}$, i.e., the composite of t'_A with the inverse of the order isomorphism $\langle \text{pr}_x, \text{pr}_Y \rangle : A^X \rightarrow A \times A^Y$.

Thus, if $t = \mu x.t'$, then for each $a \in A^X$, $(\mu x.t)_A(a)$ is the least pre-fixed point of the continuous function $A \rightarrow A$,

$$b \mapsto t'_A(a_b^x),$$

where $a_b^x \in A^X$ agrees with a except that it maps x to b .

A *preiteration* $\Sigma_{+,0}$ -algebra is a set A together with an assignment of a function $t_A : A^X \rightarrow A$ to each term t over $\Sigma_{+,0}$ subject to the following rules:

1. For each variable x and $a \in A^X$, $x_A(a) = a(x)$, i.e., x_A is the projection $A^X \rightarrow A$ corresponding to x .
2. If $a, b \in A^X$ are such that $a(x) = b(x)$ for all variables x with a free occurrence in t , then $t_A(a) = t_A(b)$.
3. For all terms t, t_1, \dots, t_n and $a \in A^X$,

$$(t[t_1/x_1, \dots, t_n/x_n])_A(a) = t_A(b),$$

where $b(x_i) = (t_i)_A$, $i \in [n]$, and $b(x) = a(x)$, if $x \notin \{x_1, \dots, x_n\}$.

4. For all terms t, t' over $\Sigma_{+,0}$, if $t_A(a) = t'_A(a)$ for all $a \in A^X$, then also $(\mu x.t)_A(a) = (\mu x.t')_A(a)$, for all $a \in A^X$, i.e., if $t_A = t'_A$, then $(\mu x.t)_A = (\mu x.t')_A$.

A morphism $A \rightarrow B$ of preiteration $\Sigma_{+,0}$ -algebras is a function $h : A \rightarrow B$ such that the following square commutes for each term t :

$$\begin{array}{ccc} A^X & \xrightarrow{t_A} & A \\ h^X \downarrow & & \downarrow h \\ B^X & \xrightarrow{t_B} & B \end{array}$$

Here, h^X denotes the function $a \mapsto a'$ such that $a'(x) = h(a(x))$, for all $x \in X$. If A and B are preiteration $\Sigma_{+,0}$ -algebras such that A is a subset of B and the inclusion of A into B is a morphism, we call A a *preiteration subalgebra* of B . Moreover, if h is a surjective morphism $A \rightarrow B$, then we call B a *quotient* of A .

An *ordered preiteration $\Sigma_{+,0}$ -algebra* is a preiteration $\Sigma_{+,0}$ -algebra A which is a partially ordered set such that the function t_A induced by any term t is monotonic, and for any terms t, t' , if $t_A \leq t'_A$ in the pointwise order, then $(\mu x.t)_A \leq (\mu x.t')_A$. Thus, any ordered preiteration $\Sigma_{+,0}$ -algebra satisfies the implication

$$x \leq y \Rightarrow t \leq t[y/x].$$

(See below for a general treatment of (in)equations and implications.) A morphism of ordered preiteration Σ -algebras is a monotonic preiteration $\Sigma_{+,0}$ -algebra morphism. If A and B are ordered preiteration $\Sigma_{+,0}$ -algebras, we call A an *ordered preiteration subalgebra* of B if A is a preiteration subalgebra of B and the partial order on A is the partial order induced by the order on B , i.e., when the inclusion of A into B is order reflecting. Moreover, we call B a *quotient* of A if there is a surjective ordered preiteration algebra morphism $A \rightarrow B$. Note that every preiteration $\Sigma_{+,0}$ -algebra can be regarded as an ordered preiteration $\Sigma_{+,0}$ -algebra equipped with the discrete partial order which is preserved by any morphism. The following fact is clear.

PROPOSITION 2.2 *Every ω -continuous additive Σ -algebra is an ordered preiteration $\Sigma_{+,0}$ -algebra. If A and B are ω -continuous additive Σ -algebras, and if $h : A \rightarrow B$ is an ω -continuous additive Σ -algebra morphism, then h is an ordered preiteration $\Sigma_{+,0}$ -algebra morphism.*

Suppose that A and B are preiteration $\Sigma_{+,0}$ -algebras and h is a morphism $A \rightarrow B$. The *kernel* of h , denoted θ_h , is defined in the usual way. It is immediate that θ_h is an equivalence relation on A that satisfies the following condition: If $a, b \in A^X$ such that $a(x) \theta_h b(x)$, for all $x \in X$, then for any μ -term t over $\Sigma_{+,0}$, $t_A(a) \theta_h t_A(b)$. Moreover, if $t_A(a) \theta_h t'_A(a)$, for all $a \in A^X$, then $(\mu x.t)_A(a) \theta_h (\mu x.t')_A(a)$ for all $a \in A^X$. An equivalence relation θ on A with these properties is called a *preiteration $\Sigma_{+,0}$ -algebra congruence*, or just

congruence, for short. It is clear that the quotient set A/θ can be turned into a preiteration $\Sigma_{+,0}$ -algebra in a unique way such that the natural map $A \rightarrow A/\theta$ becomes a morphism. If A and B are ordered preiteration $\Sigma_{+,0}$ -algebras and $h : A \rightarrow B$ is a morphism of ordered preiteration $\Sigma_{+,0}$ -algebras, then h determines a preorder \leq_h on A defined by $a \leq_h b$ iff $h(a) \leq h(b)$. This preorder is *compatible* in the sense that if $a, b \in A^X$ such that $a(x) \leq_h b(x)$, for all $x \in X$, then for any μ -term t over $\Sigma_{+,0}$, $t(a) \leq_h t(b)$. Moreover, if $a, b \in A$ with $a \leq b$, then $a \leq_h b$, and if $t_A(a) \leq_h t'_A(a)$, for all $a \in A^X$, then $(\mu x.t)_A(a) \leq_h (\mu x.t')_A(a)$ for all $a \in A^X$. If \leq' is a preorder on A satisfying these conditions, then let θ denote the equivalence relation induced by \leq' on A . The set A/θ , equipped with the partial order induced by \leq' on the θ -equivalence classes, also denoted \leq' , can be turned into an ordered preiteration algebra in a unique way such that the natural map $A \rightarrow A/\theta$ becomes an ordered preiteration $\Sigma_{+,0}$ -algebra morphism.

Suppose that t and t' are terms over $\Sigma_{+,0}$. We say that the *equation* (or *identity*) $t = t'$ holds in a preiteration $\Sigma_{+,0}$ -algebra A , or that A *satisfies* the equation $t = t'$, if for all $a \in A^X$,

$$t_A(a) = t'_A(a),$$

i.e., when the functions t_A and t'_A are equal. Similarly, we say that an in-equation $t \leq t'$ holds in an ordered preiteration $\Sigma_{+,0}$ -algebra A , or that A satisfies $t \leq t'$, if $t_A \leq t'_A$ in the pointwise order of functions. Thus, $t = t'$ holds in an ordered preiteration $\Sigma_{+,0}$ -algebra iff both $t \leq t'$ and $t' \leq t$ hold. More generally, we say that the implication

$$t_1 = t'_1 \wedge \dots \wedge t_n = t'_n \Rightarrow t = t' \quad (13)$$

holds in a preiteration $\Sigma_{+,0}$ -algebra A , where t, t', t_i, t'_i are terms over $\Sigma_{+,0}$, if for all $a \in A^X$, if $(t_i)_A(a) = (t'_i)_A(a)$, for all $i \in [n]$, then $t_A(a) = t'_A(a)$. In a similar way, we can define that an implication

$$t_1 \leq t'_1 \wedge \dots \wedge t_n \leq t'_n \Rightarrow t \leq t' \quad (14)$$

holds in an ordered preiteration $\Sigma_{+,0}$ -algebra.

2.5 Iteration algebras

Some nontrivial (in)equations that hold in all ω -continuous additive Σ -algebras are the (in)equations (1) – (9) given in the Introduction. To define the group-equations, we need to extend the μ -notation to term vectors $t = (t_1, \dots, t_n)$. Let $x = (x_1, \dots, x_n)$ be a vector of distinct variables. When $n = 1$, $\mu x.t$ is just the term vector whose unique component is $\mu x_1.t_1$. We identify any term vector of dimension one with its component. If $n > 1$, let $x' = (x_1, \dots, x_{n-1})$, $t' = (t_1, \dots, t_{n-1})$ and $s = t'[\mu x_n.t_n/x_n]$. (Substitution into a term vector is defined component-wise.) We define

$$\mu x.t = (\mu x'.s, (\mu x_n.t_n)[\mu x'.s/x']). \quad (15)$$

The definition is motivated by the Bekić–de Bakker–Scott identity [6, 3]. Thus, for any ω -continuous $\Sigma_{+,0}$ -algebra A , term vector $t = (t_1, \dots, t_n)$ of dimension n , and for any $x = (x_1, \dots, x_n)$ and $a \in A^X$, $(\mu x.t)_A(a)$ is the least pre-fixed point of the map $A^n \rightarrow A^n$,

$$b = (b_1, \dots, b_n) \mapsto t_A(a_b^x),$$

where of course $a_b^x(x_i) = b_i$, for all $i \in [n]$, and $a_b^x(x) = a(x)$, if $x \notin \{x_1, \dots, x_n\}$.

Suppose that $t = (t_1, \dots, t_n)$ and $t' = (t'_1, \dots, t'_n)$ are term vectors. We will say that $t \leq t'$ holds in an ordered preiteration $\Sigma_{+,0}$ -algebra if each inequation $t_i \leq t'_i$ does. For later use, we note the following fact.

LEMMA 2.3 *If $t \leq t'$ holds in an ordered preiteration $\Sigma_{+,0}$ -algebra A , then $\mu x.t \leq \mu x.t'$ also holds in A .*

Proof. By induction on n using (15). The basis case $n = 1$ is part of the assumption that A is an ordered preiteration $\Sigma_{+,0}$ -algebra. \square

Suppose now that G is a finite group of order n with multiplication denoted \circ . Moreover, suppose that the elements of G are the integers in the set $[n]$. Given a vector $x = (x_1, \dots, x_n)$ of distinct variables and an integer $i \in [n]$, define

$$i \circ x = (x_{i \circ 1}, \dots, x_{i \circ n}).$$

Thus, $i \circ x$ is obtained by permuting the components of x according to the i th row of the multiplication table of G . The *group-identity* or *group-equation* associated with G is the equation

$$(\mu x.(t[1 \circ x/x], \dots, t[n \circ x/x]))_1 = \mu y.t[y/x_1, \dots, y/x_n],$$

where t is any μ -term over $\Sigma_{+,0}$ and y is a (new) variable, and where $(\mu x.(t[1 \circ x/x], \dots, t[n \circ x/x]))_1$ is the first component of the term vector $\mu x.(t[1 \circ x/x], \dots, t[n \circ x/x])$.

An *iteration* $\Sigma_{+,0}$ -*algebra* is a preiteration $\Sigma_{+,0}$ -algebra satisfying the equations (7), (8), as well as any group-identity. A morphism of iteration algebras is a preiteration algebra morphism. An *ordered iteration* $\Sigma_{+,0}$ -*algebra* is an iteration $\Sigma_{+,0}$ -algebra which is an ordered preiteration $\Sigma_{+,0}$ -algebra. Morphisms of ordered iteration $\Sigma_{+,0}$ -algebras also preserve the partial order. We call an (ordered) preiteration subalgebra of an (ordered) iteration algebra an (*ordered*) *iteration subalgebra*. Note that any quotient of an (ordered) iteration algebra is an (ordered) iteration algebra, and every iteration $\Sigma_{+,0}$ -algebra may be regarded as an ordered iteration $\Sigma_{+,0}$ -algebra equipped with the discrete partial order.

We mention a generalization of the group-identities, called the *commutative identities*. Suppose that ρ_1, \dots, ρ_n are mappings $[n] \rightarrow [n]$, where $n \geq 1$. For each ρ_i , let $x_{\rho_i} = (x_{1\rho_i}, \dots, x_{n\rho_i})$. The commutative identity associated with the ρ_i is the equation

$$(\mu x.(t[x_{\rho_1}/x], \dots, t[x_{\rho_n}/x]))_1 = \mu y.t[y/x_1, \dots, y/x_n],$$

where t is any term over $\Sigma_{+,0}$ and y is a new variable. It is shown in [19] that the commutative identities hold in all iteration algebras. (In fact, the original axiomatization of iteration algebras was based on the Conway identities and the “vector forms” of the commutative identities, cf. [9].)

PROPOSITION 2.4 *Any ω -continuous additive Σ -algebra is an ordered iteration $\Sigma_{+,0}$ -algebra satisfying (1) – (6). Moreover, the least pre-fixed point rule (10) holds in any ω -continuous additive Σ -algebra.*

Proof. It is shown in [9, 17] that for any signature Δ , every ω -continuous Δ -algebra is an ordered iteration Δ -algebra. In fact, when $\Delta = \Sigma_{+,0}$, or-

dered iteration Δ -algebras satisfying (5) form the variety generated by the ω -continuous Δ -algebras. (See below for the definition of varieties of (ordered) preiteration algebras.) Applying the easy direction of this general result, we obtain that every ω -continuous additive Σ -algebra is an ordered iteration $\Sigma_{+,0}$ -algebra satisfying (5). It is also immediate that any ω -continuous additive Σ -algebra A satisfies (1) – (4) and the least pre-fixed point rule (10). As for (6), note that for any $a \in A$, both $\mu x.\mu y.x + y + z$ and $\mu x.x + z$ evaluate to the sum of a countable number of copies of a , when z is given the value a . \square

The least pre-fixed point rule may be weakened. We will say that an ordered preiteration $\Sigma_{+,0}$ -algebra satisfies the *weak least pre-fixed point rule* if for all terms $t = t[x]$ and t' , if the inequation $t[t'/x] \leq t'$ holds in A , then so does $\mu x.t \leq t'$, i.e. when $(t[t'/x])_A \leq t'_A$ implies $(\mu x.t)_A \leq t'_A$.

THEOREM 2.5 [17] *Any ordered preiteration $\Sigma_{+,0}$ -algebra A satisfying the fixed point identity (9) and the (weak) least pre-fixed point rule is an ordered iteration algebra.*

A *variety of (ordered) preiteration $\Sigma_{+,0}$ -algebras* is a class \mathcal{V} of (ordered) $\Sigma_{+,0}$ -algebras consisting of the models of some set E of (in)equations between terms over $\Sigma_{+,0}$, i.e., such that an (ordered) preiteration $\Sigma_{+,0}$ -algebra A belongs to \mathcal{V} iff A satisfies any (in)equation in E . The set E is called an *(in)equational basis*, or an *(in)equational axiomatization* of \mathcal{V} . Thus, iteration $\Sigma_{+,0}$ -algebras form a variety of preiteration $\Sigma_{+,0}$ -algebras. This variety is axiomatized by the equations (7), (8) and the group-equations corresponding to the finite groups. Similarly, ordered iteration $\Sigma_{+,0}$ -algebras are axiomatized, in the ordered setting, by the equational axioms of iteration algebras. A variety of (ordered) iteration algebras is a variety of (ordered) preiteration algebras contained in the variety of (ordered) iteration algebras. For a general theory of varieties of iteration algebras we refer to [9]. It is shown in op. cit. that all infinitely generated free algebras exist in any variety of (pre)iteration algebras, but the finitely generated free algebras do not always exist. In fact, for any subset M of the nonnegative integers there is a variety \mathcal{V} of iteration algebras such that for any nonnegative integer n , there is an n -generated free algebra in \mathcal{V} iff $n \in M$.

3 The completeness results

Our main completeness result is the following theorem.

THEOREM 3.1 *An inequation holds in all ω -continuous additive Σ -algebras iff it holds in all ordered iteration $\Sigma_{+,0}$ -algebras satisfying the (in)equations (1) – (6).*

Proof. One direction is a restatement of one part of Proposition 2.4. For the opposite direction, we need to show that every inequation $t \leq t'$ that holds in all ω -continuous additive $\Sigma_{+,0}$ -algebras holds in any ordered iteration $\Sigma_{+,0}$ -algebra A satisfying (1) – (6). But any such A is a quotient of $(\Sigma, A)\mathbf{ISR}$, the free ordered iteration $\Sigma_{+,0}$ -algebra on the set A satisfying (1) – (6), see Theorem 7.6. By the Embedding Theorem, Theorem 8.2, $(\Sigma, B)\mathbf{ISR}$ is an ordered iteration $\Sigma_{+,0}$ -subalgebra of a continuous additive $\Sigma_{+,0}$ -algebra C . Since $t \leq t'$ holds in C , it also holds in A . \square

REMARK 3.2 *The same results holds for continuous additive Σ -algebras having an underlying cpo such that the operations preserve the sup of any directed set. This follows from Theorem 3.1 and the fact that each ω -continuous additive Σ -algebra can be embedded in a continuous additive Σ -algebra.*

COROLLARY 3.3 *An inequation holds in all ω -continuous additive Σ -algebras iff it holds in all ordered preiteration $\Sigma_{+,0}$ -algebras satisfying (1) – (4), (6), the fixed point equation (9), and the least pre-fixed point rule (10).*

Proof. Suppose that an inequation $t \leq t'$ holds in all ω -continuous additive Σ -algebras. Then, by Theorem 3.1, $t \leq t'$ holds in all ordered iteration $\Sigma_{+,0}$ -algebras satisfying the (in)equations (1) – (6). But by Theorem 2.5, any ordered preiteration $\Sigma_{+,0}$ -algebra A satisfying (9) and (10) is an ordered iteration algebra. Moreover, by (4) and (9), each such algebra A satisfies the inequation $0 = \mu x.x \leq y$. It follows that $t \leq t'$ holds in any ordered preiteration $\Sigma_{+,0}$ -algebras satisfying (1) – (4), (6), (9), (10).

Suppose now that $t \leq t'$ holds in all ordered preiteration $\Sigma_{+,0}$ -algebras satisfying (1) – (4), (6), (9), (10). Then, in particular, $t \leq t'$ holds in all ω -continuous additive Σ -algebras. \square

The same argument proves

COROLLARY 3.4 *An inequation holds in all ω -continuous additive Σ -algebras iff it holds in all ordered preiteration $\Sigma_{+,0}$ -algebras satisfying (1) – (4), (6), the fixed point equation (9), and the weak least pre-fixed point rule.*

One can phrase the above results in several other ways. E.g., one possible formulation of Theorem 3.1 is that the defining (in)equations of ordered iteration $\Sigma_{+,0}$ -algebras together with (1) – (6) form an axiomatization of the variety generated by the ω -continuous additive Σ -algebras. Corollary 3.3 may be rephrased as the assertion that the variety of ordered preiteration $\Sigma_{+,0}$ -algebras generated by the continuous additive Σ -algebras and the variety generated by the ordered preiteration $\Sigma_{+,0}$ -algebras satisfying (1) – (4), (6), the fixed point equation (9) and the least pre-fixed point rule (10) are the same. Syntactic formulations may be obtained by using deductive systems. An equivalent syntactic formulation of Theorem 3.1 is the assertion that an inequation holds in all ω -continuous additive Σ -algebras iff it can be derived from the (in)equations (1) – (6) and those defining ordered iteration algebras using the usual rules of reflexivity and transitivity for manipulating inequalities, the rule of substitution, and the congruence rules

$$s \leq s' \vdash t[s/x] \leq t[s'/x] \quad (16)$$

$$t \leq t' \vdash \mu x.t \leq \mu x.t'. \quad (17)$$

(Here, we consider each equation $t = t'$ as an abbreviation for the inequations $t \leq t'$ and $t' \leq t$.) In the same way, Corollary 3.3 is equivalent to the assertion that an inequation holds in all ω -continuous additive Σ -algebras iff it can be derived from (1) – (4), (6) and (9) by the above rules and the rule

$$t[s/x] \leq s \vdash \mu x.t \leq s.$$

(This system is irredundant, one may leave out the second congruence rule (17).)

4 Trees

Suppose that Σ is a signature and A is a set disjoint from Σ . We extend the rank function on Σ to $\Sigma \cup A$ by defining the rank of each letter in A to

be 0. A (Σ, A) -labeled tree $S = (V, E, r, \lambda)$ consists of a countable set V of vertices, a (countable) set E of (hyper)edges, each having a source in V and a target in V^n , for some $n \geq 1$, a distinguished vertex $r \in V$, and a labeling function $\lambda : E \rightarrow \Sigma \cup A$. We require that (V, E, r) is a hypertree with root r , so that each vertex $v \in V$ is accessible from r by a unique path in the underlying directed tree. We write $e : v \rightarrow (v_1, \dots, v_n)$ to indicate that e is an edge with source v and target (v_1, \dots, v_n) . The integer n is called the rank of e . We assume that for each edge e of rank n , $e\lambda \in \Sigma_n$ if $n \neq 1$, and $e\lambda \in \Sigma_0 \cup A \cup \Sigma_1$ if $n = 1$. Moreover, we require that the target of any edge labeled in $\Sigma_0 \cup A$ is a leaf, i.e., not the source of any edge. Note that each (Σ, A) -labeled tree determines an underlying directed tree obtained by replacing each hyperedge $e : v \rightarrow (v_1, \dots, v_n)$ by n edges $e_i : v \rightarrow v_i$, $i \in [n]$. If $S = (V, E, r, \lambda)$ and $S' = (V', E', r', \lambda')$ are (Σ, A) -labeled trees, a *simulation* [33] $S \rightarrow S'$ is a relation $\rho : V \rightarrow V'$ satisfying the following conditions.

- The roots are related, i.e., $r \rho r'$.
- For all $e : v \rightarrow (v_1, \dots, v_n)$ in S and $v' \in V'$, if $v \rho v'$ then there is an edge $e' : v' \rightarrow (v'_1, \dots, v'_n)$ with $e\lambda = e'\lambda'$ and $v_i \rho v'_i$, for all $i \in [n]$.

It follows that the domain of any simulation $S \rightarrow S'$ is the set V . A *functional simulation* is a simulation which is a function. An injective functional simulation is a functional simulation which is injective. Injective functional simulations are closely related to the resource simulation of [14, 15]. The following fact is immediate.

PROPOSITION 4.1 *If $\rho : S \rightarrow S'$ and $\rho' : S' \rightarrow S''$ are (functional) simulations then the composite $\rho' \circ \rho$ is a (functional) simulation $S \rightarrow S''$. Moreover, if ρ and ρ' are injective functional simulations, then so is their composition.*

Thus (Σ, A) -labeled trees and their (functional) simulations form a category, as do (Σ, A) -labeled trees and injective functional simulations. In either category, the isomorphisms are those simulations which are bijective functions.

Suppose that $S = (V, E, r, \lambda)$ is a (Σ, A) -labeled tree and $v \in V$. Let V_v denote the set of all vertices accessible from v , so that a vertex u is in V_v iff

there is a directed path from v to u in the underlying directed tree of S . Let E_v denote the set of all edges $e : u \rightarrow (u_1, \dots, u_n)$ such that $u \in V_v$ (and hence $u_1, \dots, u_n \in V_v$), and let λ_v denote the restriction of λ to E_v . The resulting tree $S_v = (V_v, E_v, v, \lambda_v)$ is called the subtree of S rooted at v .

We let $(\Sigma, A)T$ denote the category of all (Σ, A) -labeled trees and functional simulations. We will sometimes refer to a functional simulation as a morphism. Two subcategories of $(\Sigma, A)T$ are also of interest: the category $(\Sigma, A)F$ determined by the finite trees, and the category $(\Sigma, A)R$ determined by the *regular trees*. We call a tree regular if it has, up to isomorphism, a finite number of subtrees.

PROPOSITION 4.2 [10] *The category $(\Sigma, A)T$ is countably cocomplete.*

Colimits can be constructed in the expected way. For example, if $S_i = (V_i, E_i, r_i, \lambda_i)$, $i \in I$ are trees, where I is a nonempty countable set, then the coproduct $\sum_{i \in I} S_i$ is the disjoint union of the S_i with the roots identified. The coproduct injections are the obvious embeddings. Formally, the coproduct is the tree $S = (V, E, r, \lambda)$, where

$$\begin{aligned} V &= \{r\} \cup \bigcup_{i \in I} (V_i - \{r_i\}) \\ E &= \bigcup_{i \in I} E_i \end{aligned}$$

and for every $e \in E_i$, $e\lambda = e\lambda_i$. Moreover, each edge $e \in E_i$ has the same source and target in S as in S_i , except when the source of e in S_i is r_i . In that case, the source of e in S is the new root r . The empty coproduct, i.e., initial object is the tree \perp , also denoted 0 , with one vertex and no edges.

In addition to coproducts, we will use colimits of ω -chains $(S_n, f_n)_{n \geq 0}$, where $S_n = (V_n, E_n, r_n, \lambda_n)$, but only in the particular case when the morphisms $f_n : S_n \rightarrow S_{n+1}$ are injective. In fact, we may as well assume that $S_n \subseteq S_{n+1}$, for all $n \geq 0$, i.e., that $V_n \subseteq V_{n+1}$, $E_n \subseteq E_{n+1}$, $r_n = r_{n+1}$, and that each edge $e \in E_n$ has the same source, target and label in S_n as in S_{n+1} . Moreover, we may assume that each f_n is the inclusion. Then the colimit $S = (V, E, r, \lambda)$ is the union $\bigcup_{n \geq 0} S_n$, where $V = \bigcup_{n \geq 0} V_n$, $E = \bigcup_{n \geq 0} E_n$, $r = r_0$ and each edge $e \in E_n \subseteq E$, e has the same sort, target and label in S as in S_n . The colimit morphisms are the obvious injections.

Let $S = (V, E, r, \lambda)$, $S' = (V', E', r', \lambda')$ denote (Σ, A) -labeled trees. We call a functional simulation $\rho : S \rightarrow S'$ *normal* if for all vertices $v_1, v_2 \in V$, if S_{v_1} and S_{v_2} are isomorphic and $S'_{v_1\rho}$ and $S'_{v_2\rho}$ are also isomorphic, then there exist isomorphisms $\pi : S_{v_2} \rightarrow S_{v_1}$ and $\pi' : S'_{v_1\rho} \rightarrow S'_{v_2\rho}$ such that $\rho_{v_2} = \pi' \circ \rho_{v_1} \circ \pi$. In the next lemma, we write $S \leq S'$ to denote that there is an injective simulation $S \rightarrow S'$.

LEMMA 4.3 *Suppose that $S = (V, E, r, \lambda)$, $S' = (V', E', r', \lambda')$ are (Σ, A) -labeled trees with $S \leq S'$. Then there exists an injective normal functional simulation $\rho : S \rightarrow S'$.*

Proof. Let $\{S_1 = S, S_2, \dots\}$ and $\{S'_1 = S', S'_2, \dots\}$ be full sets of representatives of isomorphism classes of subtrees of S and S' , respectively. Thus, for each vertex $u \in V$ there exist an integer $j \geq 1$ and an isomorphism $\varphi_u : S_u \rightarrow S_j$, and similarly, for each vertex $u' \in V'$ there exist an integer $k \geq 1$ and an isomorphism $\varphi'_{u'} : S'_{u'} \rightarrow S'_k$. For each pair of integers $i, j \geq 1$ such that $S_i \leq S'_j$, let ρ_{ij} denote an injective functional simulation $S_i \rightarrow S'_j$. We define $v\rho$, for $v \in V$, by induction on the depth¹ of v . It will follow that $S_v \leq S'_{v\rho}$. When the depth is 0, i.e., when $v = r$, we define $v\rho = r'$. Suppose now that the depth of v is positive. Then there is an edge $e : u \rightarrow (v_1, \dots, v_p)$ of S such that v appears in (v_1, \dots, v_p) . Let S_i denote the representative of the isomorphism class of the subtree S_u , and let S'_j denote the representative of the isomorphism class of $S'_{u\rho}$. We define

$$v\rho = v\varphi_u\rho_{ij}\varphi'_{u\rho}.$$

It is clear that ρ is an injective simulation $S \rightarrow S'$. To prove that ρ is normal, suppose that $u_1, u_2 \in V$ such that S_{u_1} is isomorphic to S_{u_2} and $S'_{u_1\rho}$ is isomorphic to $S'_{u_2\rho}$. We need to construct isomorphisms $\pi : S_{u_2} \rightarrow S_{u_1}$ and $\pi' : S'_{u_1\rho} \rightarrow S'_{u_2\rho}$ with $\rho_{u_2} = \pi' \circ \rho_{u_1} \circ \pi$. We only show how to construct π .

So let v denote a vertex of S_{u_2} , i.e., let $v \in V_{u_2}$. When $v = u_2$, define $v\pi = u_1$. We proceed by induction on the depth of v . Our construction will guarantee that S_v is isomorphic to $S_{v\pi}$. In the induction step, there exist an edge $w \rightarrow (v_1, \dots, v_n)$ such that v is one of the v_1, \dots, v_n . Let S_w be isomorphic

¹The depth of v is the length of the unique path from the root to v in the undelying directed graph.

to S_i , so that we have the isomorphisms $\varphi_w : S_w \rightarrow S_i$ and $\varphi_{w\pi} : S_{w\pi} \rightarrow S_i$. We define $v\pi = v\varphi_w\varphi_{w\pi}^{-1}$. \square

5 From categories to algebras

Recall from [5] that if \mathcal{C} is a category and F is a functor $\mathcal{C} \rightarrow \mathcal{C}$, then an F -algebra (c, f) consists of an object c and a morphism $f : Fc \rightarrow c$. Given F -algebras (c, f) and (c', f') , an F -algebra morphism $(c, f) \rightarrow (c', f')$ is a \mathcal{C} -morphism $g : c \rightarrow c'$ such that the following square commutes:

$$\begin{array}{ccc} Fc & \xrightarrow{f} & c \\ Fg \downarrow & & \downarrow g \\ Fc' & \xrightarrow{f'} & c' \end{array}$$

Suppose that \mathcal{C} is *countably cocomplete*, so that \mathcal{C} has all countable colimits. Moreover, assume that the functor $+$: $\mathcal{C}^2 \rightarrow \mathcal{C}$ produces binary coproducts. Since \mathcal{C} is countably cocomplete, so is any direct power of \mathcal{C} . Call a functor $F : \mathcal{C} \rightarrow \mathcal{D}$ ω -*continuous*, where \mathcal{C} and \mathcal{D} are countably cocomplete, if F preserves colimits of ω -diagrams. In particular, $+_{\mathcal{C}}$ is a continuous functor $\mathcal{C}^2 \rightarrow \mathcal{C}$. The following fact is well-known (see, e.g., [9]).

PROPOSITION 5.1 *Suppose that \mathcal{C} and \mathcal{D} are countably cocomplete categories and F is an ω -continuous functor $\mathcal{C} \times \mathcal{D} \rightarrow \mathcal{C}$. Then for each \mathcal{D} -object d , the endofunctor $F(-, d) : \mathcal{C} \rightarrow \mathcal{C}$ has an initial algebra $(F^\dagger d, \mu_d)$. Moreover, the assignment $d \mapsto F^\dagger d$ is the object map of a unique functor $\mathcal{D} \rightarrow \mathcal{C}$ such that $\mu = (\mu_d : d \in \mathcal{D})$ is a natural transformation, in fact a natural isomorphism. Finally, the functor F^\dagger is also ω -continuous.*

The functor F^\dagger can be constructed as follows. Let $g : d \rightarrow d'$ in \mathcal{D} and let 0 denote a fixed initial object of \mathcal{C} . Then let $c_0 = c'_0 = 0$ and define i_0 and i'_0 to be the unique morphisms $c_0 \rightarrow F(c_0, d)$ and $c'_0 \rightarrow F(c'_0, d')$, respectively.

For $n > 0$, define

$$\begin{aligned} c_n &= F(c_{n-1}, d) \\ c'_n &= F(c'_{n-1}, d') \\ i_n &= F(i_{n-1}, g) \\ i'_n &= F(i'_{n-1}, g). \end{aligned}$$

Thus, $i_n : c_n \rightarrow c_{n+1}$ and $i'_n : c'_n \rightarrow c'_{n+1}$, for all $n \geq 0$. Now $F^\dagger d$ is the colimit of the ω -diagram determined by the morphisms i_n , and $F^\dagger d'$ is defined in the same way. Let $(\kappa_n : c_n \rightarrow F^\dagger d)$ and $(\kappa'_n : c'_n \rightarrow F^\dagger d')$ denote the appropriate colimit cocones. As for $F^\dagger g$, let $f_0 : c_0 \rightarrow c'_0$ be the identity morphism $0 \rightarrow 0$, and let

$$f_n = F(f_{n-1}, g),$$

for all $n > 0$. It follows by induction that

$$f_{n+1} \circ i_n = j_n \circ f_n,$$

for all $n \geq 0$. Thus, there results a unique morphism $h : F^\dagger d \rightarrow F^\dagger d'$ with

$$h \circ \kappa_n = \kappa'_n \circ f_n,$$

for all $n \geq 0$. We define $F^\dagger g$ to be this morphism h .

Let Σ denote a signature and suppose that for each $\sigma \in \Sigma_n$, $n \geq 0$ there is an ω -continuous functor $\sigma_{\mathcal{C}} : \mathcal{C}^n \rightarrow \mathcal{C}$. Then, to any term t over $\Sigma_{+,0}$, we can assign an ω -continuous functor $t_{\mathcal{C}} : \mathcal{C}^X \rightarrow \mathcal{C}$ in a canonical way. (Recall that X denotes the set of variables.)

1. If t is a variable x , then $t_{\mathcal{C}}$ is the corresponding projection functor $\text{pr}_x : \mathcal{C}^X \rightarrow \mathcal{C}$.
2. If $t = \sigma(t_1, \dots, t_n)$, where $\sigma \in \Sigma_n$ and t_1, \dots, t_n are terms, then $t_{\mathcal{C}} = \sigma_{\mathcal{C}} \circ \langle (t_1)_{\mathcal{C}}, \dots, (t_n)_{\mathcal{C}} \rangle$, i.e., $t_{\mathcal{C}}$ is the composite of $\sigma_{\mathcal{C}}$ and the target tupling of the functors $(t_1)_{\mathcal{C}}, \dots, (t_n)_{\mathcal{C}}$.
3. If $t = t_1 + t_2$, where t_1 and t_2 are terms, then $t_{\mathcal{C}} = +_{\mathcal{C}} \circ \langle (t_1)_{\mathcal{C}}, (t_2)_{\mathcal{C}} \rangle$.

4. If $t = 0$, then $t_{\mathcal{C}}$ is the constant functor $\mathcal{C}^X \rightarrow \mathcal{C}$ with value $0_{\mathcal{C}}$, a fixed initial object of \mathcal{C} .
5. If $t = \mu x.t'$, then let $Y = X - \{x\}$ and let pr_x and pr_Y denote the projections $\mathcal{C}^X \rightarrow \mathcal{C}$ and $\mathcal{C}^X \rightarrow \mathcal{C}^Y$, respectively. Thus, $\langle \text{pr}_x, \text{pr}_Y \rangle$ is an isomorphism $\mathcal{C}^X \rightarrow \mathcal{C} \times \mathcal{C}^Y$ with inverse $\langle \text{pr}_x, \text{pr}_Y \rangle^{-1}$. We define $t_{\mathcal{C}} = F^\dagger \circ \text{pr}_Y$, where F is the functor

$$t'_{\mathcal{C}} \circ \langle \text{pr}_x, \text{pr}_Y \rangle^{-1} : \mathcal{C} \times \mathcal{C}^Y \rightarrow \mathcal{C}.$$

PROPOSITION 5.2 *For any countably cocomplete category \mathcal{C} , and for any interpretation of the symbols in Σ by ω -continuous functors, all of the iteration algebra identities hold up to isomorphism, as do the equations (1) – (4) and (6).*

Proof. For a proof of the fact that the iteration algebra identities hold, see [9]. The fact that (1) – (4) and (6) hold is obvious. \square

Thus, identifying any two isomorphic objects, we obtain an iteration $\Sigma_{+,0}$ -algebra. For a generalization of Proposition 5.2, see [18].

COROLLARY 5.3 *Suppose that \mathcal{C} is a small countably cocomplete category and for each $\sigma \in \Sigma_n$, $n \geq 0$, $\sigma_{\mathcal{C}}$ is an ω -continuous functor $\mathcal{C}^n \rightarrow \mathcal{C}$. Then the object maps of the functors $t_{\mathcal{C}}$, for all terms t over $\Sigma_{+,0}$, determine an iteration $\Sigma_{+,0}$ -algebra on the isomorphisms classes of the objects of \mathcal{C} , satisfying the equations (1) – (4) and (6).*

The above construction may be applied to the category $\mathcal{C} = (\Sigma, A)T$ of (Σ, A) -labeled trees. The functor $+(\Sigma, A)T : (\Sigma, A)T^2 \rightarrow (\Sigma, A)T$ forms binary coproducts as given above. When $\varphi_i : S_i \rightarrow R_i$, $i = 1, 2$, $\varphi_1 + \varphi_2 : S_1 + S_2 \rightarrow R_1 + R_2$ takes the root of $S_1 + S_2$ to the root of $R_1 + R_2$ and maps each nonroot vertex v of S_i to $v\varphi_i$, $i = 1, 2$. The tree 0 was defined above. For each $\sigma \in \Sigma_n$, we define the functor $\sigma_{(\Sigma, A)T} : (\Sigma, A)T^n \rightarrow (\Sigma, A)T$ as follows. Given trees S_1, \dots, S_n with distinguished vertices r_1, \dots, r_n , respectively, $\sigma(S_1, \dots, S_n)$ is the tree obtained from the S_i by taking their disjoint union and adding a new vertex r and a new edge $e : r \rightarrow (r_1, \dots, r_n)$ labeled σ . Vertex r is the new root. When $\sigma \in \Sigma_0$, we define $\sigma_{(\Sigma, A)T}$ to be the constant functor

determined by the tree which has a unique edge, which is labeled σ . On morphisms, $\sigma_{(\Sigma, A)T}$ is defined in the expected way. See [9] for details.

PROPOSITION 5.4 [10] *Each functor $\sigma_{(\Sigma, A)T}$ is ω -continuous.*

COROLLARY 5.5 [10] *The isomorphism classes of (Σ, A) -labeled trees form an iteration $\Sigma_{+,0}$ -algebra satisfying the equations (1) – (4) and (6).*

We let $(\Sigma, A)\mathbf{T}$ denote this iteration $\Sigma_{+,0}$ -algebra. It was shown in [10] that the regular trees determine an iteration $\Sigma_{+,0}$ -subalgebra of $(\Sigma, A)\mathbf{T}$. We denote this algebra by $(\Sigma, A)\mathbf{R}$. Moreover, the finite trees determine an additive Σ -algebra $(\Sigma, A)\mathbf{F}$.

COROLLARY 5.6 *$(\Sigma, A)\mathbf{R}$ is an iteration $\Sigma_{+,0}$ -algebra satisfying the equations (1) – (4) and (6). $(\Sigma, A)\mathbf{F}$ is a $\Sigma_{+,0}$ -algebra satisfying the equations (1) – (3).*

The following result gives an algebraic characterization of the iteration $\Sigma_{+,0}$ -algebra $(\Sigma, A)\mathbf{R}$. Let us identify each letter $a \in A$ with the tree which has a unique vertex and a unique edge, which is labeled a .

THEOREM 5.7 [10] *The iteration $\Sigma_{+,0}$ -algebra $(\Sigma, A)\mathbf{R}$ is freely generated by the set A in the variety of all iteration $\Sigma_{+,0}$ -algebras satisfying the equations (1) – (4) and (6).*

The meaning of this result is that for any iteration $\Sigma_{+,0}$ -algebra satisfying the equations (1) – (4), (6), and for any function $h : A \rightarrow B$, there is a unique iteration $\Sigma_{+,0}$ -algebra morphism $h^\sharp : (\Sigma, A)\mathbf{R} \rightarrow B$ extending h .

COROLLARY 5.8 *An equation holds in all (pre)iteration $\Sigma_{+,0}$ -algebras derived from small countably complete categories as in Corollary 5.3 iff it holds in all iteration $\Sigma_{+,0}$ -algebras satisfying (1) – (4) and (6).*

Suppose that t is a μ -term over $\Sigma_{+,0}$ and n is a nonnegative integer. Below we will write nt for the n -fold sum of t with itself, where we take advantage of the associativity of $+$. When $n = 0$, we define nt to be the term 0 . Moreover, we define $\infty t = \mu x.x + t$, where x does not occur in t . We let $\infty + n = n + \infty = \infty$ and $n \leq \infty$ for all $n \in \{0, 1, \dots, \infty\}$. For later use, we note two simple facts.

COROLLARY 5.9 *The following equation holds in any iteration $\Sigma_{+,0}$ -algebra satisfying (1) – (4) and (6):*

$$ny + my = (n + m)y \quad (18)$$

where n, m are nonnegative integers or ∞ . When $n, m \neq \infty$, (18) holds in all additive Σ -algebras.

Proof. Equation (18) holds in all algebras $(\Sigma, A)\mathbf{R}$. □

COROLLARY 5.10 *For all $n \leq m$, the inequation*

$$ny \leq my \quad (19)$$

holds in all ordered iteration $\Sigma_{+,0}$ -algebras satisfying (1) – (6). When $n, m \neq \infty$, (19) holds in all ordered additive Σ -algebras.

Proof. If $n \leq m$, then there exists some k with $n + k = m$. Thus, by Corollary 5.9, $ny + ky = (n + k)y = my$ holds. But then $ny \leq my$ also holds, by (12). □

5.1 The injective simulation preorder

We will consider a preorder on trees. Suppose that S and T are (Σ, A) -labeled trees. We define $S \leq T$ if there is an injective functional simulation $S \rightarrow T$. Moreover, we define $S \equiv T$ if both $S \leq T$ and $T \leq S$ hold. The relation \leq is called the *injective simulation preorder* and \equiv the *injective simulation equivalence*.

EXAMPLE 5.11 There exist nonisomorphic regular trees S, T with $S \equiv T$. For one example, let $\sigma \in \Sigma_1$, and let $S = \sigma(0) + \infty\sigma(\sigma(0))$ and $T = \infty\sigma(\sigma(0))$.

The following facts are obvious.

LEMMA 5.12 Suppose that $\sigma \in \Sigma_n$ and $S_i \leq R_i$, for $i \in [n]$. Then $\sigma(S_1, \dots, S_n) \leq \sigma(R_1, \dots, R_n)$. In fact, if $\varphi_i : S_i \rightarrow R_i$, $i \in [n]$ are injective, then so is $\sigma(\varphi_1, \dots, \varphi_n) : \sigma(S_1, \dots, S_n) \rightarrow \sigma(R_1, \dots, R_n)$.

LEMMA 5.13 If $S_1 \leq S_2$ and $R_1 \leq R_2$, then $S_1 + S_2 \leq R_1 + R_2$. Moreover, if $\varphi_i : S_i \rightarrow R_i$, $i = 1, 2$ are injective, then so is $\varphi_1 + \varphi_2 : S_1 + S_2 \rightarrow R_1 + R_2$.

PROPOSITION 5.14 For all μ -terms t over $\Sigma_{+,0}$ and for all families of (Σ, A) -labeled trees $S = (S_y)_{y \in X}$ and $R = (R_y)_{y \in X}$ with $S_y \leq R_y$ for $y \in X$, it holds that $t_{(\Sigma, A)T}(S) \leq t_{(\Sigma, A)T}(R)$. Moreover, if $\varphi = (\varphi_y)_{y \in X}$ such that each functional simulation $\varphi_y : S_y \rightarrow R_y$, $y \in X$ is injective, then so is $t_{(\Sigma, A)T}(\varphi) : t_{(\Sigma, A)T}(S) \rightarrow t_{(\Sigma, A)T}(R)$.

Proof. We argue by induction on the structure of t . Our claim is obvious when t is a variable or $t = 0$. If t is of the form $\sigma(t_1, \dots, t_n)$, or $t_1 + t_2$, the result follows by the previous lemmas. Assume finally that $t = \mu x.t'$, for some variable x and term t' . Let F denote the functor $(\Sigma, A)T \rightarrow (\Sigma, A)T$ defined by

$$f : U \rightarrow V \mapsto t'(S_f^x) : t'(S_U^x) \rightarrow t'(S_V^x).$$

Here, S_f^x denotes the assignment that takes x to f and any variable $y \neq x$ to the identity $S_y \rightarrow S_y$. Then $t(S)$ is the colimit of the ω -diagram $i_n : U_n \rightarrow U_{n+1}$, where $U_0 = 0$ is the initial tree, $U_{n+1} = FU_n$ and $i_{n+1} = Fi_n$. Of course, i_0 is the unique morphism $U_0 \rightarrow U_1$. In the same way, $t(R)$ is the colimit of the ω -diagram $j_n : V_n \rightarrow V_{n+1}$ constructed in the same way using the functor $G : (\Sigma, A)T \rightarrow (\Sigma, A)T$

$$f : U \rightarrow V \mapsto t'(R_f^x) : t'(R_U^x) \rightarrow t'(R_V^x).$$

Since i_0 and j_0 are injective, it follows by the induction assumption that each of the morphisms i_n and j_n is also injective. Assume that $\varphi_y : S_y \rightarrow R_y$ is an

injective functional simulation, for each y . Then let g_0 denote the identity functional simulation $0 \rightarrow 0$, and let

$$g_{n+1} = t'(\varphi_{g_n}^x), \quad n \geq 0.$$

By the induction assumption we have that these morphisms are also injective. Since

$$g_{n+1} \circ i_n = j_{n+1} \circ g_n,$$

for all $n \geq 0$, there results a mediating morphism $g : t(S) \rightarrow t(R)$. We may as well assume that the i_n and j_n are inclusions, in which case $t(S)$ is the union of the U_n and $t(R)$ the union of the V_n , moreover, for any vertex u of U_n , $n \geq 0$, we have $g(u) = g_n(u)$. Thus, g is injective, proving that $t(\varphi)$ is also injective. \square

PROPOSITION 5.15 *Let s, t denote μ -terms over $\Sigma_{+,0}$. Suppose that for all families $R = (R_y)_{y \in X}$ of (regular) (Σ, A) -labeled trees we have $s(R) \leq t(R)$. Then also $(\mu x.t)(R) \leq (\mu x.s)(R)$, for all families R of regular (Σ, A) -labeled trees and for all $x \in X$.*

Proof. For a family $R = (R_y)_{y \in X}$ of regular trees, let F denote the functor $(\Sigma, A)T \rightarrow (\Sigma, A)T$, defined on objects by $S \mapsto s(R_S^x)$. On morphisms, F is defined in a similar way. Let G denote the corresponding functor using term t . Then $F^\dagger = (\mu x.s)(R_y/y)$ is the colimit of the ω -diagram $F^n(i_0) : F^n(0) \rightarrow F^{n+1}(0)$, where 0 is the initial tree and i_0 denotes the unique functional simulation $0 \rightarrow F(0)$. Also, $G^\dagger = (\mu x.t)(R_y/y)$ is the colimit of the ω -diagram $G^n(j_0) : G^n(0) \rightarrow G^{n+1}(0)$ defined in the same way. By Proposition 5.14, each $F^n(i_0)$ and $G^n(j_0)$ is injective, so that we may assume as before that each $F^n(i_0)$ and $G^n(j_0)$ is an inclusion, and that F^\dagger and G^\dagger are the unions of the $F^n(0)$ and the $G^n(0)$, respectively. Suppose that S is a finite tree with $S \leq F^\dagger$. Then there is some n such that $S \leq F^n(0)$. But it follows from our assumption on s and t that $F^n(0) \leq G^n(0)$, so that $S \leq G^n(0)$ and $S \leq G^\dagger$. Since F^\dagger and G^\dagger are regular, by Lemma 8.1, proved independently, we have $F^\dagger \leq G^\dagger$. \square

By the previous facts, the injective simulation equivalence classes of regular (Σ, A) -labeled trees form an iteration $\Sigma_{+,0}$ -algebra. Let $(\Sigma, A)\mathbf{ISR}$ denote

this iteration $\Sigma_{+,0}$ -algebra. In fact, identifying any tree with its isomorphism class, \leq induces a compatible preorder on $(\Sigma, A)\mathbf{R}$, so that $(\Sigma, A)\mathbf{ISR}$ is the quotient of $(\Sigma, A)\mathbf{R}$ under the natural map taking a tree T to its injective simulation equivalence class T/\equiv . Equipped with the partial order induced by \leq , $(\Sigma, A)\mathbf{ISR}$ is an ordered iteration $\Sigma_{+,0}$ -algebra. In fact, we have:

PROPOSITION 5.16 *$(\Sigma, A)\mathbf{ISR}$ is an ordered iteration $\Sigma_{+,0}$ -algebra satisfying (1) – (6).*

Proof. We have already noted that $(\Sigma, A)\mathbf{ISR}$ is an ordered iteration $\Sigma_{+,0}$ -algebra which is a quotient of $(\Sigma, A)\mathbf{R}$. Thus, by Corollary 5.6, (1) – (4) and (6) hold in $(\Sigma, A)\mathbf{ISR}$. The inequation (5) is obvious. \square

Let $(\Sigma, A)\mathbf{ISF}$ denote the subalgebra of $(\Sigma, A)\mathbf{ISR}$ determined by the finite trees. Clearly, $(\Sigma, A)\mathbf{ISF}$ is an ordered additive Σ -algebra.

6 Finite trees

In this section we show that the injective simulation equivalence classes of finite (Σ, A) -labeled trees form the free ordered additive Σ -algebra on A . In fact, we show that two finite trees are injective simulation equivalent iff they are isomorphic. Thus, $(\Sigma, A)\mathbf{ISF}$ is just $(\Sigma, A)\mathbf{F}$ equipped the partial order induced by injective simulations. First we consider the unordered case.

PROPOSITION 6.1 [10] *The algebra $(\Sigma, A)\mathbf{F}$ of isomorphism classes of finite (Σ, A) -labeled trees is freely generated by A in the variety of additive Σ -algebras.*

PROPOSITION 6.2 *If S and S' are finite (Σ, A) -labeled trees with $S \equiv S'$ then S and S' are isomorphic.*

Proof. Let φ and ψ be injective morphisms $S \rightarrow S'$ and $S' \rightarrow S$, respectively. Then the composite $\psi \circ \varphi$ is necessarily an isomorphism $S \rightarrow S$, and $\varphi \circ \psi$ is an isomorphism $S' \rightarrow S'$. But then φ and ψ are also isomorphisms. \square

REMARK 6.3 *In the same way, for any “finitely branching” (Σ, A) -labeled trees S, T , if $S \equiv T$ then S is isomorphic to T .*

We use the above results to prove

PROPOSITION 6.4 *The algebra $(\Sigma, A)\mathbf{ISF}$, equipped with the partial order \leq , is freely generated by A in the variety of ordered additive Σ -algebras.*

Proof. We already know that $(\Sigma, A)\mathbf{ISF}$ is an ordered additive Σ -algebra. By Proposition 6.2, $(\Sigma, A)\mathbf{ISF}$ may be represented as the ordered additive Σ -algebra of isomorphism classes of finite trees, equipped with the partial order \leq . We may as well identify each tree with its isomorphism class and regard the elements of $(\Sigma, A)\mathbf{ISF}$ as finite (Σ, A) -labeled trees.

Suppose that B is an ordered additive Σ -algebra and h is a function $A \rightarrow B$. By Proposition 6.1, h extends to a unique additive Σ -algebra morphism $h^\# : (\Sigma, A)\mathbf{ISF} \rightarrow B$. We need to show that $h^\#$ preserves the partial order. Let S and S' be finite trees with $S \leq S'$. We argue by induction on the depth of S to show that $h^\#(S) \leq h^\#(S')$. When $S = 0$, our claim holds by (5). Suppose now that the depth of S' is positive. We can write S as

$$S = \sum_{i \in I} \sigma_i(S_{i1}, \dots, S_{in_i}),$$

where I is a finite nonempty set, and for each $i \in I$, σ_i is a symbol in $\Sigma(A)_{n_i}$ and S_{i1}, \dots, S_{in_i} are finite trees of depth less than the depth of S . Since $S \leq S'$, we can also write

$$S' = T + \sum_{i \in I} \sigma_i(S'_{i1}, \dots, S'_{in_i}),$$

where T and the S'_{ij} , $j \in [n_i]$, $i \in I$ are finite trees such that $S_{ij} \leq S'_{ij}$, for all i, j . Thus, using the induction hypothesis and the fact that the operations are monotonic, as well as (12),

$$\begin{aligned} h^\#(S) &= \sum_{i \in I} \sigma_i(h^\#(S_{i1}), \dots, h^\#(S_{in_i})) \\ &\leq \sum_{i \in I} \sigma_i(h^\#(S'_{i1}), \dots, h^\#(S'_{in_i})) \\ &\leq h^\#(T) + \sum_{i \in I} \sigma_i(h^\#(S'_{i1}), \dots, h^\#(S'_{in_i})) \\ &= h^\#(S'). \quad \square \end{aligned}$$

COROLLARY 6.5 *Additive Σ -algebras and ordered additive Σ -algebras satisfy the same equations.*

7 Regular trees

Let Σ denote a signature and let A be a set. In this section we prove that the algebra $(\Sigma, A)\mathbf{ISR}$ of injective simulation equivalence classes of regular (Σ, A) -labeled trees, equipped with the injective simulation order, is freely generated by A in the variety of ordered iteration $\Sigma_{+,0}$ -algebras satisfying (1) – (6).

Given (Σ, A) -labeled trees S and S' and a functional simulation $\rho : S \rightarrow S'$, for any vertex u of S let ρ_u denote the functional simulation $S_u \rightarrow S'_{u\rho}$ induced by ρ , so that ρ_u is the restriction of ρ to the vertices of S_u .

A (Σ, A) -normal representation of dimension n in the variables $x_1, \dots, x_n, y_1, \dots, y_p$ is an ordered pair $D = (t, a)$, where $t = (t_1, \dots, t_n)$ is an n -dimensional vector of terms over $\Sigma_{+,0}$ in the free variables $x_1, \dots, x_n, y_1, \dots, y_p$, and where $a = (a_1, \dots, a_p) \in A^p$. Moreover, each term t_i is *primitive*, i.e., a finite sum of terms of the form $k\sigma(x_{j_1}, \dots, x_{j_m})$ or ky_j , where $k \neq 0$, $\sigma \in \Sigma_m$, $m \geq 0$, $j_1, \dots, j_m \in [n]$ and $j \in [p]$. (It is allowed that $k = \infty$, and we take advantage of the associativity and commutativity of $+$.) Let us denote $x = (x_1, \dots, x_n)$ and $y = (y_1, \dots, y_p)$. The *behavior* of D , denoted $|D|$ is the first component of $(\mu x.t)_{(\Sigma, A)\mathbf{R}}(a/y)$. Thus, the behavior of D is a regular tree in $(\Sigma, A)\mathbf{R}$. The term $\mu x.t$ itself will be called normal of dimension n . The variables y_j are called the *parameters*.

Each regular tree $T \in (\Sigma, A)\mathbf{R}$ is known to be the behavior of a normal representation $D = (t, a)$. To construct D , let T_1, \dots, T_n be an enumeration of the subtrees of T with $T = T_1$, and let a_1, \dots, a_p be an enumeration of those elements of A which appear as labels of some edges of T . We define $t = (t_1, \dots, t_n)$ and $a = (a_1, \dots, a_p)$, where each t_i corresponds to T_i in the following manner. Each edge $e : v \rightarrow (v_1, \dots, v_m)$ whose source is the root of T_i is labeled by some symbol $\sigma \in \Sigma$, or some component of a . Suppose that e is labeled $\sigma \in \Sigma_m$, $m > 0$. Then let T_{j_1}, \dots, T_{j_m} denote the subtrees rooted at the vertices v_1, \dots, v_m , respectively, and let k denote the total

number of edges $v \rightarrow (v'_1, \dots, v'_m)$ labeled $\sigma \in \Sigma_m$ such that the subtrees rooted at the vertices v'_1, \dots, v'_m are isomorphic to the trees T_{j_1}, \dots, T_{j_m} , respectively. (If there are an infinite number of such edges, then $k = \infty$.) Then $k\sigma(x_{j_1}, \dots, x_{j_m})$ is a summand of t_i . If $\sigma \in \Sigma_0$, then $k\sigma$ is a summand of t_i , where k is the number of those edges with source v labeled σ . Similarly, for each a_j , ky_j is a summand of t_i , where k is determined in the same way. The term t_i is the sum of all such summands. We call the normal representation D constructed above the *minimal representation of T* . (Note that the minimal representation is unique only up to a rearrangement of the components of t and a and renaming of the variables.)

EXAMPLE 7.1 Suppose that $\sigma \in \Sigma_1$ and that T is a regular tree with non-isomorphic subtrees T_1, T_2 , where T_1 is T , T_2 is the tree 0 , and where there are 4 edges leaving the root. Two of the edges, labeled σ , lead to vertices whose corresponding subtrees are isomorphic to T , a third edge, also labeled σ , leads to a vertex with no outgoing edge, and the last edge is labeled $a \in A$. Then the normal representation for T is

$$(\mu(x_1, x_2).(2\sigma(x_1) + \sigma(x_2) + a, 0), a).$$

It is known that for each i , the i th component of $(\mu x.t)(a)$ is the tree T_i . In particular, we have:

PROPOSITION 7.2 [9] *For the normal representation $D = (t, a)$ of the tree $T \in (\Sigma, A)\mathbf{R}$, it holds that $|D| = T$.*

Given normal terms $\mu x.t$ and $\mu y.t'$, of dimension n and m respectively, and a function $\rho : [n] \rightarrow [m]$, where $x = (x_1, \dots, x_n)$, $y = (y_1, \dots, y_m)$, $t = (t_1, \dots, t_n)$ and $t' = (t'_1, \dots, t'_m)$, we define $\rho \circ t'$ to be the term vector $(t'_{1\rho}, \dots, t'_{n\rho})$. Moreover, we define

$$t \circ \rho = (t_1[y_{1\rho}/x_1, \dots, y_{n\rho}/x_n], \dots, t_n[y_{1\rho}/x_1, \dots, y_{n\rho}/x_n]).$$

Recall that when $t = (t_1, \dots, t_n)$, $t' = (t'_1, \dots, t'_n)$ and A is a preiteration $\Sigma_{+,0}$ -algebra, we say that $t = t'$ holds in A if each $t_i = t'_i$ does. We use a similar convention for inequations $t \leq t'$.

LEMMA 7.3 *Suppose that S and S' are trees with $S \leq S'$. Then there exist normal representations $(\mu x.t, a)$ and $(\mu y.t', a)$ for S and S' and a function $\rho : [n] \rightarrow [m]$, where n denotes the dimension of $\mu x.t$ and m denotes the dimension of $\mu y.t'$, such that $1\rho = 1$ and*

$$t \circ \rho \leq \rho \circ t'$$

holds in all ordered preiteration $\Sigma_{+,0}$ -algebras satisfying (1), (2), (3), (18) and (19).

Proof. Let $\{S_1 = S, S_2, \dots, S_k\}$ and $\{S'_1 = S', S'_2, \dots, S'_m\}$ denote full sets of representatives of isomorphism classes of subtrees of S and S' , respectively. By Lemma 4.3, there is an injective normal morphism $\varphi : S \rightarrow S'$. Let I denote the set of all ordered pairs (i, j) , $i \in [k]$, $j \in [m]$ such that there is a vertex u of S such that S_u is isomorphic to S_i and $S'_{u\varphi}$ is isomorphic to S'_j . Let n denote the number of elements of the set I and let us fix an enumeration of I starting with the pair $(1, 1)$. Call an edge $e : u \rightarrow (v_1, \dots, v_p)$ of S an

$$(i, j) \rightarrow ((i_1, j_1), \dots, (i_p, j_p))$$

type edge, where $i, i_1, \dots, i_p \in [k]$ and $j, j_1, \dots, j_p \in [m]$, if S_u is isomorphic to S_i , $S_{u\varphi}$ is isomorphic to S'_j , and for each ℓ , S_{v_ℓ} is isomorphic to S_{i_ℓ} and $S'_{v_\ell\varphi}$ is isomorphic to S'_{j_ℓ} . Note that (i, j) is necessarily in I as are the ordered pairs $(i_1, j_1), \dots, (i_p, j_p)$.

We define the term $\mu x.t$. The components of the vector x correspond to the elements of I in a bijective manner according to the fixed enumeration of I . Let x_{ij} denote the variable corresponding to the ordered pair (i, j) , and let t_{ij} denote the corresponding component of t . Moreover, let a_1, \dots, a_q be all the letters of A that appear in S or S' , in some fixed enumeration, and let z_1, \dots, z_q be variables corresponding to these letters. (We assume that these variables are different from the components of x .) To define t_{ij} , consider a vertex u of S such that S_u is isomorphic to S_i and $S'_{u\varphi}$ is isomorphic to S'_j . For each $\sigma \in \Sigma_p$, $p > 0$, which appears as the label of an outgoing edge of u , the root of S_u , and for each sequence $\alpha = ((i_1, j_1), \dots, (i_p, j_p)) \in I^p$, let $s = s(\sigma, (i, j) \rightarrow \alpha)$ denote the number of edges $e : u \rightarrow (v_1, \dots, v_p)$ labeled σ and having type $(i, j) \rightarrow ((i_1, j_1), \dots, (i_p, j_p))$. When $\sigma_0 \in \Sigma_0$, let $s(\sigma_0, i)$ denote the total number of edges with source u labeled σ_0 . Moreover, for

each $\ell \in [q]$, let $s_{i\ell}$ denote the number of edges with source u labeled a_ℓ . We define

$$t_{ij} = \sum_{\sigma, \alpha} s(\sigma, (i, j) \rightarrow \alpha) \sigma(x_{i_1 j_1}, \dots, x_{i_p j_p}) + \sum_{\sigma_0} s(\sigma_0, i) \sigma + \sum_{\ell} s_{i\ell} z_\ell.$$

As for $\mu y.t'$, we take the minimal normal representation of S' . Thus, the components of y and t' correspond to the subtrees of S' , so that $y = (y_1, \dots, y_m)$ and $t' = (t'_1, \dots, t'_m)$ with y_j and t'_j corresponding to S'_j , for each $j \in [m]$. Moreover, each t'_j is given by

$$t'_j = \sum_{\sigma, \beta} s'(\sigma, j \rightarrow \beta) \sigma(y_{j_1}, \dots, y_{j_p}) + \sum_{\sigma_0} s'(\sigma_0, j) + \sum_{\ell} s'_{\ell} z_\ell.$$

Here, $\sigma \in \Sigma_p$, $p > 0$ ranges over the symbols which appear as labels of outedges of the root of S'_j , $\beta = (j_1, \dots, j_p) \in [m]^p$, and where $s'(\sigma, j \rightarrow \beta)$ is the number of outedges of the root of S'_j labeled σ of type $j \rightarrow (j_1, \dots, j_p)$, i.e., whose targets determine a sequence of vertices whose corresponding subtrees are isomorphic to $S'_{j_1}, \dots, S'_{j_p}$, respectively. Moreover, σ_0 ranges over those symbols in Σ_0 that appear as labels of outedges of the root of S'_j , $s'(\sigma_0, j)$ is the corresponding multiplicity, and for each $\ell \in [q]$, s'_{ℓ} is the number of outedges of the root of S'_j labeled a_ℓ .

By Proposition 7.2, we have that $|(\mu y.t', a)| = S'$. Define the function $\rho : [n] \rightarrow [m]$ such that it maps any integer in $[n]$ corresponding to an ordered pair (i, j) in the fixed enumeration of the set I to the integer j . Suppose that $(i, j) \in I$. Since $S_i \leq S_j$, it is clear that for any $\sigma \in \Sigma_p$ with $p > 0$ and j_1, \dots, j_p ,

$$\sum_{i_1, \dots, i_p} s(\sigma, (i, j) \rightarrow (i_1, j_1), \dots, (i_p, j_p)) \leq s'(\sigma, j \rightarrow (j_1, \dots, j_p)).$$

And for all $\sigma_0 \in \Sigma_0$ and $\ell \in [q]$,

$$s(\sigma_0, i) \leq s'(\sigma_0, j),$$

and

$$s_{i\ell} \leq s'_{\ell},$$

It follows that $t \circ \rho \leq \rho \circ t'$ holds modulo (1), (2), (3), (18) and (19) and the monotonicity of $+$, i.e., $t \circ \rho \leq \rho \circ t'$ holds in any ordered preiteration $\Sigma_{+,0}$ -algebra satisfying the (in)equations (1), (2), (3), (18) and (19).

To complete the proof, we still need to show that $|(\mu x.t, a)| = S$. But consider the minimal normal representation $(\mu x''.t'', a)$ for S . Here $x'' = (x''_1, \dots, x''_k)$, $t'' = (t''_1, \dots, t''_k)$ such that for each i , x''_i and t''_i correspond to S_i , i.e., the i th component of $(\mu x''.t'')(a)$ is isomorphic to S_i . If τ denotes the function $[n] \rightarrow [k]$ corresponding to the assignment $I \rightarrow [k]$, $(i, j) \mapsto i$, then we clearly have $t \circ \tau = \tau \circ t''$ modulo (1), (2), (3) and (18). Thus, by Lemma 7.5, for each (i, j) in I , the (i, j) th component of $(\mu x.t)(a)$ is isomorphic to the i th component of $(\mu x''.t'')(a)$. \square

LEMMA 7.4 *Let $\mu x.t$ and $\mu y.t'$ be normal terms of dimension n and m , respectively. Suppose that there is a function $\rho : [n] \rightarrow [m]$ with $t \circ \rho \leq \rho \circ t'$ modulo (1), (2), (3), (18) and (19). Then there is a normal term $\mu x.s$ of dimension n such that $t \leq s$ and $s \circ \rho = \rho \circ t'$ hold modulo (1), (2), (3), (18) and (19).*

Proof. Assume that $t = (t_1, \dots, t_n)$ and $t' = (t'_1, \dots, t'_m)$. For each $i \in [n]$ there is a term t''_i with $(t_i + t''_i) \circ \rho = t'_{i\rho}$ modulo (1), (2), (3) and (18). Moreover, t''_i is the sum of primitive terms. Define $s_i = t_i + t''_i$ and $s = (s_1, \dots, s_n)$. Then $t \leq s$ modulo (1), (2), (3), (18) and (19). \square

LEMMA 7.5 *Suppose that $\mu x.t$ and $\mu y.t'$ are normal terms of dimension n and m in the parameters $z = (z_1, \dots, z_p)$. Let ρ denote a function $[n] \rightarrow [m]$. If $t \circ \rho = \rho \circ t'$, modulo (1), (2), (3) and (18), then for any $a \in A^p$ and $i \in [n]$, the i th component of $(\mu x.t)(a)$ is isomorphic to the $(i\rho)$ th component of $(\mu y.t')(a)$. Thus, if $1\rho = 1$, then trees represented by $(\mu x.t, a)$ and $(\mu y.t', a)$ are isomorphic.*

Proof. We know that $(\mu x.t)(a)$ is the colimit of the ω -diagram $(i_k : s_k \rightarrow s_{k+1})_{k \geq 0}$, where s_0 is the n -tuple $(0, \dots, 0)$, $s_{k+1} = t(s_k, a/x, z)$, $i_{k+1} = t(i_k, a/x, z)$, $k \geq 0$. Also, $(\mu y.t')(a)$ is the colimit of an ω -diagram $(i'_k : s'_k \rightarrow s'_{k+1})_{k \geq 0}$ constructed in the same way. But since $t \circ \rho = \rho \circ t'$, modulo the associativity and commutativity of $+$ and (18), we have that $s_k = \rho \circ s'_k$,

for all $k \geq 1$, i.e., for any $i \in [n]$, the i th component of s_k is the same as the $(i\rho)$ th component of s'_k . In the same way, $i_k = \rho \circ i'_k$. It follows that $(\mu x.t)(a) = \rho \circ (\mu x.t')(a)$. \square

THEOREM 7.6 *For each signature Σ and set A , the ordered iteration $\Sigma_{+,0}$ -algebra $(\Sigma, A)\mathbf{ISR}$ is freely generated by A in the variety of ordered iteration $\Sigma_{+,0}$ -algebras satisfying the (in)equations (1) – (6).*

Proof. We already know that $(\Sigma, A)\mathbf{ISR}$ is an ordered iteration $\Sigma_{+,0}$ -algebra satisfying (1) – (6), cf. Proposition 5.16. Suppose that B is another ordered iteration $\Sigma_{+,0}$ -algebra satisfying the equations (1) – (6). Let φ be a function $A \rightarrow B$. By Theorem 5.7, there is an iteration $\Sigma_{+,0}$ -algebra morphism $\bar{\varphi} : (\Sigma, A)\mathbf{R} \rightarrow B$ extending φ . We show that for all trees S, S' in $(\Sigma, A)\mathbf{R}$, if $S \leq S'$, then $\bar{\varphi}(S) \leq \bar{\varphi}(S')$. By Lemma 7.3 there exist normal representations $(\mu x.t, a)$ and $(\mu y.t', a)$ of dimensions n and m in the parameters $z = (z_1, \dots, z_p)$ with $|(\mu x.t, a)| = S$, $|(\mu y.t', a)| = S'$ and such that for some $\rho : [n] \rightarrow [m]$ with $1\rho = 1$, $t \circ \rho \leq \rho \circ t'$ holds modulo (1), (2), (3), (18), (19). Moreover, by Lemma 7.4, there is a normal representation $(\mu x.s, a)$ of dimension n such that $t \leq s$ modulo the same inequations, and $s \circ \rho = \rho \circ t'$ modulo (1), (2), (3) and (18). By Lemma 7.5, $|(\mu x.s, a)| = S'$. Thus, denoting $b = (b_1, \dots, b_p)$ with $b_i = a_i\bar{\varphi}$, $i \in [p]$, by Lemma 2.3,

$$\begin{aligned} \bar{\varphi}(S) &= \bar{\varphi}((\mu x.t)_{(\Sigma, A)\mathbf{R}}(a)) \\ &= (\mu x.t)_B(b) \\ &\leq (\mu x.s)_B(b) \\ &= \bar{\varphi}((\mu x.s)_{(\Sigma, A)\mathbf{R}}(a)) \\ &= \bar{\varphi}(S'). \end{aligned}$$

It is now easy to complete the proof. For each \equiv -equivalence class $S/\equiv \in (\Sigma, A)\mathbf{R}/\equiv$, define $\varphi^\sharp(S/\equiv) = \bar{\varphi}(S)$. By the above argument, φ^\sharp is a well-defined ordered iteration $\Sigma_{+,0}$ -algebra morphism $(\Sigma, A)\mathbf{ISR} \rightarrow B$ extending φ . The uniqueness of φ^\sharp is obvious. \square

8 An embedding theorem

In this section we show that for each set A , $(\Sigma, A)\mathbf{ISR}$ can be embedded in an ω -continuous additive Σ -algebra. By Proposition 6.4, $(\Sigma, A)\mathbf{ISF}$ is freely generated by A in the variety of all ordered additive Σ -algebras. It is known, cf. [8, 24], that for each signature Δ and each variety \mathcal{V} of ordered Δ -algebras, the ω -continuous algebras in \mathcal{V} and their ω -continuous morphisms form a reflective subcategory of \mathcal{V} . Moreover, for each $B \in \mathcal{V}$, there is an order reflecting morphism from B into the free ω -continuous algebra generated by B . The free ω -continuous algebra on B may be constructed as the completion of B by ω -ideals generated by countable directed subsets (or ω -chains) of B . We show that when Δ is $\Sigma_{+,0}$ and \mathcal{V} is the variety of ordered additive Σ -algebras, the ordered iteration $\Sigma_{+,0}$ -algebra $(\Sigma, A)\mathbf{ISR}$ embeds in the completion of $(\Sigma, A)\mathbf{ISF}$.

LEMMA 8.1 *Suppose that S and T are (Σ, A) -labeled trees such that T is regular and $F \leq T$ holds for all finite trees F with $F \leq S$. Then $S \leq T$.*

Proof. Our claim is obvious when S is finite, so that below we assume that S is infinite. Let F_0, F_1, \dots be an enumeration of finite trees $F_i \subseteq S$ with the property that every vertex of S appears in all but a finite number of the F_i . By our assumption, for each i there is an injective functional simulation $\varphi_i : F_i \rightarrow T$. We show how to construct an injective functional simulation $\varphi : S \rightarrow T$. When v is the root of S , define $v\varphi$ to be the root of T . Suppose now that φ has been defined on all vertices of S of depth not exceeding d such that φ is injective on these vertices and for every vertex u of depth $\leq d$, $T_{u\varphi}$ is isomorphic to $T_{u\varphi_i}$ for all i in an infinite set I of nonnegative integers. Suppose that u has depth d . For simplicity, in the argument below we will assume that each edge leaving u is labeled in Σ_1 . The generalization of the argument is left to the reader. Let v_0, v_1, v_2, \dots be an enumeration of the successors of u . We define the pairwise different successors $v_0\varphi, v_1\varphi, v_2\varphi \dots$ of $u\varphi$ such that for each k there exists an infinite set $I_k \subseteq I$ with the following property:

For all $j < k$ and $i \in I_k$, the edges $u \rightarrow v_j$, $u\varphi \rightarrow v_j\varphi$ and $u\varphi_i \rightarrow v_j\varphi_i$ have the same label and $T_{v_j\varphi}$ is isomorphic to $T_{v_j\varphi_i}$.

When $k = 0$, let $I_0 = I$. Suppose that $k > 0$ and that we have already defined $v_0\varphi, \dots, v_{k-1}\varphi$ and I_k . Since v_k is a vertex of all but a finite number of the trees F_i , $i \in I$, there is an infinite set $I' \subseteq I_k$ such that $v_k\varphi_i$ is defined for all $i \in I'$. Since T is regular, we can select an infinite set $I_{k+1} \subseteq I'$ such that the trees $T_{v_k\varphi_i}$, $i \in I_{k+1}$ are all isomorphic and the edges $u\varphi_i \rightarrow v_k\varphi_i$, $i \in I_{k+1}$ have the same label. Let $i_0 \in I_{k+1}$. The vertices $v_0\varphi_{i_0}, \dots, v_k\varphi_{i_0}$ are all distinct because φ_{i_0} is injective. Since $T_{u\varphi}$ is isomorphic to $T_{u\varphi_{i_0}}$ and for each $j = 0, \dots, k-1$, $T_{v_j\varphi}$ is isomorphic to $T_{v_j\varphi_{i_0}}$, and the edges $u\varphi \rightarrow v_j\varphi$ and $u\varphi_{i_0} \rightarrow v_j\varphi_{i_0}$ have the same label, there exists a successor v' of $u\varphi$ such that $v' \notin \{v_0\varphi, \dots, v_{k-1}\varphi\}$, $T_{v'}$ is isomorphic to $T_{v_k\varphi_0}$ and $u\varphi \rightarrow v'$ and $u\varphi_{i_0} \rightarrow v_k\varphi_{i_0}$ are labeled by the same letter. Define $v_k\varphi = v'$. \square

Let $I(A)$ denote the collection of all ω -ideals of injective simulation equivalence classes of finite (Σ, A) -labeled trees, i.e., sets of the form

$$(U] = \{F \in (\Sigma, A)\mathbf{ISF} : \exists T \in U \ F \leq T\},$$

where U is any countable directed set of finite trees in $(\Sigma, A)\mathbf{ISF}$. In fact, it is easy to see that a set is in $I(A)$ iff it is a countable ideal which is also a directed set with respect to the injective simulation order. Given $\sigma \in \Sigma_m$ and $I_1 = (U_1], \dots, I_m = (U_m]$, we define

$$\sigma(I_1, \dots, I_m) = (\{\sigma(F_1, \dots, F_m) : F_i \in U_i\}).$$

Moreover, we define $0 = \{0\} = (\{0\})$ and

$$I_1 + I_2 = (\{F_1 + F_2 : F_i \in U_i\}),$$

for all $I_1 = (U_1]$ and $I_2 = (U_2]$ in $I(A)$. It can be shown that the operations are well-defined and that $I(A)$, equipped with the subset order is an ω -continuous additive Σ -algebra.

THEOREM 8.2 *For each set A , there exist an ω -continuous additive Σ -algebra B and an injective order reflecting iteration $\Sigma_{+,0}$ -algebra morphism $(\Sigma, A)\mathbf{ISR} \rightarrow B$.*

Proof. Let $B = I(A)$ and consider the map $A \rightarrow B$ taking each letter a into $\{a\} = (\{a\})$. Since B is an ordered iteration $\Sigma_{+,0}$ -algebra satisfying

(1) – (6), by Theorem 7.6 this map extends to an ordered additive iteration $\Sigma_{+,0}$ -algebra morphism h . We need to show that for any two regular trees S, T , if $Sh \subseteq Th$, then $S \leq T$. But it is easy to see that Sh is the set of all finite trees $\leq S$, and similarly for T . Thus, by Lemma 8.1, $S \leq T$. \square

9 Further results

In this section we outline some further results. First we consider decidability and nonfinite axiomatizability.

Suppose that $t \leq t'$ is an inequation between μ -terms t, t' over $\Sigma_{+,0}$, where Σ is a finite signature. Let T and T' denote the corresponding regular trees in $(\Sigma, X)\mathbf{R}$. Thus, T is the value of the function $t_{(\Sigma, X)\mathbf{R}}$ induced by T when each variable x is assigned the corresponding tree having a single edge which is labeled x . By our previous results, the inequation $t \leq t'$ holds in ω -continuous additive Σ -algebras iff $T \leq T'$. This latter property is easily seen to be decidable. Thus we have

PROPOSITION 9.1 *When Σ is finite, it is decidable whether an inequation between μ -terms over Σ holds in all ω -continuous additive Σ -algebras.*

Since each term is finite, the assumption on the finiteness of Σ is not important. It can be shown that the problem is in fact in **NP**.

We have seen that for each Σ , the variety generated by ω -continuous additive Σ -algebras can be axiomatized by an infinite number of equation schemes.

THEOREM 9.2 *Suppose that Σ contains a symbol of positive rank. Then the variety of ordered iteration algebras generated by the ω -continuous additive Σ -algebras cannot be axiomatized by a finite number of equation schemes.*

Proof. It is sufficient to prove this fact in the case that Σ contains a single symbol, say σ , and σ has rank 1.

Let \mathbf{Reg}_a denote the collection of all regular subsets of a^* . We may turn \mathbf{Reg}_a into an ω -continuous additive Σ -algebra by defining

$$\begin{aligned} L_1 + L_2 &= L_1 \cup L_2 \\ 0 &= \emptyset \\ \sigma(L) &= aL = \{au : u \in L\} \end{aligned}$$

and

$$L_1 \leq L_2 \Leftrightarrow L_1 \subseteq L_2,$$

for all $L_1, L_2 \in \mathbf{Reg}_a$.

The result now follows from two observations. First, it is proved in [2, 16] that the variety of ordered iteration algebras generated by \mathbf{Reg}_a cannot be defined by a finite number of equation schemes. (Actually, this is proved in [2, 16] in an equivalent form using the regular operations of set union, concatenation and Kleene star.) Second, it follows from results proved in [34, 19, 16] that the valid inequations of \mathbf{Reg}_a can be axiomatized, relatively to the equations of ω -continuous Σ -algebras, by a finite number of equation schemes. \square

When Σ is an arbitrary signature, define

$$\sigma(L_1, \dots, L_n) = \begin{cases} aL_1 & \text{if } n > 0 \\ \emptyset & \text{otherwise,} \end{cases}$$

for all $\sigma \in \Sigma_n$, $n \geq 0$, and $L_1, \dots, L_n \in \mathbf{Reg}_a$. The same argument shows that when $\Sigma \neq \Sigma_0$, *no* variety of ordered iteration $\Sigma_{+,0}$ -algebras containing the algebra \mathbf{Reg}_a has an axiomatization by a finite number of equation schemes.

It is natural to ask how our results change when the operations induced by the letters in Σ are commutative, so that all equations

$$\sigma(x_1, \dots, x_n) = \sigma(x_{1\pi}, \dots, x_{n\pi}) \tag{20}$$

hold, where $\sigma \in \Sigma_n$ and π is any permutation $[n] \rightarrow [n]$. It can be seen that the variety of ordered preiteration $\Sigma_{+,0}$ -algebras generated by the ω -continuous additive Σ -algebras stisfying equations (20) can be axiomatized

by the inequations that hold in ω -continuous additive Σ -algebras and the equations (20). The free algebras in this variety can be described by injective simulations of the variant of (regular) synchronization trees where each hyperedge has an unordered n -tuple as its target.

References

- [1] L. Aceto, W.J. Fokkink and A. Ingólfssdóttir, A menagerie of nonfinitely based process semantics over BPA*: from ready simulation to completed traces, *Math. Struct. Comput. Sci.*, 8(1998), 193-230.
- [2] L. Aceto, W.J. Fokkink and A. Ingólfssdóttir, On a question of A. Salomaa: The equational theory of regular expressions over a singleton alphabet is not finitely based, *Theoret. Comput. Sci.*, 209(1998), 163-178.
- [3] J.W. De Bakker and D. Scott, A theory of programs, IBM Seminar, Vienna, 1969.
- [4] J. Balcazar, J. Gabarro and M. Santha, Deciding bisimilarity is P-complete, *Formal Aspects of Computing*, 4(1992), 638-648.
- [5] M. Barr and Ch. Wells, *Category Theory for Computing Science*, Prentice-Hall International Series in Computer Science. Prentice-Hall International, New York, 1990.
- [6] H. Bekić, Definable operations in general algebras, and the theory of automata and flowcharts, *Technical Report*, IBM Laboratory, Vienna, 1969.
- [7] D. Benson and J. Tiuryn, Fixed points in free process algebras, *Theoret. Comput. Sci.*, 63(1989), 275-294.
- [8] S.L. Bloom, Varieties of ordered algebras, *J. Comput. System Sci.*, 13(1976), 200-212.
- [9] S.L. Bloom and Z. Ésik, *Iteration Theories*, Springer-Verlag, 1993.

- [10] S.L. Bloom, Z. Ésik and D. Taubner, Iteration theories of synchronization trees, *Information and Computation*, 102(1993), 1–55.
- [11] S.L. Bloom and Z. Ésik, The equational logic of fixed points, *Theoret. Comput. Sci.*, 179(1997), 1–60.
- [12] J.H. Conway, *Regular Algebra and Finite Machines*, Chapman and Hall, London, 1971.
- [13] F. Corradini, R. De Nicola and A. Labella, Tree morphisms and bisimulations, in: Proc. *MFCS'98 Workshop on Concurrency, ENTCS*, 18(1998).
- [14] F. Corradini, R. De Nicola and A. Labella, A finite axiomatization of nondeterministic regular expressions, *Theoret. Inform. Appl.*, 33(1999), 447–465.
- [15] F. Corradini, R. De Nicola and A. Labella, Models of nondeterministic regular expressions. *J. Comput. Sys. Sci.*, 59(1999), 412–449.
- [16] S. Crvenković, I. Dolinka, Z. Ésik, A note on equations for commutative regular languages, *Inform. Process. Lett.*, 70(1999), 265–267.
- [17] Z. Ésik, Completeness of Park induction, *Theoret. Comput. Sci.*, 177(1997), 217–283.
- [18] Z. Ésik and A. Labella, Equational properties of iteration in algebraically complete categories. Mathematical foundations of computer science (Cracow, 1996). *Theoret. Comput. Sci.*, 195(1998), 61–89.
- [19] Z. Ésik, Group axioms for iteration, *Inform. and Comput.*, 148(1999), 131–180.
- [20] Z. Ésik, Axiomatizing the least fixed point operation and binary supremum, in: proc. *CSL 2000*, LNCS 1862, Springer–Verlag, 2000, 302–316.
- [21] W. Fokkink and H. Zantema, Basic process algebra with iteration: Completeness of its equational axioms. *Computer Journal*, 37(1994), 259–267.

- [22] J. Goguen, J. Thatcher, E. Wagner, and J. Wright, Initial algebra semantics and continuous algebras, *J. ACM*, 24(1977), 68–95.
- [23] G. Gratzer, *Universal Algebra*. Second edition, Springer-Verlag, New York-Heidelberg, 1979.
- [24] I. Guessarian, *Algebraic Semantics*, LNCS 99, Springer-Verlag, 1981.
- [25] P.C. Kanellakis and S.A. Smolka, CCS expressions, finite state processes and three problems of equivalence, *Inform. and Comput.*, 86(1990), 43–68.
- [26] D. Kozen, A completeness theorem for Kleene algebras and the algebra of regular events, *Inform. and Comput.*, 110(1994), 366–390.
- [27] D. Krob, Monoides et semi-anneaux complets, *Semigroup Forum*, 36(1987), 323–339,
- [28] D. Krob, Complete systems of B-rational identities, *Theoret. Comput. Sci.*, 89(1991), 207–343.
- [29] E.G. Manes and M.A. Arbib, *Algebraic Approaches to Program Semantics*, Springer-Verlag, 1986.
- [30] R. Milner, A complete inference system for a class of regular behaviours, *J. Comput. Syst. Sci.*, 28(1984), 439–466.
- [31] R. Milner, *Communication and Concurrency*, Prentice-Hall, 1989.
- [32] D.M.R. Park, Fixpoint induction and proofs of program properties, in: *Machine Intelligence 5*, D. Michie and B. Meltzer, Eds., Edinburgh Univ. Press, 1970, 59–78.
- [33] D.M.R. Park, Concurrency and automata on infinite sequences, in: *Proc. GI Conference*, P. Deussen, Ed., LNCS 104, Springer-Verlag, 1981, 167–183.
- [34] V.N. Redko, On algebra of commutative events, *Ukrainian Math. J.*, 16(1964), 185–195 (in Russian).

- [35] A. Salomaa, Two complete axiom systems for the algebra of regular events. *J. Assoc. Comput. Mach.*, 13(1966), 158–169.
- [36] P. Sewell, Nonaxiomatisability of equivalences over finite state processes, *Ann. Pure Appl. Logic*, 90(1997), 163–191.

Recent BRICS Report Series Publications

- RS-00-25 Zoltán Ésik. *Continuous Additive Algebras and Injective Simulations of Synchronization Trees*. September 2000. 41 pp.
- RS-00-24 Claus Brabrand and Michael I. Schwartzbach. *Growing Languages with Metamorphic Syntax Macros*. September 2000.
- RS-00-23 Luca Aceto, Anna Ingólfssdóttir, Mikkel Lykke Pedersen, and Jan Poulsen. *Characteristic Formulae for Timed Automata*. September 2000. 23 pp.
- RS-00-22 Thomas S. Hune and Anders B. Sandholm. *Using Automata in Control Synthesis — A Case Study*. September 2000. 20 pp. Appears in Maibaum, editor, *Fundamental Approaches to Software Engineering: First International Conference, FASE '00 Proceedings*, LNCS 1783, 2000, pages 349–362.
- RS-00-21 M. Oliver Möller and Rajeev Alur. *Heuristics for Hierarchical Partitioning with Application to Model Checking*. August 2000. 30 pp.
- RS-00-20 Luca Aceto, Willem Jan Fokkink, and Anna Ingólfssdóttir. *2-Nested Simulation is not Finitely Equationally Axiomatizable*. August 2000. 13 pp.
- RS-00-19 Vinodchandran N. Variyam. *A Note on $NP \cap coNP/poly$* . August 2000. 7 pp.
- RS-00-18 Federico Crazzolaro and Glynn Winskel. *Language, Semantics, and Methods for Cryptographic Protocols*. August 2000. ii+42 pp.
- RS-00-17 Thomas S. Hune. *Modeling a Language for Embedded Systems in Timed Automata*. August 2000. 26 pp. Earlier version entitled *Modelling a Real-Time Language* appeared in Gnesi and Latella, editors, *Fourth International ERCIM Workshop on Formal Methods for Industrial Critical Systems, FMICS '99 Proceedings of the FLoC Workshop, 1999*, pages 259–282.
- RS-00-16 Jiří Srba. *Complexity of Weak Bisimilarity and Regularity for BPA and BPP*. June 2000. 20 pp. To appear in Aceto and Victor, editors, *Expressiveness in Concurrency: Fifth International Workshop EXPRESS '00 Proceedings*, ENTCS, 2000.