



Basic Research in Computer Science

BRICS RS-97-23 U. Kohlenbach: On the arithmetical content of restricted forms of comprehension

**On the Arithmetical Content of
Restricted Forms of Comprehension,
Choice and General Uniform Boundedness**

Ulrich Kohlenbach

BRICS Report Series

RS-97-23

ISSN 0909-0878

August 1997

**Copyright © 1997, BRICS, Department of Computer Science
University of Aarhus. All rights reserved.**

**Reproduction of all or part of this work
is permitted for educational or research use
on condition that this copyright notice is
included in any copy.**

**See back inner page for a list of recent BRICS Report Series publications.
Copies may be obtained by contacting:**

**BRICS
Department of Computer Science
University of Aarhus
Ny Munkegade, building 540
DK-8000 Aarhus C
Denmark
Telephone: +45 8942 3360
Telefax: +45 8942 3255
Internet: BRICS@brics.dk**

**BRICS publications are in general accessible through the World Wide
Web and anonymous FTP through these URLs:**

`http://www.brics.dk`
`ftp://ftp.brics.dk`
This document in subdirectory RS/97/23/

On the arithmetical content of restricted forms of comprehension, choice and general uniform boundedness

Ulrich Kohlenbach

BRICS*

Department of Computer Science
University of Aarhus
Ny Munkegade, Bldg. 540
DK-8000 Aarhus C, Denmark
kohlenb@brics.dk

August 5, 1997

Abstract

In this paper the numerical strength of fragments of arithmetical comprehension, choice and general uniform boundedness is studied systematically. These principles are investigated relative to base systems \mathcal{T}_n^ω in all finite types which are suited to formalize substantial parts of analysis but nevertheless have provably recursive function(al)s of low growth. We reduce the use of instances of these principles in \mathcal{T}_n^ω -proofs of a large class of formulas to the use of instances of certain arithmetical principles thereby determining faithfully the arithmetical content of the former. This is achieved using the method of elimination of Skolem functions for monotone formulas which was introduced by the author in a previous paper.

As corollaries we obtain new conservation results for fragments of analysis over fragments of arithmetic which strengthen known purely first-order conservation results.

*Basic Research in Computer Science, Centre of the Danish National Research Foundation.

1 Introduction

This paper studies the numerical strength of fragments Γ of arithmetical comprehension, choice and uniform boundedness relative to weak base systems, formulated in the language of all finite types, which are suited to formalize substantial parts of analysis.

In a previous paper ([12]) we have introduced a hierarchy $G_n A^\omega$ of systems where the definable functions correspond to the well-known Grzegorzczuk hierarchy. These systems extended by the schema of full quantifier-free choice

$$AC^{\rho,\tau}\text{-qf} : \forall x^\rho \exists y^\tau A_0(x, y) \rightarrow \exists Y^{\tau(\rho)} \forall x^\rho A_0(x, Yx),$$

$$AC\text{-qf} := \bigcup_{\rho,\tau \in \mathbf{T}} \{AC^{\rho,\tau}\text{-qf}\},$$

where A_0 is a quantifier-free formula,¹ and various non-constructive analytical axioms Δ , having the form

$$\forall x^\rho \exists y \leq_\tau s x \forall z^\delta A_0(x, y, z),$$

including a generalized version of the binary König's lemma, allow to carry out a great deal of classical analysis even for $n = 2, 3$. The axioms Δ and $AC\text{-qf}$ do not contribute to the growth of extractable uniform bounds which in the particular case of $G_2 A^\omega$ are polynomials (see [12],[14] and in particular [10] for more information).

In contrast to this, fragments of arithmetical comprehension and choice as well as generalizations of our principle of uniform Σ_1^0 -boundedness (from [12]) to more complex formulas do contribute significantly to the arithmetic strength of the base systems. In [13] we developed a general method to calibrate faithfully this contribution and applied it to instances of Π_1^0 -comprehension and Π_1^0 -choice. These results were then used in [15] to determine the arithmetical strength of single sequences of instances of the Bolzano-Weierstraß theorem for bounded sequences in \mathbb{R}^d , the Ascoli-lemma and others.

In this paper we give a systematic treatment of the whole arithmetical hierarchy for comprehension, choice and uniform boundedness and determine precisely their arithmetical strength. We also consider much more complex formulas to be proved in these systems than we did in our previous papers.

¹Throughout this paper A_0, B_0, C_0, \dots denote quantifier-free formulas. We allow bounded number quantifiers $\forall x \leq_0 t, \exists x \leq_0 t$ to occur in A_0, B_0, C_0, \dots since they can be expressed in a quantifier-free way using the bounded search-functional μ_b from $G_n A^\omega$. \mathbf{T} denotes the set of all finite types.

In the following let us discuss now some of the difficulties one has to deal with in order to achieve this goal and which indicate already the type of results one can expect. For simplicity we restrict ourselves for the moment to the second-order system $EA^2 + AC^{0,0}\text{-qf}$ instead of $G_nA^\omega + AC\text{-qf} + \Delta$ (which we actually are going to consider below).

EA^2 is an extension of Kalmar-elementary arithmetic (with number quantifiers) EA obtained by adding n -ary function quantifiers (for every $n \geq 1$)² and the schema of explicit definition of functions

$$ED : \exists f \forall \underline{x} (f(\underline{x}) = t[\underline{x}]),$$

where t is a number term of EA^2 and \underline{x} is a tuple of number variables. Furthermore EA^2 contains the schema of quantifier-free induction for all quantifier-free formulas of EA^2 which may contain function parameters. Finally EA^2 contains constants and their defining equations for all elementary recursive functionals of type-level 2.

In EA^2 the schema of quantifier-free induction can be expressed equivalently as a single axiom

$$\text{QF-IA} : \forall f (f(0) = 0 \wedge \forall x (f(x) = 0 \rightarrow f(x') = 0) \rightarrow \forall x (f(x) = 0)).$$

Let us consider furthermore the restriction of arithmetical choice to Π_1^0 - (or equivalently to Σ_2^0 -) formulas of $\mathcal{L}(EA^2)$ which like QF-IA can be expressed as a single second-order axiom $\forall f \Pi_1^0\text{-AC}(f)$, where³

$$\Pi_1^0\text{-AC}(f) : \equiv \forall a^0 (\forall x^0 \exists y^0 \forall z^0 (f(a, x, y, z) = 0) \rightarrow \exists g \forall x, z (f(a, x, gx, z) = 0)).$$

Now by iteration one easily verifies that $EA^2 + \forall f \Pi_1^0\text{-AC}(f)$ proves already full arithmetical choice. So in order to prevent the arithmetical hierarchy of choice principles from collapsing we restrict ourselves to single instances of $\forall f \Pi_1^0\text{-AC}(f)$ which later on are allowed however to depend on the parameters of the theorem to be proved. For the moment we forbid completely the occurrence of function parameters in $\Pi_1^0\text{-AC}$, i.e. we consider the schema

$$\Pi_1^0\text{-AC}^- : \forall x^0 \exists y^0 A(x, y) \rightarrow \exists g \forall x A(x, gx),$$

where $A(x, y)$ is a Π_1^0 -formula without **function parameters**.

²Since coding of finite tuples of numbers is available in EA one can in fact restrict oneself to unary function variables.

³The universal closure with respect to number parameters a^0 is superfluous for $\forall f \Pi_1^0\text{-AC}(f)$ since it can be captured by the universal closure $\forall f$. However below we consider single instances $\Pi_1^0\text{-AC}(\xi)$ of $\forall f \Pi_1^0\text{-AC}(f)$ where it does make a difference.

As a starting point for the introduction into our general program let us consider now the following question:

What arithmetical statements are provable in $EA^2 + AC^{0,0}\text{-qf} + \Pi_1^0\text{-AC}^-$?

A first observation is that $\Pi_1^0\text{-AC}^-$ proves $\Pi_1^0\text{-CA}^-$, i.e.

$$\exists f \forall x (f(x) = 0 \leftrightarrow A(x)),$$

where $A(x)$ is a Π_1^0 -formula without function parameters. Combined with the axiom QF-IA this yields every function parameter-free instance of Σ_1^0 -IA. Hence the first-order system $EA + \Sigma_1^0\text{-IA}$ is a subsystem of $EA^2 + AC^{0,0}\text{-qf} + \Pi_1^0\text{-AC}^-$.

What is the precise relationship between $EA^2 + AC^{0,0}\text{-qf} + \Pi_1^0\text{-AC}^-$ and $EA + \Sigma_1^0\text{-IA}$?

It will turn out that the former theory is conservative over the latter for **some** formulas, including Π_3^0 -sentences, but not for all formulas.

That $EA^2 + AC^{0,0}\text{-qf}$ cannot be conservative over $EA + \Sigma_1^0\text{-IA}$ without some restriction imposed on the formulas follows from the following observation:

By applying the functional $\Phi_{\max} f x := \max_{i \leq x} (f(i))$ to the function g in $\Pi_1^0\text{-AC}^-$ one obtains the corresponding instance of the so-called (bounded) collection principle for Π_1^0 -formulas

$$\Pi_1^0\text{-CP} : \forall x \leq a \exists y A(x, y) \rightarrow \exists z \forall x \leq a \exists y \leq z A(x, y),$$

where $A \in \Pi_1^0$.

So $EA^2 + AC^{0,0}\text{-qf} + \Pi_1^0\text{-AC}^-$ proves every function parameter-free instance of $\Pi_1^0\text{-CP}$, i.e. $EA + \Pi_1^0\text{-CP}$ is a subsystem of $EA^2 + AC^{0,0}\text{-qf} + \Pi_1^0\text{-AC}^-$.

It is well-known (see [18]) that there exists an instance A of $\Pi_1^0\text{-CP}$ which is not provable in $EA + \Sigma_1^0\text{-IA}$. On the other hand $EA + \Pi_1^0\text{-CP}$ is Π_3^0 -conservative over $EA + \Sigma_1^0\text{-IA}$ by a result due to H. Friedman and (implicitly) J.Paris/L.Kirby [17] (see e.g. [7] for details). The universal closure of the instance A of $\Pi_1^0\text{-CP}$ can be shown to be equivalent to a Π_4^0 -sentence in $EA + \Sigma_1^0\text{-IA}$. Hence $EA^2 + AC^{0,0}\text{-qf} + \Pi_1^0\text{-AC}^-$ is not Π_4^0 -conservative over $EA + \Sigma_1^0\text{-IA}$.

Here is another arithmetical use of $\Pi_1^0\text{-AC}^-$ we can make relative to $EA^2 + AC^{0,0}\text{-qf}$:

As mentioned above, $\Pi_1^0\text{-CA}^-$ is a trivial consequence of $\Pi_1^0\text{-AC}^-$ (in the presence of classical logic). Now combining $\Pi_1^0\text{-CA}^-$ with $\text{AC}^{0,0}\text{-qf}$ one can easily prove $\Delta_2^0\text{-CA}^-$ and therefore every function parameter-free instance of $\Delta_2^0\text{-IA}$. Hence $\text{EA} + \Delta_2^0\text{-IA}$ is a subsystem of $\text{EA}^2 + \text{AC}^{0,0}\text{-qf} + \Pi_1^0\text{-AC}^-$ as well even if the functional Φ_{\max} were not included in EA^2 .

So the arithmetical strength of $\Pi_1^0\text{-AC}^-$ depends heavily on the second-order axioms, like QF-IA , $\text{AC}^{0,0}\text{-qf}$ and the characterizing axioms for functionals as Φ_{\max} , which are available in the context in which $\Pi_1^0\text{-AC}^-$ is considered.⁴

As a special corollary of the results of this paper it follows that $\text{EA}^2 + \text{AC}^{0,0}\text{-qf} + \Pi_k^0\text{-AC}^-$ is Π_{k+2}^0 -conservative over $\text{EA} + \Sigma_k^0\text{-IA}$, which implies the result of H. Friedman, J.Paris/L.Kirby. Furthermore we show that $\text{EA}^2 + \text{AC}^{0,0}\text{-qf} + \Pi_k^0\text{-AC}^-$ is conservative over $\text{EA} + \Sigma_k^0\text{-IA}$ w.r.t. monotone formulas of arbitrary complexity. These results are sensitive to small changes of the base system EA^2 : E.g. if we add the primitive recursive functional Φ_{it} defined by

$$\Phi_{it}fg0 := g(0) \quad \Phi_{it}fgx' := f(x, \Phi_{it}fgx)$$

to EA^2 , then the Ackermann-function becomes provably total in $\text{EA}^2 + \Phi_{it} + \text{AC}^{0,0}\text{-qf} + \Pi_1^0\text{-AC}^-$ and the resulting system proves the consistency of $\text{EA} + \Sigma_1^0\text{-IA}$: $\text{EA}^2 + \Phi_{it} + \text{AC}^{0,0}\text{-qf}$ proves the second-order axiom of Σ_1^0 -induction. Combined with $\Pi_1^0\text{-CA}^-$ one obtains every function parameter-free instance of $\Sigma_2^0\text{-IA}$. Hence $\text{EA} + \Sigma_2^0\text{-IA}$ (which is known to prove the totality of the Ackermann-function as well as the consistency of $\text{EA} + \Sigma_1^0\text{-IA}$) is a subsystem of $\text{EA}^2 + \Phi_{it} + \text{AC}^{0,0}\text{-qf} + \Pi_1^0\text{-AC}^-$.

Using a more involved argument one can show that already $\text{EA}^2 + \Phi_{it} + \Pi_1^0\text{-AC}^-$ proves the totality of the Ackermann function (see chapter 12 of [10] for details on this).

So any proof of conservation of systems based on $\Pi_k^0\text{-AC}^-$ over $\Sigma_k^0\text{-IA}$ has to take into account carefully the structure of the functionals of type level 2 which are definable in the given system.

Things become of course even more complicated for the systems $G_nA^\omega + \text{AC-qf} + \Delta$ instead of $\text{EA}^2 + \text{AC}^{0,0}\text{-qf}$ which we are treating in this paper. In particular we show the following result:

⁴Both aspects are not taken into account appropriately in [21] where $\Pi_k^0\text{-CA}^-$ and $\Pi_k^0\text{-AC}^-$ are studied systematically for the first time. As a consequence of this, theorems 5.8,5.13 and corollaries 5.9,5.14 in [21] are not correct (see [11] and in particular chapter 12 of [10] for a thorough investigation of this matter).

Let t, ξ_1, ξ_2 be closed terms of $G_n A^\omega$ and $B := \forall u^1 \forall v \leq_\tau tu B_{ar}(u, v)$ a sentence of $G_n A^\omega$ where $B_{ar}(u, v) \in \Pi_\infty^0$. Then for every $k \in \mathbb{N}$ the following rule holds

$$\left\{ \begin{array}{l} \text{If} \\ G_n A^\omega + \text{AC-qf} + \Delta \vdash \\ \quad \forall u^1 \forall v \leq_\tau tu (\Delta_{k+1}^0\text{-CA}(\xi_1 uv) \wedge \Pi_k^0\text{-AC}(\xi_2 uv) \rightarrow B_{ar}(u, v)) \\ \text{then} \\ G_n A^\omega + \Sigma_k^0\text{-IA} + \tilde{\Delta} + \text{Mon}(B) \vdash \forall u^1 \forall v \leq_\tau tu B_{ar}(u, v), \end{array} \right.$$

where $\text{Mon}(B)$ expresses that B is monotone in the sense of definition 2.3 below and

$$\tilde{\Delta} := \{ \exists Y^{\rho\delta} \leq s \forall x^\delta, z^\gamma A_0(x, Yx, z) : \forall x^\delta \exists y \leq_\rho sx \forall z^\gamma A_0(x, y, z) \in \Delta \}.$$

(Δ and consequently $\tilde{\Delta}$ may be empty).

If $B_{ar} \in \Pi_{k+2}^0$, then the monotonicity assumption $\text{Mon}(B)$ can be avoided, i.e. the conclusion can be strengthened to

$$G_n A^\omega + \Sigma_k^0\text{-IA} + \tilde{\Delta} \vdash \forall u^1 \forall v \leq_\tau tu B_{ar}(u, v).$$

These results will be used also to prove new conservation results for $\text{EA} + \Pi_k^0\text{-CP}$ over $\text{EA} + \Sigma_k^0\text{-IA}$ which strengthen the Friedman-Paris-Kirby result.⁵

Finally we consider generalizations $\Pi_k^0\text{-UB}^-$ of the principle of uniform Σ_1^0 -boundedness $\Sigma_1^0\text{-UB}^-$ which was studied in [12].⁶ In [14] we showed that $\Sigma_1^0\text{-UB}^-$ proves already relative to $G_2 A^\omega + \text{AC-qf}$ many important analytical theorems (like Dini's theorem, the attainment of the maximum for $f \in C([0, 1]^d, \mathbb{R})$, the sequential Heine-Borel property for $[0, 1]^d$, the existence of an inverse function for every strictly monotone function $f \in C[0, 1]$ and others) but does not contribute to the growth of extractable bounds, thereby guaranteeing the extractability of polynomial bounds when applied in the context of $G_2 A^\omega + \text{AC-qf}$.

⁵A proof-theoretic treatment of the Friedman-Paris-Kirby result was first given in [21]. However the proof in [21] contains a serious gap. See [1] for a correction of Sieg's proof. Another proof-theoretic treatment can be found in [3].

⁶Whereas we generally use the superscript ' $-$ ' to denote the restriction S^- of a schema S to function parameter-free instances of S , this superscript has a different meaning in the context of principles of uniform boundedness. Although this might be troublesome we wish to stick to the notation for these principles from [12] where they were introduced.

Whereas the straightforward generalization of $\Sigma_1^0\text{-UB}^-$ to Π_k^0 -formulas is inconsistent with G_2A^ω already for $k = 1$, our restricted version $\Pi_k^0\text{-UB}^- \upharpoonright$ (introduced in the present paper) is consistent. In [15] we implicitly used (a special case of) $\Pi_1^0\text{-UB}^- \upharpoonright$ to prove the Bolzano-Weierstraß principle and the Ascoli-lemma and it were these proofs which were used to calibrate faithfully the arithmetical strength of these principles.

One of the results on $\Pi_k^0\text{-UB}^- \upharpoonright$ to be proved in the present paper is that we may strengthen the assumption of the rule stated above by adding $\Pi_k^0\text{-UB}^- \upharpoonright(\xi_3 uv)$ to $\Delta_{k+1}^0\text{-CA}(\xi_1 uv) \wedge \Pi_k^0\text{-AC}(\xi_2 uv)$.

2 Monotone formulas and their Skolem normal forms

In this section we review some of the proof-theoretic tools from [13] on which the present paper is based and also recall some of the basic concepts and definitions from [12].

The set \mathbf{T} of all finite types is defined inductively by

$$(i) 0 \in \mathbf{T} \text{ and } (ii) \rho, \tau \in \mathbf{T} \Rightarrow \tau(\rho) \in \mathbf{T}.$$

Terms which denote a natural number have type 0. Elements of type $\tau(\rho)$ are functions which map objects of type ρ to objects of type τ .

The set $\mathbf{P} \subset \mathbf{T}$ of pure types is defined by

$$(i) 0 \in \mathbf{P} \text{ and } (ii) \rho \in \mathbf{P} \Rightarrow 0(\rho) \in \mathbf{P}.$$

Brackets whose occurrences are uniquely determined are often omitted, e.g. we write $0(00)$ instead of $0(0(0))$. Furthermore we write for short $\tau\rho_k \dots \rho_1$ instead of $\tau(\rho_k) \dots (\rho_1)$. Pure types can be represented by natural numbers: $0(n) := n + 1$. The types $0, 00, 0(00), 0(0(00)) \dots$ are so represented by $0, 1, 2, 3 \dots$. For arbitrary types $\rho \in \mathbf{T}$ the degree of ρ (for short $\text{deg}(\rho)$) is defined by $\text{deg}(0) := 0$ and $\text{deg}(\tau(\rho)) := \max(\text{deg}(\tau), \text{deg}(\rho) + 1)$. For pure types the degree is just the number which represents this type.

Description of the theories $(\mathbf{E})\text{-G}_nA^\omega$

Our theories \mathcal{T}^ω used in this paper are based on many-sorted classical logic formulated in the language of all finite types plus the combinators $\Pi_{\rho,\tau}, \Sigma_{\delta,\rho,\tau}$ which allow the definition of λ -abstraction. \mathcal{T}_i^ω denotes the intuitionistic variant of \mathcal{T}^ω .

The systems G_nA^ω (for all $n \geq 1$) are introduced in [12] to which we refer for details. G_nA^ω has as primitive relations $=_0, \leq_0$ for objects of type 0, the

constant 0^0 , functions \min_0, \max_0, S^{00} (successor), A_0, \dots, A_n , where A_i is the i -th branch of the Ackermann function (i.e. $A_0(x, y) = y', A_1(x, y) = x + y, A_2(x, y) = x \cdot y, A_3(x, y) = x^y, \dots$), functionals of type level 2: Φ_1, \dots, Φ_n , where $\Phi_1 f x = \max_0(f 0, \dots, f x)$ and Φ_i is the iteration of A_{i-1} on the f -values for $i \geq 2$, i.e. $\Phi_2 f x = \sum_{i=0}^x f i, \Phi_3 f x = \prod_{i=0}^x f i, \dots$. We also have a bounded search functional μ_b and bounded predicative recursion provided by recursor constants \tilde{R}_ρ (where ‘predicative’ means that recursion is possible only at the type-0-level as in the case of the (unbounded) Kleene-Feferman recursors \hat{R}_ρ). Moreover $G_n A^\omega$ contains a quantifier-free rule of extensionality QF-ER.

In addition to the defining axioms for the constants of our theories all true sentences having the form $\forall x^\rho A_0(x)$, where A_0 is quantifier-free and $\text{deg}(\rho) \leq 2$, are added as axioms. By ‘true’ we refer to the full set-theoretic model \mathcal{S}^ω . In given proofs however only very special universal axioms will be used which can be proved in suitable extensions of our theories. Nevertheless we include them all as axioms in order to emphasize that (proofs of) universal sentences do not contribute to the growth of extractable bounds. In particular this covers all instances of the schema of quantifier-free induction (The main results in this paper are also valid for the variant of $G_n A_i^\omega$ where the universal axioms are replaced by the schema of quantifier-free induction). The restriction $\text{deg}(\rho) \leq 2$ has a technical reason discussed in [12].

$$G_\infty A^\omega := \bigcup_{n \in \mathbb{N}} G_n A^\omega.$$

PA^ω, PA_i^ω are the extensions of $G_n A^\omega, G_n A_i^\omega$ obtained by the addition of the schema of full induction and all (impredicative) primitive recursive functionals in the sense of [5].

$E\text{-}\mathcal{T}_{(i)}^\omega$ denotes the theory which results from $\mathcal{T}_{(i)}^\omega$ when the quantifier-free rule of extensionality is replaced by the axioms of extensionality (E)

$$\forall x^\rho, y^\rho, z^{\tau\rho} (x =_\rho y \rightarrow z x =_\tau z y)$$

for all finite types ($x =_\rho y$ is defined as $\forall z_1^{\rho_1}, \dots, z_k^{\rho_k} (x z_1 \dots z_k =_0 y z_1 \dots z_k)$ where $\rho = 0\rho_k \dots \rho_1$).

$G_n R^\omega$ and T denote the sets of all closed terms of (E)- $G_n A_{(i)}^\omega$ and (E)- $PA_{(i)}^\omega$. T_k is the subset of all closed terms of T which contain the Gödel-recursors R_ρ for ρ of type level $\leq k$ only.

Definition 2.1 *Between functionals of type ρ we define relations \leq_ρ (‘less*

or equal') and $s\text{-maj}_\rho$ ('strongly majorizes') by induction on the type:

$$\begin{cases} x_1 \leq_0 x_2 \equiv (x_1 \leq_0 x_2), \\ x_1 \leq_{\tau\rho} x_2 \equiv \forall y^\rho (x_1 y \leq_\tau x_2 y); \end{cases}$$

$$\begin{cases} x^* s\text{-maj}_0 x \equiv x^* \geq_0 x, \\ x^* s\text{-maj}_{\tau\rho} x \equiv \forall y^{*\rho}, y^\rho (y^* s\text{-maj}_\rho y \rightarrow x^* y^* s\text{-maj}_\tau x^* y, xy). \end{cases}$$

Remark 2.2 ' $s\text{-maj}$ ' is a variant of W.A. Howard's relation ' maj ' from [6] which is due to [2]. For more details see [8].

Let $A(\underline{a})$ be a formula of $G_n A^\omega$ (\underline{a} are all free variables of A) and $\exists \underline{x} \forall \underline{y} A_D(\underline{x}, \underline{y}, \underline{a})$ its Gödel functional interpretation (see e.g. [24] for details on Gödel's functional interpretation). We say that a tuple of closed terms \underline{t} realizes the **monotone** functional interpretation of $A(\underline{a})$ if⁷

$$(*) \exists \underline{x} (\underline{t} \text{ s-maj } \underline{x} \wedge \forall \underline{a}, \underline{y} A_D(\underline{x} \underline{a}, \underline{y}, \underline{a}))$$

(Monotone functional interpretation which directly extracts a tuple \underline{t} satisfying $(*)$ from a proof of $A(\underline{a})$ was introduced in [9]. See also [12] for details.)

Definition 2.3 ([13]) Let $A \in \mathcal{L}(G_n A^\omega)$ be a formula having the form

$$A \equiv \forall u^1 \forall v \leq_\tau t u \exists y_1^0 \forall x_1^0 \dots \exists y_k^0 \forall x_k^0 \exists w^\gamma A_0(u, v, y_1, x_1, \dots, y_k, x_k, w),$$

where A_0 is quantifier-free and contains only $u, v, \underline{y}, \underline{x}, w$ free, $t \in G_n R^\omega$ and τ, γ are arbitrary finite types.

1) A is called (arithmetically) **monotone** if

$$\text{Mon}(A) := \begin{cases} \forall u^1 \forall v \leq_\tau t u \forall x_1, \tilde{x}_1, \dots, x_k, \tilde{x}_k, y_1, \tilde{y}_1, \dots, y_k, \tilde{y}_k \\ \left(\bigwedge_{i=1}^k (\tilde{x}_i \leq_0 x_i \wedge \tilde{y}_i \geq_0 y_i) \wedge \exists w^\gamma A_0(u, v, y_1, x_1, \dots, y_k, x_k, w) \right) \\ \rightarrow \exists w^\gamma A_0(u, v, \tilde{y}_1, \tilde{x}_1, \dots, \tilde{y}_k, \tilde{x}_k, w). \end{cases}$$

⁷Here $\underline{t} \text{ s-maj } \underline{x}$ means $\bigwedge_i (t_i \text{ s-maj } x_i)$.

2) The **Herbrand normal form** A^H of A is defined to be

$$A^H := \forall u^1 \forall v \leq_\tau tu \forall h_1^{\rho_1}, \dots, h_k^{\rho_k} \exists y_1^0, \dots, y_k^0, w^\gamma$$

$$\underbrace{A_0(u, v, y_1, h_1 y_1, \dots, y_k, h_k y_1 \dots y_k, w)}_{A_0^H :=}$$

where $\rho_i = 0 \underbrace{(0) \dots (0)}_i$.

Remark 2.4 In definition 2.3 (and theorems 2.5, 2.7 below) one may also have tuples ' $\exists \underline{w}$ ' instead of ' $\exists w^\gamma$ ' in A where $\underline{w} = w_1^{\gamma_1}, \dots, w_l^{\gamma_l}$ and γ_i is arbitrary. Also instead of ' $\forall u^1$ ' we may have ' $\forall \underline{u}$ ' where $\underline{u} = u_1^{\rho_1}, \dots, u_q^{\rho_q}$ with $\deg(\rho_i) \leq 1$ for $1 \leq i \leq q$. In particular we can consider an innermost existential number quantifier $\exists y_{k+1}^0$ as part of $\exists \underline{w}$ and an outermost universal number quantifier $\forall x_0^0$ as part of $\forall \underline{u}$. So for $\forall x_0^0$ and $\exists y_{k+1}^0$ no monotonicity is required in definition 2.3.1).

Theorem 2.5 ([13]) Let $n \geq 1$ and $\Psi_1, \dots, \Psi_k \in G_n R^\omega$. Then

$$G_n A^\omega + \text{Mon}(A) \vdash \forall u^1 \forall v \leq_\tau tu \forall h_1, \dots, h_k \left(\bigwedge_{i=1}^k (h_i \text{ monotone}) \right. \\ \left. \rightarrow \exists y_1 \leq_0 \Psi_1 u \underline{h} \dots \exists y_k \leq_0 \Psi_k u \underline{h} \exists w^\gamma A_0^H \right) \rightarrow A,$$

where $(h_i \text{ monotone}) := \forall x_1, \dots, x_i, y_1, \dots, y_i \left(\bigwedge_{j=1}^i (x_j \geq_0 y_j) \rightarrow h_i \underline{x} \geq_0 h_i \underline{y} \right)$.

Definition 2.6 (Bounded choice) $b\text{-}AC := \bigcup_{\delta, \rho \in \mathbf{T}} \{(b\text{-}AC^{\delta, \rho})\}$ denotes the schema of bounded choice

$$(b\text{-}AC^{\delta, \rho}) : \forall Z^{\rho\delta} (\forall x^\delta \exists y \leq_\rho Zx \ A(x, y, Z) \rightarrow \exists Y \leq_{\rho\delta} Z \forall x A(x, Yx, Z)).$$

In general $G_n A^\omega \vdash A^H$ does not imply $G_n A^\omega \vdash A$ (see [13] for a detailed discussion of this phenomenon), which is in contrast to the first-order case where the derivability of A^H follows from that of A by Herbrand's theorem (see [20]). If however A is monotone then this rule is valid also for $G_n A^\omega$ (but for very different reasons):

Theorem 2.7 ([13]) Let A be as in thm.2.5 and Δ be a set of sentences $\forall x^\delta \exists y \leq_\rho s x \forall z^\eta G_0(x, y, z)$ where s is a closed term of $G_n A^\omega$ and G_0 a

quantifier-free formula, and let A' denote the negative translation⁸ of A . Then the following rule holds:

$$\left\{ \begin{array}{l} G_n A^\omega + AC\text{-}qf + \Delta \vdash A^H \wedge \text{Mon}(A) \Rightarrow \\ G_n A^\omega + \tilde{\Delta} \vdash A, \text{ and by monotone functional interpretation one can} \\ \text{extract a tuple } \underline{\Psi} \in G_n R^\omega \text{ such that} \\ G_n A_i^\omega + \tilde{\Delta} \vdash \underline{\Psi} \text{ satisfies the monotone functional interpretation of } A', \end{array} \right.$$

where $\tilde{\Delta} := \{\exists Y \leq_{\rho\delta} s \forall x^\delta, z^\eta G_0(x, Yx, z) : \forall x^\delta \exists y \leq_\rho sx \forall z^\eta G_0(x, y, z) \in \Delta\}$. (In particular the second conclusion can be proved in $G_n A_i^\omega + \Delta + b\text{-}AC$).

The weakened conclusion $G_n A^\omega + \tilde{\Delta} + \text{Mon}(A) \vdash A$ follows already from $G_n A^\omega + AC\text{-}qf + \Delta \vdash A^H$.⁹

3 Making arithmetical comprehension monotone

In this section we consider the arithmetical content of instances $\Pi_k^0\text{-CA}(\xi uv)$ of $\Pi_k^0\text{-CA}$ which are used in given proofs of sentences $\forall u^1 \forall v \leq_\tau tu B_{ar}(u, v)$ as discussed in the introduction.

Definition 3.1

$$\Pi_k^0\text{-CA}(f) := \exists g^1 \forall x^0 (gx =_0 0 \leftrightarrow \forall u_1^0 \exists u_2^0 \dots \exists^{(d)} u_k^0 (f(x, \underline{u}) =_0 0)).^{10}$$

Remark 3.2 *There is no need here to incorporate closure under number parameters in the definition of $\Pi_k^0\text{-CA}(f)$, i.e. by defining*

$$\Pi_k^0\text{-CA}(f) := \forall l^0 \exists g^1 \forall x^0 (gx =_0 0 \leftrightarrow \forall u_1^0 \exists u_2^0 \dots \exists^{(d)} u_k^0 (f(l, x, \underline{u}) =_0 0)),$$

since the latter can be reduced to the former (relative to $G_n A^\omega$ for $n \geq 2$) by coding l, x together and applying comprehension without number parameters to this pair.

⁸Here we can use Gödel's [4] translation or any other of the various negative translations. For a systematical treatment of negative translations see [16].

⁹This last assertion is not stated in the formulation of the theorem in [13] but does follow immediately from its proof.

¹⁰Whether one has here ' $\exists u_k^0$ ' or ' $\forall u_k^0$ ' depends of course on whether k is even or odd.

In order to be able to apply the method of elimination of Skolem functions for monotone formulas from section 2 we follow this strategy:

Construct an arithmetical principle $A_{ar}(f)$ such that for suitable $\xi_1, \xi_2 \in G_n R^\omega$:

- 1) $G_n A^\omega \vdash Mon(\forall f A_{ar}(f))$,
- 2) $G_n A^\omega \vdash \forall f (A_{ar}^S(\xi_1 f) \rightarrow \Pi_k^0\text{-CA}(f))$ and
- 3) $G_n A^\omega \vdash \forall f (\Pi_k^0\text{-CA}(\xi_2 f) \rightarrow A_{ar}(f))$.

Because of 2) the use of $\Pi_k^0\text{-CA}(\xi uv)$ in a given proof of a monotone sentence $\forall u^1 \forall v \leq_\tau tu B_{ar}(u, v)$ can be reduced to the use of $A_{ar}^S(\xi' uv)$ (where $\xi' uv := \xi_1(\xi uv)$) which in turn (by 1) and theorem 2.7) can be reduced to the use of $A_{ar}(\xi' uv)$. Because of 3) nothing is lost by this reduction.

It will turn out that the correct principle $A_{ar}(f)$ is a ‘monotone version’ $\Pi_k^0\text{-TND}^{mon}(f)$ of the tertium-non-datur principle for Π_k^0 -formulas.

Definition 3.3 *In the following $m := \frac{k}{2}$ if k is even (resp. $m := \frac{k-1}{2}$ if k is odd).*

- 1) *The Π_k^0 -tertium-non-datur axiom is given by the following formula (where f is a function variable of appropriate type)¹¹*

$$\begin{aligned} \underline{\Pi_k^0\text{-TND}}(f) &::= \\ &\left\{ \begin{array}{l} \forall x^0 (\forall y_1^0 \exists z_1^0 \dots \forall y_m^0 \exists z_m^0 (\forall y_{m+1}^0) (f(x, y_1, z_1, \dots, y_m, z_m, (y_{m+1})) =_0 0) \\ \vee \exists u_1^0 \forall v_1^0 \dots \exists u_m^0 \forall v_m^0 (\exists u_{m+1}^0) (f(x, u_1, v_1, \dots, u_m, v_m, (u_{m+1})) \neq 0), \end{array} \right. \end{aligned}$$

- 2) *We also need the following prenex normal form of $\Pi_k^0\text{-TND}(f)$:*

$$\begin{aligned} \underline{\Pi_k^0\text{-TND}}(f)^{pr} &::= \\ &\left\{ \begin{array}{l} \forall x^0 \exists u_1^0 \forall y_1^0 \exists z_1^0 \forall v_1^0 \dots \exists u_m^0 \forall y_m^0 \exists z_m^0 \forall v_m^0 (\exists u_{m+1}^0 \forall y_{m+1}^0) \\ \quad (f(x, y_1, z_1, \dots, y_m, z_m, (y_{m+1})) =_0 0 \vee \\ \quad f(x, u_1, v_1, \dots, u_m, v_m, (u_{m+1})) \neq 0), \end{array} \right. \end{aligned}$$

¹¹Here and in the following the quantifiers $\forall y_{m+1}^0, \exists u_{m+1}^0$ are only present if k is odd.

3) The Skolem normal form of $\Pi_k^0\text{-TND}(f)^{pr}$ is given by

$$\underbrace{(\Pi_k^0\text{-TND}(f)^{pr})^S}_{\text{S}} := \left\{ \begin{array}{l} \exists h_1, \dots, h_m, (h_{m+1}), g_1, \dots, g_m \forall x^0, y_1^0, v_1^0, \dots, y_m^0, v_m^0, (y_{m+1}) \\ (f(x, y_1, g_1(x, y_1), \dots, y_m, g_m(x, y_1, \dots, y_m, v_1, \dots, v_{m-1}), (y_{m+1})) =_0 0 \vee \\ f(x, h_1 x, v_1, \dots, h_m(x, y_1, \dots, y_{m-1}, v_1, \dots, v_{m-1}), v_m, \\ (h_{m+1}(x, y_1, \dots, y_m, v_1, \dots, v_m))) \neq 0). \end{array} \right.$$

Remark 3.4 For $n \geq 2$ we have coding of finite tuples (of fixed length) available in $G_n A^\omega$. Hence quantifier-blocks can be contracted to a single quantifier. Since in all of our results we assume that (at least) $n \geq 2$, it is no restriction in the definition above to consider only single quantifiers.

Lemma 3.5 For every $k \in \mathbb{N}$ the following implication can be proved in $G_1 A^\omega$:

$$\forall f((\Pi_k^0\text{-TND}(f)^{pr})^S \rightarrow \Pi_k^0\text{-CA}(f)).$$

Proof:

For notational simplicity we confine ourselves to the case $k = 4$ which well shows the general pattern of the proof for arbitrary k :

$(\Pi_4^0\text{-TND}(f)^{pr})^S$ yields the existence of functions g_1, g_2, h_1, h_2 such that

$$(1) \left\{ \begin{array}{l} \forall x, y_1, v_1, y_2 (f(x, y_1, g_1(x, y_1), y_2, g_2(x, y_1, y_2, v_1)) = 0 \\ \vee \forall v_2 (f(x, h_1 x, v_1, h_2(x, y_1, v_1), v_2) \neq 0)). \end{array} \right.$$

(1) in turn yields

$$(2) \left\{ \begin{array}{l} \forall x, y_1, v_1 (\forall y_2 \exists z_2 f(x, y_1, g_1(x, y_1), y_2, z_2) = 0 \\ \vee \forall v_2 (f(x, h_1 x, v_1, h_2(x, y_1, v_1), v_2) \neq 0)), \end{array} \right.$$

$$(3) \left\{ \begin{array}{l} \forall x, y_1, v_1 (\forall y_2 \exists z_2 f(x, y_1, g_1(x, y_1), y_2, z_2) = 0 \\ \vee \exists u_2 \forall v_2 (f(x, h_1 x, v_1, u_2, v_2) \neq 0)), \end{array} \right.$$

$$(4) \left\{ \begin{array}{l} \forall x, y_1 (\forall y_2 \exists z_2 f(x, y_1, g_1(x, y_1), y_2, z_2) = 0 \\ \vee \forall v_1 \exists u_2 \forall v_2 (f(x, h_1 x, v_1, u_2, v_2) \neq 0)), \end{array} \right.$$

$$(5) \begin{cases} \forall x, y_1 (\exists z_1 \forall y_2 \exists z_2 f(x, y_1, z_1, y_2, z_2) = 0 \\ \quad \forall \forall v_1 \exists u_2 \forall v_2 (f(x, h_1 x, v_1, u_2, v_2) \neq 0)) \end{cases}$$

and finally

$$(6) \forall x (\forall y_1 \exists z_1 \forall y_2 \exists z_2 f(x, y_1, z_1, y_2, z_2) = 0 \vee \forall v_1 \exists u_2 \forall v_2 (f(x, h_1 x, v_1, u_2, v_2) \neq 0)).$$

(1) applied to $y_1 := h_1 x, v_1 := g_1(x, h_1 x), y_2 := h_2(x, h_1 x, g_1(x, h_1 x))$ gives

(*) $:=$

$$\forall x^0 \left(f(x, h_1 x, g_1(x, h_1 x), h_2(x, h_1 x, g_1(x, h_1 x))), g_2(x, h_1 x, h_2(x, h_1 x, g_1(x, h_1 x))), \right. \\ \left. g_1(x, h_1 x)) \right) = 0 \vee \forall v_2 (f(x, h_1 x, g_1(x, h_1 x), h_2(x, h_1 x, g_1(x, h_1 x))), v_2) \neq 0).$$

We now show (+) $:=$

$$\forall x^0 \left(f(x, h_1 x, g_1(x, h_1 x), h_2(x, h_1 x, g_1(x, h_1 x))), g_2(x, h_1 x, h_2(x, h_1 x, g_1(x, h_1 x))), \right. \\ \left. g_1(x, h_1 x)) \right) = 0 \leftrightarrow \forall y_1 \exists z_1 \forall y_2 \exists z_2 (f(x, y_1, z_1, y_2, z_2) = 0).$$

(+) yields the claim of the lemma with

$$gx := \Phi x h_1 h_2 g_1 g_2 := \begin{cases} f(x, h_1 x, g_1(x, h_1 x), h_2(x, h_1 x, g_1(x, h_1 x))), \\ g_2(x, h_1 x, h_2(x, h_1 x, g_1(x, h_1 x))), g_1(x, h_1 x)). \end{cases}$$

Proof of (+):

' \rightarrow ': $\Phi x f h_1 h_2 g_1 g_2 = 0$ implies

$$\neg \forall v_2 (f(x, h_1 x, g_1(x, h_1 x), h_2(x, h_1 x, g_1(x, h_1 x))), v_2) \neq 0).$$

Hence by (2) (putting $y_1 := h_1 x, v_1 := g_1(x, h_1 x)$)

$$\forall y_2 \exists z_2 (f(x, h_1 x, g_1(x, h_1 x), y_2, z_2) = 0)$$

and therefore

$$\exists z_1 \forall y_2 \exists z_2 (f(x, h_1 x, z_1, y_2, z_2) = 0),$$

i.e.

$$\neg \forall v_1 \exists u_2 \forall v_2 (f(x, h_1 x, v_1, u_2, v_2) \neq 0).$$

By (6) this implies

$$\forall y_1 \exists z_1 \forall y_2 \exists z_2 (f(x, y_1, z_1, y_2, z_2) = 0).$$

' \leftarrow ': $\Phi x f h_1 h_2 g_1 g_2 \neq 0$ implies by (*)

$$\forall v_2 (f(x, h_1 x, g_1(x, h_1 x), h_2(x, h_1 x, g_1(x, h_1 x))), v_2) \neq 0)$$

and therefore

$$\exists u_2 \forall v_2 (f(x, h_1 x, g_1(x, h_1 x), u_2, v_2) \neq 0),$$

i.e.

$$\neg \forall y_2 \exists z_2 (f(x, h_1 x, g_1(x, h_1 x), y_2, z_2) = 0).$$

By (4) this yields (putting $y_1 := h_1 x$)

$$\forall v_1 \exists u_2 \forall v_2 (f(x, h_1 x, v_1, u_2, v_2) \neq 0)$$

and therefore

$$\exists u_1 \forall v_1 \exists u_2 \forall v_2 (f(x, u_1, v_1, u_2, v_2) \neq 0),$$

which concludes the proof of (+) and hence of the lemma.

Definition 3.6 For a Π_k^0 -formula

$A(\underline{a}) \equiv \forall x_1^0 \exists x_2^0 \dots \exists^{(d)} x_k^0 A_0(\underline{a}, x_1, x_2, \dots, x_k)$ of $G_n A^\omega$ (where \underline{a} are all free variables of A which may have arbitrary type) we define

$$\tilde{A}(\underline{a}) :=$$

$$\forall x_1^0 \exists x_2^0 \dots \exists^{(d)} x_k^0 \forall \tilde{x}_1 \leq x_1 \exists \tilde{x}_2 \leq x_2 \dots \exists^{(d)} \tilde{x}_k \leq x_k A_0(\underline{a}, \tilde{x}_1, \tilde{x}_2, \dots, \tilde{x}_k).$$

In the following we need a variant Mon^* of Mon where monotonicity is required for **all** number quantifiers (compare this with remark 2.4):

Definition 3.7 Let $A(\underline{a}) \equiv \forall x_1^0 \exists y_1^0 \dots \forall x_k^0 \exists y_k^0 A_0(\underline{a}, x_1, y_1, \dots, x_k, y_k)$.¹²

Then

$$\begin{aligned} Mon^*(A(\underline{a})) &:= \forall x_1, \tilde{x}_1, y_1, \tilde{y}_1, \dots, x_k, \tilde{x}_k, y_k, \tilde{y}_k \left(\bigwedge_{i=1}^k (\tilde{x}_i \leq_0 x_i \wedge \tilde{y}_i \geq_0 y_i) \rightarrow \right. \\ &\quad \left. (A_0(\underline{a}, x_1, y_1, \dots, x_k, y_k) \rightarrow A_0(\underline{a}, \tilde{x}_1, \tilde{y}_1, \dots, \tilde{x}_k, \tilde{y}_k)) \right). \end{aligned}$$

Lemma 3.8 For $\tilde{A}(\underline{a})$ as in the previous definition we have

$$G_n A^\omega \vdash Mon^*(\tilde{A}(\underline{a})).$$

Proof: Trivial.

The Π_k^0 -collection principle is the schema

$$\underline{\Pi_k^0\text{-CP}} : \forall x \leq_0 a \exists y^0 A(x, y) \rightarrow \exists z^0 \forall x \leq_0 a \exists y \leq_0 z A(x, y),$$

for all Π_k^0 -formulas $A(x, y)$.

¹²Here the quantifiers $\forall x_1^0$ and $\exists y_k^0$ may be empty ('dummy') quantifiers.

Convention 3.9 In Π_k^0 -CP (and other axiom schemas which we will consider below) $A(x, y)$ may contain arbitrary parameters (besides x, y) of the language we consider. E.g. if we write $G_n A^\omega + \Pi_k^0$ -CP then instances of Π_k^0 -CP may contain parameters of arbitrary type. In $EA + \Pi_k^0$ -CP however (where EA denotes first-order elementary recursive arithmetic) instances of Π_k^0 -CP of course contain only number parameters.

Π_k^0 -CP is equivalent over many systems (e.g. $G_n A^\omega$ for $n \geq 3$) to the axiom schema of finite choice for Π_k^0 -formulas

$$\underline{\Pi_k^0\text{-FAC}} : \forall x \leq_0 a \exists y^0 A(x, y) \rightarrow \exists z^0 \forall x \leq_0 a A(x, (z)_x),$$

for all Π_k^0 -formulas $A(x, y)$ (with the convention stated above).

In the presence of function variables as in $G_n A^\omega$ the schema Π_k^0 -CP can be expressed as a single second-order axiom $\forall f \Pi_k^0\text{-CP}(f)$, where

$$\underline{\Pi_k^0\text{-CP}(f)} := \begin{cases} \forall l^0, a^0 (\forall x \leq_0 a \exists y^0 \forall u_1^0 \exists u_2^0 \dots \exists^{(d)} u_k^0 (f(l, a, x, y, \underline{u}) =_0 0) \\ \rightarrow \exists z^0 \forall x \leq_0 a \exists y \leq_0 z \forall u_1^0 \exists u_2^0 \dots \exists^{(d)} u_k^0 (f(l, a, x, y, \underline{u}) =_0 0) \end{cases}.$$

By incorporating the universal closure w.r.t. to **arithmetical** parameters $\forall l^0, a^0$ in $\Pi_k^0\text{-CP}(f)$, we achieve that the universal closure of every instance of Π_k^0 -CP which contains only number parameters can be written as a sentence $\Pi_k^0\text{-CP}(\xi)$ in $G_n A^\omega$ where ξ is a closed term (essentially the characteristic function of the quantifier-free matrix of the Π_k^0 -formula $A(x, y)$) which will be of importance below.

The same is true for the principle of Σ_k^0 -induction $\Sigma_k^0\text{-IA}(f)$ which we need below:

$$\underline{\Sigma_k^0\text{-IA}(f)} := \begin{cases} \forall l^0 (\exists u_1^0 \forall u_2^0 \dots \forall^{(d)} u_k^0 (f(l, 0, \underline{u}) =_0 0) \wedge \\ \forall x^0 (\exists u_1^0 \forall u_2^0 \dots \forall^{(d)} u_k^0 (f(l, x, \underline{u}) =_0 0) \rightarrow \\ \exists u_1^0 \forall u_2^0 \dots \forall^{(d)} u_k^0 (f(l, x', \underline{u}) =_0 0)) \\ \rightarrow \forall x^0 \exists u_1^0 \forall u_2^0 \dots \forall^{(d)} u_k^0 (f(l, x, \underline{u}) =_0 0) \end{cases}.$$

Lemma 3.10 Let $A(\underline{a}), \tilde{A}(\underline{a})$ be as in definition 3.6. Then for suitable $\xi_1, \dots, \xi_l, \tilde{\xi}_1, \dots, \tilde{\xi}_l \in G_n R^\omega$ the following holds:

$$G_n A^\omega \vdash \bigwedge_{i=1}^l \Pi_{k-2}^0\text{-CP}(\xi_i \underline{a}) \rightarrow (A(\underline{a}) \rightarrow \tilde{A}(\underline{a}))$$

and

$$G_n A^\omega \vdash \bigwedge_{i=1}^{\bar{l}} \Pi_{k-3}^0\text{-CP}(\tilde{\xi}_i \underline{a}) \rightarrow (\tilde{A}(\underline{a}) \rightarrow A(\underline{a}))$$

(Here and in the following we use the convention that $\Pi_k^0\text{-S}$ is empty (i.e. $\equiv (0 = 0)$) for an axiom schema S if $k < 0$).

Proof: Induction on k : For $k = 0, 1$ the lemma is trivial. So let $k \geq 1$.
 $k \mapsto k + 1$: Consider

$$A(\underline{a}) \equiv \forall x_1^0 \exists x_2^0 \dots \exists^{(d)} x_{k+1}^0 A_0(\underline{a}, x_1, x_2, \dots, x_{k+1}) \in \Pi_{k+1}^0.$$

By the induction hypothesis applied to the Π_k^0 -formula

$$\forall x_2 \exists x_3 \dots \forall^{(d)} x_{k+1} \neg A_0(\underline{a}, x_1, \dots, x_{k+1})$$

we have instances $\Pi_{k-2}^0\text{-CP}(\xi_i \underline{a})$ (note that instances of $\Pi_{k-3}^0\text{-CP}$ can be considered as instance of $\Pi_{k-2}^0\text{-CP}$ as well) such that $\bigwedge_i \Pi_{k-2}^0\text{-CP}(\xi_i \underline{a})$ implies (relative to $G_n A^\omega$)

$$\begin{aligned} \exists x_2 \forall x_3 \dots \exists^{(d)} x_{k+1} A_0 &\leftrightarrow \\ \exists x_2 \forall x_3 \dots \exists^{(d)} x_{k+1} \exists \tilde{x}_2 \leq x_2 \forall \tilde{x}_3 \leq x_3 \dots A_0(\underline{a}, x_1, \tilde{x}_2, \dots, \tilde{x}_{k+1}). \end{aligned}$$

Hence

$$\begin{aligned} A(\underline{a}) &\leftrightarrow \forall x_1 \exists x_2 \dots \exists^{(d)} x_{k+1} \exists \tilde{x}_2 \leq x_2 \dots \exists^{(d)} \tilde{x}_{k+1} \leq x_{k+1} A_0(\underline{a}, x_1, \tilde{x}_2, \dots, \tilde{x}_{k+1}) \\ &\leftrightarrow \forall x_1 \forall \tilde{x}_1 \leq x_1 \exists x_2 \dots \exists^{(d)} x_{k+1} \exists \tilde{x}_2 \leq x_2 \dots A_0(\underline{a}, \tilde{x}_1, \tilde{x}_2, \dots, \tilde{x}_{k+1}) \\ &\stackrel{\Pi_{k-1}^0\text{-CP}(\hat{\xi} \underline{a})}{\stackrel{\text{logic}}{\leftrightarrow}} \\ &\forall x_1 \exists x_2 \forall \tilde{x}_1 \leq x_1 \exists \hat{x}_2 \leq x_2 \forall x_3 \dots \exists^{(d)} x_{k+1} \exists \tilde{x}_2 \leq \hat{x}_2 \dots \exists^{(d)} \tilde{x}_{k+1} \leq x_{k+1} A_0(\underline{a}, \tilde{x}) \\ &\leftrightarrow \forall x_1 \exists x_2 \forall \tilde{x}_1 \leq x_1 \forall x_3 \dots \exists^{(d)} x_{k+1} \exists \tilde{x}_2 \leq x_2 \dots \exists^{(d)} \tilde{x}_{k+1} \leq x_{k+1} A_0(\underline{a}, \tilde{x}) \\ &\leftrightarrow \forall x_1 \exists x_2 \forall x_3 \forall \tilde{x}_1 \leq x_1 \exists x_4 \dots \exists^{(d)} x_{k+1} \exists \tilde{x}_2 \leq x_2 \dots \exists^{(d)} \tilde{x}_{k+1} \leq x_{k+1} A_0(\underline{a}, \tilde{x}). \end{aligned}$$

In the same way as we shifted $\forall \tilde{x}_1 \leq x_1$ over $\exists x_2$ we now move $\forall \tilde{x}_1 \leq x_1$ over $\exists x_4$, then permute $\forall \tilde{x}_1 \leq x_1$ with $\forall x_5$, move over $\exists x_6$ and so on until we obtain $\tilde{A}(\underline{a})$. This requires only Π_{k-3}^0 -instances (or simpler ones) of CP

which can be considered a fortiori as instances $\Pi_{k-2}^0\text{-CP}(\zeta_j \underline{a})$. Putting things together we have shown that (relative to $G_n A^\omega$):

$$\Pi_{k-1}^0\text{-CP}(\widehat{\xi} \underline{a}) \wedge \bigwedge_i \Pi_{k-2}^0\text{-CP}(\xi_i \underline{a}) \wedge \bigwedge_j \Pi_{k-2}^0\text{-CP}(\zeta_j \underline{a}) \rightarrow (A(\underline{a}) \rightarrow \tilde{A}(\underline{a}))$$

and

$$\bigwedge_i \Pi_{k-2}^0\text{-CP}(\xi_i \underline{a}) \wedge \bigwedge_j \Pi_{k-2}^0\text{-CP}(\zeta_j \underline{a}) \rightarrow (\tilde{A}(\underline{a}) \rightarrow A(\underline{a})),$$

which concludes the proof of the lemma.

Since in our main results we assume $n \geq 2$ or $n \geq 3$ for the level n of $G_n A^\omega$ we also use for simplicity $G_2 A^\omega$ in the following definition and lemmas although some of them can be carried out even in $G_1 A^\omega$.

Definition 3.11 (and lemma) *For $m \in \mathbb{N}$ let $\Phi \in G_2 R^\omega$ be such that*

$$\begin{aligned} G_2 A^\omega \vdash & \forall f^{(0)\dots(0)}, x^0, y_1^0, z_1^0, \dots, y_m^0, z_m^0, (y_{m+1}) \\ & (\Phi f x y_1 z_1 \dots y_m z_m (y_{m+1}) =_0 0 \\ & \leftrightarrow \forall \tilde{y}_1 \leq y_1 \exists \tilde{z}_1 \leq z_1 \dots \forall \tilde{y}_m \leq y_m \exists \tilde{z}_m \leq z_m (\forall \tilde{y}_{m+1} \leq y_{m+1}) \\ & (f(x, \tilde{y}_1, \tilde{z}_1, \dots, \tilde{y}_m, \tilde{z}_m, (\tilde{y}_{m+1})) =_0 0)). \end{aligned}$$

We denote Φf by f' .

Lemma 3.12 *Let $k \geq 1$. There are (effectively) finitely many terms $\xi_1, \dots, \xi_l \in G_2 R^\omega$ such that*

$$G_2 A^\omega \vdash \forall f \left(\left(\bigwedge_{i=1}^l \Pi_{k-2}^0\text{-CP}(\xi_i f) \right) \rightarrow (\Pi_k^0\text{-CA}(f) \leftrightarrow \Pi_k^0\text{-CA}(f')) \right).$$

Proof: The lemma follows from lemma 3.10.

Definition 3.13 *The ‘monotone’ tertium-non-datur is given by*

$$\begin{aligned} \underline{\Pi_k^0\text{-TND}^{mon}}(f) & := \\ & \left\{ \begin{array}{l} \forall x^0 \exists u_1^0 \forall y_1^0 \exists z_1^0 \forall v_1^0 \dots \exists u_m^0 \forall y_m^0 \exists z_m^0 \forall v_m^0 (\exists u_{m+1}^0 \forall y_{m+1}^0) \forall \tilde{x} \leq x \\ (f'(\tilde{x}, y_1, z_1, \dots, y_m, z_m, (y_{m+1})) =_0 0 \vee f'(\tilde{x}, u_1, v_1, \dots, u_m, v_m, (u_{m+1})) \neq 0), \end{array} \right. \end{aligned}$$

Lemma 3.14 1) $G_2 A^\omega \vdash \forall f ((\underline{\Pi_k^0\text{-TND}^{mon}}(f))^S \rightarrow (\Pi_k^0\text{-TND}(f')^{pr})^S)$.

2) $G_2A^\omega \vdash \forall f(\text{Mon}^*(\Pi_k^0\text{-TND}^{\text{mon}}(f)))$.

Proof: 1) follows by putting $\tilde{x} := x$.

2) Follows immediately from the definition of $\Pi_k^0\text{-TND}^{\text{mon}}(f)$.

Proposition 3.15 $G_2A^\omega \vdash \forall f((\Pi_k^0\text{-TND}^{\text{mon}}(f))^S \rightarrow \Pi_k^0\text{-CA}(f'))$.

Proof: Lemmas 3.5 and 3.14.1.

Lemma 3.16 *One can construct a $\xi \in G_2R^\omega$ such that*

$$G_2A^\omega + AC^{0,0}\text{-qf} \vdash \forall f(\Pi_k^0\text{-CA}(\xi f) \rightarrow \Pi_k^0\text{-CP}(f)).$$

Proof:

Using $\Pi_k^0\text{-CA}(\xi f)$ for a suitable $\xi \in G_2R^\omega$ one can reduce $\Pi_k^0\text{-CP}(f)$ to $\Pi_0^0\text{-CP}$ which is provable in $G_2A^\omega + AC^{0,0}\text{-qf}$.

Proposition 3.17 *For a suitable $\xi \in G_2R^\omega$ one has*

$$G_2A^\omega + AC^{0,0}\text{-qf} \vdash \forall f((\Pi_k^0\text{-TND}^{\text{mon}}(\xi f))^S \rightarrow \Pi_k^0\text{-CA}(f)).$$

Proof: Induction on k : $k = 0, 1$: easy. Let $k > 1$ and let's assume that the proposition holds for all $m < k$. $\Pi_{k-2}^0\text{-CP}(\xi_i f)$ denote the instances of Π_{k-2}^0 -collection from lemma 3.12 which are needed to show

$$\Pi_k^0\text{-CA}(f) \leftrightarrow \Pi_k^0\text{-CA}(f').$$

Let $\hat{\xi} \in G_2R^\omega$ be (using lemma 3.16) such that¹³

$$(1) G_2A^\omega + AC^{0,0}\text{-qf} \vdash \Pi_{k-2}^0\text{-CA}(\hat{\xi} f) \rightarrow (\Pi_k^0\text{-CA}(f) \leftrightarrow \Pi_k^0\text{-CA}(f')).$$

By the induction hypothesis we have

$$(2) G_2A^\omega + AC^{0,0}\text{-qf} \vdash \forall f((\Pi_{k-2}^0\text{-TND}^{\text{mon}}(\tilde{\xi} f))^S \rightarrow \Pi_{k-2}^0\text{-CA}(f))$$

for a suitable $\tilde{\xi} \in G_2R^\omega$. So by proposition 3.15 (3) \equiv

$$G_2A^\omega + AC^{0,0}\text{-qf} \vdash (\Pi_k^0\text{-TND}^{\text{mon}}(f))^S \wedge (\Pi_{k-2}^0\text{-TND}^{\text{mon}}(\tilde{\xi}(\hat{\xi} f)))^S \rightarrow \Pi_k^0\text{-CA}(f).$$

¹³Note that two instances $\Pi_k^0\text{-CA}(\xi_1 f) \wedge \Pi_k^0\text{-CA}(\xi_2 f)$ can be coded together into one instance $\Pi_k^0\text{-CA}(\xi_3 f)$ in G_2A^ω .

Introducing dummy quantifiers, $(\Pi_{k-2}^0\text{-TND}^{mon}(\tilde{\xi}(\widehat{\xi}f)))^S$ can be reduced to $(\Pi_k^0\text{-TND}^{mon}(\xi^*f))^S$ for a suitable $\xi^* \in G_2R^\omega$. Furthermore

$$(4) (\Pi_k^0\text{-TND}^{mon}(h))^S \rightarrow (\Pi_k^0\text{-TND}^{mon}(f))^S \wedge (\Pi_k^0\text{-TND}^{mon}(g))^S$$

for

$$h(x, \underline{y}, \underline{z}) = \begin{cases} f(\tilde{x}, \underline{y}, \underline{z}) & \text{if } x = 2\tilde{x} \\ g(\tilde{x}, \underline{y}, \underline{z}) & \text{if } x = 2\tilde{x} + 1. \end{cases}$$

Hence

$$(5) (\Pi_k^0\text{-TND}^{mon}(\xi f))^S \rightarrow (\Pi_k^0\text{-TND}^{mon}(f))^S \wedge (\Pi_k^0\text{-TND}^{mon}(\xi^*f))^S$$

for a suitable $\xi \in G_2R^\omega$. By (3) and (5) we have

$$G_2A^\omega + \text{AC}^{0,0}\text{-qf} \vdash (\Pi_k^0\text{-TND}^{mon}(\xi f))^S \rightarrow \Pi_k^0\text{-CA}(f).$$

Lemma 3.18 *Let $k \geq 1$ and $A \in \Sigma_{k-1}^0$. Then*

$$G_3A^\omega + \Sigma_k^0\text{-IA} \vdash \forall x^0 \exists u^0 \forall \tilde{x} \leq_0 x (\forall y^0 A(\tilde{x}, y) \vee \exists \tilde{u} \leq u \neg A(\tilde{x}, \tilde{u})).$$

Proof: Assume

$$(+) \forall u^0 \exists \tilde{x} \leq x (\exists y \neg A(\tilde{x}, y) \wedge \forall \tilde{u} \leq u A(\tilde{x}, \tilde{u})).$$

We show by induction on n :

$$(*) \forall n \exists u, \tilde{x} \left(\overbrace{\left(\begin{array}{l} \text{lth } \tilde{x} = n + 1 \wedge \bigwedge_{\substack{i, j \leq n \\ i \neq j}} ((\tilde{x})_i \neq (\tilde{x})_j) \wedge \\ \bigwedge_{i \leq n} ((\tilde{x})_i \leq x) \wedge \forall i \leq n \exists \tilde{u} \leq u \neg A((\tilde{x})_i, \tilde{u}) \end{array} \right)}^{G(n, u, \tilde{x})} \right)$$

(For $n = x + 1$ this obviously is contradictory and so $\neg(+)$ is proved).

$n = 0$: $(+)$ applied to $u := 0$ yields an $x_0 \leq x$ such that $A(x_0, 0)$ and $\exists y_0 \neg A(x_0, y_0)$. $(*)$ is now satisfied by taking $\tilde{x} := \langle x_0 \rangle, u := y_0$.

$n \rightarrow n + 1$: Let u, \tilde{x} be such that $(*)$ is satisfied for n . By $(+)$ there exists an $x_{n+1} \leq x$ such that $\exists y_{n+1} \neg A(x_{n+1}, y_{n+1})$ and $\forall \tilde{u} \leq u A(x_{n+1}, \tilde{u})$. By $(*)$ we have $\forall i \leq n \exists \tilde{u} \leq u \neg A((\tilde{x})_i, \tilde{u})$. Hence $\forall i \leq n ((\tilde{x})_i \neq x_{n+1})$ and so $\hat{u} := \max(u, y_{n+1}), \hat{x} := \tilde{x} * \langle x_{n+1} \rangle$ satisfy $G(n + 1, \hat{u}, \hat{x})$.

It remains to show that $\exists u, \tilde{x} G(n, u, \tilde{x})$ is equivalent to a Σ_k^0 -formula:

Using Σ_{k-1}^0 -CP, $\exists \tilde{u} \leq u \neg A((\tilde{x})_i, \tilde{u})$ can be shown to be equivalent to a Π_{k-1}^0 -formula. Since Σ_{k-1}^0 -CP follows from Σ_k^0 -IA, the whole proof can be carried out in $G_3A^\omega + \Sigma_k^0$ -IA.

In contrast to Π_k^0 -TND(f) its monotone version Π_k^0 -TND^{mon}(f) does not hold logically. However it can be proved using Σ_k^0 -induction. More precisely the following proposition holds:

Proposition 3.19 *Let $k \geq 1$. There are finitely many instances Σ_k^0 -IA($\xi_i f$) such that*

$$G_3A^\omega \vdash \forall f \left(\left(\bigwedge_{i=1}^l \Sigma_k^0\text{-IA}(\xi_i f) \right) \rightarrow \Pi_k^0\text{-TND}^{\text{mon}}(f) \right).$$

Proof: By (the proof of) lemma 3.18 there are instances Σ_k^0 -IA($\xi_i f$) which prove (relatively to G_3A^ω)

$$(*) \left\{ \begin{array}{l} \forall x \exists u_1 \forall \tilde{x} \leq x (\forall y_1 \exists z_1 \dots \forall y_m \exists z_m (\forall y_{m+1}) (f'(\tilde{x}, y_1, z_1, \dots, y_m, z_m) = 0)) \\ \forall \exists \tilde{u} \leq u_1 \forall v_1 \dots \exists u_m \forall v_m (\exists u_{m+1}) (f'(\tilde{x}, \tilde{u}, v_1, \dots, u_m, v_m) \neq 0) \end{array} \right.$$

and therefore by the definition of f' (which makes

$\exists \tilde{u} \leq u_1 \forall v_1 \dots \exists u_m \forall v_m (\exists u_{m+1}) (f'(\tilde{x}, \tilde{u}, v_1, \dots, u_m, v_m, (u_{m+1})) \neq 0)$ monotone w.r.t. $\exists \tilde{u}$)

$$\left\{ \begin{array}{l} \forall x \exists u_1 \forall \tilde{x} \leq x (\forall y_1 \exists z_1 \dots \forall y_m \exists z_m (\forall y_{m+1}) (f'(\tilde{x}, y_1, z_1, \dots, y_m, z_m,) = 0)) \\ \forall \forall v_1 \dots \exists u_m \forall v_m (\exists u_{m+1}) (f'(\tilde{x}, u_1, v_1, \dots, u_m, v_m,) \neq 0)), \end{array} \right.$$

which is equivalent to $(**) :=$

$$\left\{ \begin{array}{l} \forall x \exists u_1 \forall y_1 \forall \tilde{x} \leq x \exists z_1 (\forall y_2 \dots \forall y_m \exists z_m (\forall y_{m+1}) (f'(\tilde{x}, y_1, z_1, \dots, y_m, z_m,) = 0)) \\ \forall \forall v_1 \dots \exists u_m \forall v_m (\exists u_{m+1}) (f'(\tilde{x}, u_1, v_1, \dots, u_m, v_m,) \neq 0)). \end{array} \right.$$

By a suitable instance of Π_{k-1}^0 -CP and the monotonicity of $(**)$ w.r.t. $\exists z_1$ one can ‘shift’ $\forall \tilde{x} \leq x$ over $\exists z_1$. Now one continues in this way until one obtains Π_k^0 -TND^{mon}(f) which needs only suitable instances of Π_l^0 -CP with $l < k - 1$ which can be considered as instances of Π_{k-1}^0 -CP. All the instances of Π_{k-1}^0 -CP used follow from suitable instances of Σ_k^0 -IA.

Corollary 3.20 $G_3A^\omega \vdash \forall f (\Pi_k^0\text{-CA}(\xi f) \rightarrow \Pi_k^0\text{TND}^{\text{mon}}(f))$ for a suitable $\xi \in G_3R^\omega$.

4 Conservation results for $\Pi_k^0\text{-AC}(f)$ and $\Delta_k^0\text{-CA}(f, g)$

We are now ready to determine the arithmetical content of instances $\Pi_k^0\text{-CA}(\xi uv)$ and even $\Pi_k^0\text{-AC}(\xi uv)$ and $\Delta_{k+1}^0\text{-CA}(\xi uv)$ in proofs of monotone sentences (and without monotonicity assumption if the logical complexity is restricted). It turns out that this content is given by certain instances of $\Pi_k^0\text{-TND}^{mon}$.

Definition 4.1

$$\Pi_k^0\text{-AC}(f) := \begin{cases} \forall l^0 (\forall x^0 \exists y^0 \forall u_1^0 \exists u_2^0 \dots \exists^{(d)} u_k^0 (f(l, x, y, \underline{u}) =_0 0) \\ \rightarrow \exists g^1 \forall x^0 \forall u_1^0 \exists u_2^0 \dots \exists^{(d)} u_k^0 (f(l, x, gx, \underline{u}) =_0 0) \end{cases}$$

$$\Delta_k^0\text{-CA}(f, g) := \begin{cases} \forall l^0 (\forall x^0 (\forall u_1^0 \exists u_2^0 \dots \exists^{(d)} u_k^0 (f(l, x, \underline{u}) =_0 0) \leftrightarrow \\ \exists v_1^0 \forall v_2^0 \dots \forall^{(d)} v_k^0 (g(l, x, \underline{v}) =_0 0)) \\ \rightarrow \exists h^1 \forall x^0 (gx =_0 0 \leftrightarrow \forall u_1 \exists u_2 \dots \exists^{(d)} u_k (f(l, x, \underline{u}) =_0 0)) \end{cases}$$

$$\Delta_k^0\text{-CA}(f) := \Delta_k^0\text{-CA}(j_1^1 f, j_2^1 f) \text{ for the projection functions } j_i^1 \in G_2 R^\omega.$$

Lemma 4.2 *Let $k \in \mathbb{N}$. Then for suitable $\xi_1, \xi_2 \in G_2 R^\omega$:*

- 1) $G_2 A^\omega + AC^{0,0}\text{-}qf \vdash \forall f (\Pi_k^0\text{-CA}(\xi_1 f) \rightarrow \Pi_k^0\text{-AC}(f)).$
- 2) $G_2 A^\omega + AC^{0,0}\text{-}qf \vdash \forall f (\Pi_k^0\text{-CA}(\xi_2 f) \rightarrow \Delta_{k+1}^0\text{-CA}(f)).$

Proof: Obvious.

Below we also need a certain ‘non-standard’ axiom F^-

$$F^- := \forall \Phi^{2(0)}, y^{1(0)} \exists y_0 \leq_{1(0)} y \forall k^0, z^1, n^0 (\bigwedge_{i <_0 n} (zi \leq_0 yki) \rightarrow \Phi k(\overline{z, n}) \leq_0 \Phi k(y_0 k)),$$

where, for $z^{\rho 0}$, $(\overline{z, n})(k^0) :=_\rho zk$, if $k <_0 n$ and $:= 0^\rho$, otherwise.

F^- does not hold in the full set-theoretic type-structure but can be eliminated from proofs of monotone sentences in our theories. This axiom was introduced and studied in [12] and implies the principle of uniform Σ_1^0 -boundedness which was mentioned in the introduction and which will be generalized in section 5 below.

Proposition 4.3 *Let $n \geq 2$, $k \geq 0$ and*

$B := \forall u^1 \forall v \leq_\tau tu \exists a_1^0 \forall b_1^0 \dots \exists a_l^0 \forall b_l^0 \exists w^\gamma B_0$ be a sentence in $\mathcal{L}(G_n A^\omega)$, where B_0 is quantifier-free and $t \in G_n R^\omega$. Let $\xi_1, \xi_2 \in G_n R^\omega$ (of suitable types) and Δ a set of sentences having the form $\forall x^\delta \exists y \leq_\rho sx \forall z^\eta A_0$ (A_0 quantifier-free, $s \in G_n R^\omega$). Then for a suitable $\xi \in G_n R^\omega$ the following holds:

$$\left\{ \begin{array}{l} \text{If} \\ G_n A^\omega + \Delta + AC\text{-qf} \vdash \\ \forall u^1 \forall v \leq_\tau tu (\Delta_{k+1}^0\text{-CA}(\xi_1 uv) \wedge \Pi_k^0\text{-AC}(\xi_2 uv)) \rightarrow \exists a_1^0 \forall b_1^0 \dots \exists a_l^0 \forall b_l^0 \exists w^\gamma B_0 \\ \text{then} \\ G_n A^\omega + \tilde{\Delta} + Mon(B) \vdash \\ \forall u^1 \forall v \leq_\tau tu (\Pi_k^0\text{-TND}^{mon}(\xi uv)) \rightarrow \exists a_1^0 \forall b_1^0 \dots \exists a_l^0 \forall b_l^0 \exists w^\gamma B_0 \\ \text{and in particular} \\ G_{\max(3,n)} A^\omega + \Sigma_k^0\text{-IA} + \tilde{\Delta} + Mon(B) \vdash \forall u^1 \forall v \leq_\tau tu \exists a_1^0 \forall b_1^0 \dots \exists a_l^0 \forall b_l^0 \exists w^\gamma B_0. \end{array} \right.$$

In the assumption of the rule the theory $G_n A^\omega + \Delta + AC\text{-qf}$ can be strengthened to¹⁴ $(G_n A^\omega + \Delta + AC\text{-qf}) \oplus F^-$. Then in the first conclusion $G_n A^\omega$ must be replaced by $G_{\max(3,n)} A^\omega$.

Proof: By lemma 4.2, proposition 3.17 and the fact that two instances of $\Pi_k^0\text{-CA}$ can be coded together into a single instance of $\Pi_k^0\text{-CA}$, there is a $\xi \in G_n R^\omega$ such that

$$\begin{aligned} & G_n A^\omega + AC^{0,0}\text{-qf} \\ & \vdash \forall u^1 \forall v \leq_\tau tu ((\Pi_k^0\text{-TND}^{mon}(\xi uv))^S \rightarrow \Delta_{k+1}^0\text{-CA}(\xi_1 uv) \wedge \Pi_k^0\text{-AC}(\xi_2 uv)). \end{aligned}$$

So the assumption of the rule implies

$$(1) \left\{ \begin{array}{l} G_n A^\omega + AC\text{-qf} + \Delta \vdash \\ \forall u^1 \forall v \leq_\tau tu (\Pi_k^0\text{-TND}^{mon}(\xi uv)) \rightarrow \exists a_1^0 \forall b_1^0 \dots \exists a_l^0 \forall b_l^0 \exists w^\gamma B_0. \end{array} \right.$$

By lemma 3.14.2) the prenexation¹⁵

$$A^{pr} := \forall u^1 \forall v \leq_\tau tu \exists x \forall u_1 \exists y_1 \forall z_1 \exists v_1 \dots \exists a_1 \forall b_1 \dots \exists w^\gamma (\text{TND}_0^{mon}(\xi uv) \rightarrow B_0)$$

¹⁴Here \oplus means that F^- must not be used in the proof of the premise of an application of the quantifier-free rule of extensionality QF-ER. $G_n A^\omega$ satisfies the deduction theorem w.r.t \oplus but not w.r.t $+$.

¹⁵Note that A^{pr} is not completely in prenex normal form because of the universal quantifiers hidden in $v \leq_\tau tu$. However it has

the form required in theorem 2.7 used below.

of¹⁶

$$A := \forall u^1 \forall v \leq_\tau tu(\Pi_k^0\text{-TND}^{mon}(\xi uv) \rightarrow \exists a_1^0 \forall b_1^0 \dots \exists a_l^0 \forall b_l^0 \exists w^\gamma B_0)$$

is monotone if B is:

$$G_n A^\omega \vdash \text{Mon}(B) \rightarrow \text{Mon}(A^{pr}).$$

Now (1) implies

$$G_n A^\omega + \text{AC-}qf + \Delta \vdash (A^{pr})^H$$

and therefore using theorem 2.7

$$G_n A^\omega + \tilde{\Delta} + \text{Mon}(B) \vdash A^{pr} \text{ i.e.}$$

$$G_n A^\omega + \tilde{\Delta} + \text{Mon}(B) \vdash A.$$

The second part of the claim in the proposition now follows from proposition 3.19.

The proof above can be combined with the elimination procedure for F^- given in [12](thm.4.21) yielding the claim about adding F^- .

Corollary 4.4 *Let $k \geq 1, \gamma \leq 2$ and $\xi_1, \xi_2 \in G_n R^\omega$. Then the following rule holds*

$$\left\{ \begin{array}{l} G_\infty A^\omega + \Delta + \text{AC-}qf \vdash \\ \quad \forall u^1 \forall v \leq_\tau tu(\Delta_{k+1}^0\text{-}CA(\xi_1 uv) \wedge \Pi_k^0\text{-}AC(\xi_2 uv) \rightarrow \exists w^\gamma B_0(u, v, w)) \\ \Rightarrow \exists \Phi \in T_{k-1} \text{ such that} \\ \text{PA}_i^\omega + \tilde{\Delta} \vdash \forall u^1 \forall v \leq_\tau tu \exists w \leq_\gamma \Phi u B_0(u, v, w). \end{array} \right.$$

Again we may strengthen the theory in the assumption of the rule above by $\oplus F^-$.

Proof: The corollary follows from proposition 4.3 by observing that the condition $\text{Mon}(\forall u^1 \forall v \leq_\tau tu \exists w^\gamma B_0)$ is empty and using the fact that $G_\infty A^\omega + \tilde{\Delta} + \Sigma_k^0\text{-IA}$ has a monotone functional interpretation as developed in [9] (via negative translation) in $\text{PA}_i^\omega + \tilde{\Delta}$ by terms $\in T_{k-1}$. The latter follows from the proof that the negative translation of $\Sigma_k^0\text{-IA}$ has a functional interpretation in T_{k-1} (provable in (a subsystem of) PA_i^ω) as given in [19] and the fact that every (closed) term of T_{k-1} can be majorized (in the sense of definition 2.1) by a suitable term in T_{k-1} which follows from Howard's proof of this fact for full T as given in [6].

¹⁶ TND_0^{mon} denotes the quantifier-free matrix of (some prenex normal form of) $\Pi_k^0\text{-TND}^{mon}$.

Corollary 4.5 *Let $n \geq 3$ and A be a Π_1^1 -sentence.*

*If $E\text{-}G_n A^\omega + AC^{1,0}\text{-}qf + \Delta_{k+1}^0\text{-}CA^- + \Pi_k^0\text{-}AC^- + WKL \vdash A$
then $G_n A^\omega + \Sigma_k^0\text{-}IA + Mon(A) \vdash A$.*

Proof:

Using the deduction theorem for $E\text{-}G_n A^\omega$, the fact that $E\text{-}G_3 A^\omega + AC^{1,0}\text{-}qf + F^-$ proves WKL (see [12]) and the existence of characteristic terms $\in G_n R^\omega$ for quantifier-free formulas of $E\text{-}G_n A^\omega$ the assumption implies

$$E\text{-}G_n A^\omega + AC^{1,0}\text{-}qf + F^- \vdash \bigwedge_{i=1}^l (\Delta_{k+1}^0\text{-}CA(\xi_i)) \wedge \bigwedge_{j=1}^{\bar{l}} (\Pi_k^0\text{-}AC(\tilde{\xi}_j)) \rightarrow A$$

for certain terms $\xi_i, \tilde{\xi}_j \in G_n R^\omega$ (corresponding to the universal closures of the instances of $\Delta_{k+1}^0\text{-}CA^-$ and $\Pi_k^0\text{-}AC^-$ used in the proof).

For suitable $\xi, \tilde{\xi} \in G_n R^\omega$ we have

$$G_n A^\omega \vdash \Delta_{k+1}^0\text{-}CA(\xi) \rightarrow \bigwedge_{i=1}^l (\Delta_{k+1}^0\text{-}CA(\xi_i))$$

and

$$G_n A^\omega \vdash \Pi_k^0\text{-}AC(\tilde{\xi}) \rightarrow \bigwedge_{j=1}^{\bar{l}} (\Pi_k^0\text{-}AC(\tilde{\xi}_j)).$$

Together with elimination of extensionality (see e.g. [16]) we obtain

$$(G_n A^\omega + AC^{1,0}\text{-}qf) \oplus F^- \vdash \Delta_{k+1}^0\text{-}CA(\xi) \wedge \Pi_k^0\text{-}AC(\tilde{\xi}) \rightarrow A.$$

The conclusion now follows from proposition 4.3.

Lemma 4.6 *Let $\forall u^1 \forall v \leq_\tau tu A(u, v)$ be a sentence with $A(u, v) \in \Sigma_{k+1}^0$. Then one can construct a sentence $\forall u^1 \forall v \leq_\tau tu \tilde{A}(u, v)$ with $\tilde{A}(u, v) \in \Sigma_{k+1}^0$ such that*

- 1) $G_n A^\omega \vdash Mon(\forall u^1 \forall v \leq_\tau tu \tilde{A}(u, v))$,
- 2) $G_n A^\omega \vdash \forall u^1 \forall v \leq_\tau tu (\bigwedge_{i=1}^l \Pi_{k-2}^0\text{-}CP(\xi_i uv) \rightarrow (A(u, v) \rightarrow \tilde{A}(u, v)))$,
- 3) $G_n A^\omega \vdash \forall u^1 \forall v \leq_\tau tu (\bigwedge_{i=1}^{\bar{l}} \Pi_{k-1}^0\text{-}CP(\tilde{\xi}_i uv) \rightarrow (\tilde{A}(u, v) \rightarrow A(u, v)))$,

where $\xi_i, \tilde{\xi}_j \in G_n R^\omega$ are suitable terms.

Proof: Lemmas 3.8,3.10.

Corollary 4.7 *Let $n \geq 3$, $\forall u^1 \forall v \leq_\tau tu A(u, v)$ be a sentence in $G_n A^\omega$ with $A(u, v) \in \Sigma_{k+1}^0$, $t \in G_n R^\omega$ and $\xi_1, \xi_2 \in G_n R^\omega$ of suitable types. Then the following rule holds:*

$$\left\{ \begin{array}{l} \text{If } G_n A^\omega + \Delta + AC\text{-qf} \vdash \\ \quad \forall u^1 \forall v \leq_\tau tu(\Delta_{k+1}^0\text{-CA}(\xi_1 uv) \wedge \Pi_k^0\text{-AC}(\xi_2 uv) \rightarrow A(u, v)) \\ \text{then } G_n A^\omega + \Sigma_k^0\text{-IA} + \tilde{\Delta} \vdash \forall u^1 \forall v \leq_\tau tu A(u, v). \end{array} \right.$$

We may strengthen the theory in the assumption of the rule above by $\oplus F^-$.

Proof:

Let \tilde{A} be as in lemma 4.6. $\Pi_{k-2}^0\text{-CP}(\xi_i uv)$ follows from a corresponding instance $\Pi_{k-2}^0\text{-AC}(\hat{\xi}_i uv)$ of $\Pi_{k-2}^0\text{-AC}$ which can be considered as an instance $\Pi_k^0\text{-AC}(\hat{\xi}_i uv)$ of $\Pi_k^0\text{-AC}$. All these instances $\Pi_k^0\text{-AC}(\hat{\xi}_i uv)$ ($i = 1, \dots, l$) can be combined with $\Pi_k^0\text{-AC}(\xi_2 uv)$ into a single instance $\Pi_k^0\text{-AC}(\hat{\xi}_2 uv)$. Hence the assumption of the corollary yields

$$G_n A^\omega + \Delta + AC\text{-qf} \vdash \\ \forall u^1 \forall v \leq_\tau tu(\Delta_{k+1}^0\text{-CA}(\xi_1 uv) \wedge \Pi_k^0\text{-AC}(\hat{\xi}_2 uv) \rightarrow \tilde{A}(u, v)).$$

The conclusion now follows from proposition 4.3, lemma 4.6 and the fact that $G_n A^\omega + \Sigma_k^0\text{-IA} \vdash \Pi_{k-1}^0\text{-CP}$.

Corollary 4.8 *For $n \geq 3$,*

$E\text{-}G_n A^\omega + AC^{1,0}\text{-qf} + \Delta_{k+1}^0\text{-CA}^- + \Pi_k^0\text{-AC}^- + WKL$ is conservative w.r.t. Π_{k+2}^0 -sentences over $G_n A^\omega + \Sigma_k^0\text{-IA}^-$.

Proof: The corollary follows from the proofs of corollary 4.5 and corollary 4.7.

Remark 4.9 Corollary 4.8 is optimal in the following sense. For every k there is a sentence $A \in \Pi_{k+3}^0$ such that

$$G_3 A^\omega + \Pi_k^0\text{-AC}^- \vdash A, \text{ but } G_3 A^\omega + \Sigma_k^0\text{-IA} \not\vdash A.$$

Proof: There is a first-order instance A (i.e. without parameters of type level > 0) of $\Pi_k^0\text{-FAC}$ which does not follow from $\Sigma_k^0\text{-IA}$ relative to e.g. $G_3 A^\omega$ (see

[18]). It is clear that $G_3A^\omega + \Pi_k^0\text{-AC}^- \vdash A$. Since the universal closure of A can be shown to be equivalent to a Π_{k+3}^0 -sentence in $G_3A^\omega + \Sigma_k^0\text{-IA}^-$ (and hence in $G_3A^\omega + \Pi_k^0\text{-AC}^-$), the claim follows.

Corollary 4.10 *Let $\forall u^1\forall v \leq_\tau tu A(u, v)$ be a sentence with $A(u, v) \in \Sigma_{k+2}^0$. Then for $n \geq 3$ the following rule holds:*

$$\left\{ \begin{array}{l} \text{If } G_nA^\omega + \Delta_+ \text{ AC-}qf \vdash \\ \quad \forall u^1\forall v \leq_\tau tu(\Delta_{k+1}^0\text{-CA}(\xi_1uv) \wedge \Pi_k^0\text{-AC}(\xi_2uv) \rightarrow A(u, v)) \\ \text{then } G_nA^\omega + \Pi_k^0\text{-CP} + \tilde{\Delta} \vdash \forall u^1\forall v \leq_\tau tu A(u, v). \end{array} \right.$$

We may strengthen the theory in the assumption of the rule above by $\oplus F^-$.

Proof: The corollary follows analogously to the proof of corollary 4.7 using lemma 4.6 for $k+1$ instead of k and the well-known fact (see e.g. [18]) that $G_nA^\omega + \Pi_k^0\text{-CP} \vdash \Sigma_k^0\text{-IA}$.

Corollary 4.11 *For $n \geq 3$, $E\text{-}G_nA^\omega + \text{AC}^{1,0}\text{-}qf + \Delta_{k+1}^0\text{-CA}^- + \Pi_k^0\text{-AC}^- + \text{WKL}$ is conservative w.r.t. Π_{k+3}^0 -sentences over $G_nA^\omega + \Pi_k^0\text{-CP}^-$.*

Proof: The corollary follows from corollary 4.10 analogously to the proof of corollary 4.8.

Let EA be Kalmar-elementary arithmetic EA (with number quantifiers) and let us consider the variant $G_nA_-^\omega$ of G_nA^ω where the arbitrary true universal axioms 9) from its definition in [12] are replaced by the schema of quantifier-free induction (with arbitrary parameters)¹⁷ only. The results above also hold for $G_nA_-^\omega$ since no other universal axioms from 9) were used. EA can be considered as a subsystem of $G_3A_-^\omega$ and the latter is conservative over the former. Hence we obtain the following corollaries for EA:

Corollary 4.12 *Let A be an arbitrary sentence of EA. Then the following rule holds:*

$$\text{EA} + \Pi_k^0\text{-CP} \vdash A \Rightarrow \text{EA} + \Sigma_k^0\text{-IA} + \text{Mon}(A) \vdash A.$$

In particular we have the following

Corollary 4.13 *Let A, \tilde{A} be sentences from EA such that*

¹⁷Or equivalently the second-order axiom of quantifier-free induction.

- 1) $EA + \Pi_k^0\text{-}CP \vdash A \rightarrow \tilde{A}$,
- 2) $EA + \Sigma_k^0\text{-}IA \vdash \tilde{A} \rightarrow A$ and
- 3) $EA + \Sigma_k^0\text{-}IA \vdash \text{Mon}(\tilde{A})$.

Then $EA + \Pi_k^0\text{-}CP \vdash A$ implies $EA + \Sigma_k^0\text{-}IA \vdash A$.

Combined with lemma 4.6 we finally obtain

Corollary 4.14 (Paris-Kirby [17], H. Friedman)

$EA + \Pi_k^0\text{-}CP$ is Π_{k+2}^0 -conservative over $EA + \Sigma_k^0\text{-}IA$.

5 Generalized principles of uniform boundedness and their arithmetical content

In the following we define a generalization of the principle of uniform Σ_1^0 -boundedness $\Sigma_1^0\text{-UB}^-$ which was studied in [12],[14],[15]:

$$\Sigma_1^0\text{-UB}^- := \left\{ \begin{array}{l} \forall y^1{}^{(0)} (\forall k^0 \forall x \leq_1 yk \exists z^0 A(x, y, k, z) \rightarrow \exists \chi^1 \forall k^0, x^1, n^0 \\ \quad (\bigwedge_{i <_0 n} (xi \leq_0 yki) \rightarrow \exists z \leq_0 \chi k A((\overline{x, n}), y, k, z))), \end{array} \right.$$

where $A \equiv \exists l^0 A_0(l)$ is a purely existential formula.

$\Sigma_1^0\text{-UB}^-$ follows from F^- relative to $G_n A^\omega + AC^{1,0}\text{-}qf$ (for $n \geq 2$).

In $G_2 A^\omega + \Sigma_1^0\text{-UB}^-$ and hence in $G_2 A^\omega + F^- + AC^{1,0}\text{-}qf$ one can give very short and perspicuous proofs of various important analytical theorems like

- Every pointwise continuous function $f : [0, 1]^d \rightarrow \mathbb{R}$ is uniformly continuous
- The attainment of the maximum value of $f \in C([0, 1]^d, \mathbb{R})$ on $[0, 1]^d$
- The sequential form of the Heine–Borel covering property for $[0, 1]^d$
- Dini’s theorem
- The existence of a uniformly continuous inverse function for every strictly increasing continuous function $f : [0, 1] \rightarrow \mathbb{R}$.

Since F^- does not contribute to the growth of extractable bounds one can extract polynomial bounds from proofs in $G_2A^\omega + \Sigma_1^0\text{-UB}^- + \text{AC-qf}$.

Whereas the straightforward generalization of $\Sigma_1^0\text{-UB}^-$ to Π_k^0 -formulas is not consistent with G_nA^ω (see [15]), the following restricted form is (although it does – like $\Sigma_1^0\text{-UB}^-$ – not hold in the full set-theoretic type structure):

Definition 5.1 *Let $\rho = 0(0)(0)(1(0))(1)$, $k \geq 0$.*

$$\Pi_k^0\text{-UB}^- \upharpoonright (g) := \begin{cases} \forall \Phi^\rho, y^{1(0)}, a^0 (\forall k^0 \forall x \leq_1 yk \exists z^0 A(g, \Phi(x, y, k, z), k, z, a) \rightarrow \\ \exists \chi^1 \forall k^0 \forall x \leq_1 yk \forall l^0 \exists z \leq_0 \chi k A(g, \Phi(\overline{(x, l)}, y, k, z), k, z, a)), \end{cases}$$

where $A(g, v^0, k^0, z^0, a^0) := \forall u_1^0 \exists u_2^0 \dots \exists^{(d)} u_k^0 (g(v, k, z, a, \underline{u}) =_0 0) \in \Pi_k^0$.

Remark 5.2 $G_nA^\omega \vdash \Pi_0^0\text{-UB}^- \upharpoonright (t) \rightarrow \Sigma_1^0\text{-UB}^-$, where $t \in G_1R^\omega$ such that $t(v, k, z, a) =_0 v$.

In [15] we have shown that every single (sequence of) instance(s) of the Bolzano-Weierstraß principle for bounded sequences in \mathbb{R}^d and of the Ascoli-lemma (in the sense of [22]) follows from suitable instances of $\Pi_1^0\text{-UB}^- \upharpoonright$ and used this to calibrate precisely the contribution of such instances to the growth of extractable bounds. This indicates the mathematical relevance of our generalized principles of uniform boundedness.

Proposition 5.3 *Let $n \geq 2, k \geq 0$. For suitable $\xi \in G_nR^\omega$ we have*

$$G_nA^\omega + \text{AC}^{1,0}\text{-qf} \vdash F^- + \Pi_k^0\text{-CA}(\xi g) \rightarrow \Pi_k^0\text{-UB}^- \upharpoonright (g),$$

where g is a free (function) variable.

Proof:

For a suitable $\xi \in G_2R^\omega$, $\Pi_k^0\text{-CA}(\xi g)$ yields the existence of a function h such that

$$\forall v^0, k^0, z^0, a^0 (hvkza =_0 0 \leftrightarrow A(g, v, k, z, a)),$$

where A is as in definition 5.1. Using h , the assumption of $\Pi_k^0\text{-UB}^- \upharpoonright (g)$ can be expressed as

$$\forall k^0 \forall x \leq_1 yk \exists z^0 (h(\Phi(x, y, k, z), k, z, a) =_0 0).$$

By $\Sigma_1^0\text{-UB}^-$, which follows from F^- and $\text{AC}^{1,0}\text{-qf}$ relative to G_nA^ω (see [12]), this yields

$$\exists \chi^1 \forall k^0 \forall x \leq_1 yk \forall l^0 \exists z \leq_0 \chi k (h(\Phi(\overline{(x, l)}, y, k, z), k, z, a) =_0 0)$$

We now show that $\Pi_k^0\text{-CA}(f)$ in fact is implied by suitable instances of $\Pi_k^0\text{-UB}^-$:

Proposition 5.6 *Let $n \geq 2, k \geq 1$. For suitable $\xi_1, \dots, \xi_l \in G_2R^\omega$ we have*

$$G_n A^\omega \vdash \bigwedge_{i=1}^l \Pi_k^0\text{-UB}^-(\xi_i f) \rightarrow \Pi_k^0\text{-CA}(f),$$

where f is a free (function) variable.

Proof: Induction on k . $k = 1$: $\Pi_1^0\text{-CA}(f)$ is logically equivalent to

$$(1) \exists g \leq_1 1 \forall x^0, y^0 \exists z^0 ((gx =_0 0 \rightarrow f(x, y) =_0 0) \wedge (f(x, z) =_0 0 \rightarrow gx =_0 0))$$

and hence to (2) : \equiv

$$\neg \forall g \leq_1 1 \exists x^0, y^0 \forall z^0 \neg ((gx =_0 0 \rightarrow f(x, y) =_0 0) \wedge (f(x, z) =_0 0 \rightarrow gx =_0 0)).$$

For a suitable $\xi_1 \in G_2R^\omega$, $\Pi_1^0\text{-UB}^-(\xi_1 f)$ yields the equivalence of (2) and

$$(3) \begin{cases} \neg \exists n^0 \forall g \leq_1 1 \exists x, y \leq n \forall z^0 \\ \neg ((gx =_0 0 \rightarrow f(x, y) =_0 0) \wedge (f(x, z) =_0 0 \rightarrow gx =_0 0)) \end{cases}$$

i.e.

$$(4) \begin{cases} \forall n^0 \exists g \leq_1 1 \forall x \leq n \\ ((gx =_0 0 \rightarrow \forall y \leq n f(x, y) =_0 0) \wedge (\forall z (f(x, z) =_0 0) \rightarrow gx =_0 0)). \end{cases}$$

Define

$$gx := \begin{cases} 0^0 & \text{if } \forall y \leq n (f(x, y) =_0 0) \\ 1^0 & \text{otherwise.} \end{cases}$$

Let $k \geq 1$. $k \mapsto k + 1$:

$\Pi_{k+1}^0\text{-CA}(f)$ is equivalent to

$$(*) \begin{cases} \exists g \leq_1 1 \forall x^0, y^0 \exists z^0 ((gx =_0 0 \rightarrow \exists u_1^0 \forall u_2^0 \dots \forall^{(d)} u_k^0 (f(x, y, \underline{u}) =_0 0)) \wedge \\ (\exists u_1^0 \forall u_2^0 \dots \forall^{(d)} u_k^0 (f(x, z, \underline{u}) =_0 0) \rightarrow gx =_0 0)). \end{cases}$$

By induction hypothesis there exists an instance $\Pi_k^0\text{-UB}^-(\xi_2 f)$ (which can be considered as an instance $\Pi_{k+1}^0\text{-UB}^-(\xi_2 f)$) which implies (relative to $G_n A^\omega$) $\Pi_k^0\text{-CA}(f)$ and hence the existence of an h such that

$$\forall x, a (h(x, a) =_0 0 \leftrightarrow \exists u_1 \forall u_2 \dots \forall^{(d)} u_k (f(x, a, \underline{u}) =_0 0)).$$

By $\Pi_{k+1}^0\text{-UB}^-\lrcorner(\xi_3 f)$ (for a suitable ξ_3) applied to the negation of $(*)$, $\Pi_{k+1}^0\text{-CA}(f)$ is equivalent to

$$(**) \left\{ \begin{array}{l} \forall n \exists g \leq_1 \forall x \leq n \left((gx =_0 0 \rightarrow \forall y \leq n \exists u_1^0 \forall u_2^0 \dots \forall^{(d)} u_k^0 f(x, y, \underline{u}) =_0 0) \right. \\ \left. \wedge (\forall z \exists u_1^0 \forall u_2^0 \dots \forall^{(d)} u_k^0 f(x, z, \underline{u}) = 0 \rightarrow gx =_0 0) \right), \end{array} \right.$$

which is satisfied by

$$gx := \begin{cases} 0^0 & \text{if } \forall y \leq n (h(x, y) = 0) \\ 1^0 & \text{otherwise.} \end{cases}$$

Corollary 5.7 *For $n \geq 2, k \geq 1$ the following holds:*

- 1) $G_n A^\omega \vdash \forall g \Pi_1^0\text{-UB}^-\lrcorner(g) \rightarrow \forall \tilde{g} \Pi_k^0\text{-CA}(\tilde{g})$.
- 2) $G_n A^\omega \vdash \forall g \Pi_1^0\text{-UB}^-\lrcorner(g) \leftrightarrow \forall \tilde{g} \Pi_k^0\text{-UB}^-\lrcorner(\tilde{g})$.

Proof: 1) By proposition 5.6 $\forall g \Pi_1^0\text{-UB}^-\lrcorner(g)$ implies $\forall f \Pi_1^0\text{-CA}(f)$ and hence $\forall f \Pi_k^0\text{-CA}(f)$ (by iteration).

2) follows from 1) and the proof of proposition 5.3.

Let $B_{0,1}$ be the type-0-bar recursor constant of equality rank 1, i.e. $B_{0,1}$ is characterized by the axioms

$$(\text{BR}_{0,1}) : \left\{ \begin{array}{l} x^2(\overline{y^1}, n^0) < n \rightarrow B_{0,1} x z u n y =_1 z \\ x(\overline{y}, \overline{n}) \geq n \rightarrow B_{0,1} x z u n y =_1 u(\lambda D^0. B_{0,1} x z u n'(\overline{y}, \overline{n} * D)), \end{array} \right.$$

where u is of type $1(1(0))$ and

$$(\overline{y}, \overline{n} * D)(k^0) =_1 \begin{cases} xk, & \text{if } k < n \\ D, & \text{if } k = n \\ 0^1, & \text{otherwise.} \end{cases}$$

Proposition 5.8 *Let $n \geq 3, k \geq 1$, $B_0(u, v, w)$ be a quantifier-free formula of $G_n A^\omega$ containing only u, v, w free, $t^{\tau 1} \in G_n R^\omega, \gamma \leq 2$. Then the following rule holds:*

$$\left\{ \begin{array}{l} G_n A^\omega + \Delta + AC\text{-qf} \vdash \forall g \Pi_k^0\text{-UB}^-\lrcorner(g) \rightarrow \forall u^1 \forall v \leq_\tau t u \exists w^\gamma B_0(u, v, w) \\ \Rightarrow \exists \Phi \in G_n R^\omega [B_{0,1}] \text{ such that} \\ G_n A^\omega + \tilde{\Delta} + (\text{BR}_{0,1}) + (DC^0) \vdash \forall u^1 \forall v \leq_\tau t u \exists w \leq_\gamma \Phi u B_0(u, v, w), \end{array} \right.$$

where

$$(DC^0) := \forall x^0 \exists y^0 A(x, y) \rightarrow \forall x^0 \exists z^1 (z0 =_0 x \wedge \forall z_1^0 A(z z_1, z(z'_1))).$$

Proof: By proposition 5.3 and corollary 5.7 one has

$$G_n A^\omega + AC^{1,0}\text{-qf} + \forall g \Pi_1^0\text{-CA}(g) \vdash F^- \rightarrow \forall \tilde{g} \Pi_k^0\text{-UB}^- \vdash (\tilde{g}).$$

Hence the assumption of the rule to be proved yields

$$G_n A^\omega + \Delta + AC\text{-qf} + \forall g \Pi_1^0\text{-CA}(g) \vdash F^- \rightarrow \forall u^1 \forall v \leq_\tau tu \exists w^\gamma B_0(u, v, w).$$

From the work of Spector [23] it follows that $G_n A^\omega + AC\text{-qf} + \forall g \Pi_1^0\text{-CA}(g)$ has (via negative translation) a Gödel functional interpretation in $G_n A_i^\omega + (BR_{0,1})$ by terms $\in G_n R^\omega[B_{0,1}]$. In [2] it is shown that the type structure \mathcal{M}^ω of the so-called strongly majorizable functionals forms a model of full bar recursion. From the proof of this fact (restricted to type-0-bar recursion) one obtains the construction of a term $B_{0,1}^* \in G_n R^\omega[B_{0,1}]$ such that

$$G_n A^\omega + (BR_{0,1}) + (DC^0) \vdash B_{0,1}^* \text{ s-maj } B_{0,1},$$

where ‘s-maj’ is the corresponding syntactic notion of strong majorization as defined in definition 2.1. Therefore the proof of the fact that (the negative translation of) $G_n A^\omega + AC\text{-qf} + \Delta$ has a monotone functional interpretation (in the sense of [9]) in $G_n A_i^\omega$ by terms in $G_n R^\omega$ (see [12]) extends to $G_n A^\omega + \Delta + AC\text{-qf} + \forall g \Pi_1^0\text{-CA}(g)$ yielding a monotone functional interpretation (via negative translation) in $G_n A^\omega + \tilde{\Delta} + (BR_{0,1}) + (DC^0)$ by terms in $G_n R^\omega[B_{0,1}]$. This has the consequence that as in the case of $G_n A^\omega + \Delta + AC\text{-qf}$ (see the proof of theorem 4.21 in [12]) we can eliminate F^- from the proof of $\forall u \forall v \leq tu \exists w B_0$ and extract a uniform bound Φ on ‘ $\exists w$ ’ which now of course is only in $G_n R^\omega[B_{0,1}]$ (instead of $G_n R^\omega$) and its verification can be carried out in $G_n A^\omega + \tilde{\Delta} + (BR_{0,1}) + (DC^0)$.

References

- [1] Beklemishev, L.D., A proof-theoretic analysis of collection. Preprint 1996.
- [2] Bezem, M.A., Strongly majorizable functionals of finite type: a model for bar recursion containing discontinuous functionals. *J. Symb. Logic* **50** pp. 652–660 (1985).

- [3] Clote, P., Hájek, P., Paris, J., On some formalized conservation results in arithmetic. *Arch. Math. Logic* **30**, pp. 201-218 (1990).
- [4] Gödel, K., Zur intuitionistischen Arithmetik und Zahlentheorie. *Ergebnisse eines Mathematischen Kolloquiums*, vol. 4 pp. 34–38 (1933).
- [5] Gödel, K., Über eine bisher noch nicht benutzte Erweiterung des finiten Standpunktes. *Dialectica* **12**, pp. 280–287 (1958).
- [6] Howard, W.A., Hereditarily majorizable functionals of finite type. In: Troelstra (1973).
- [7] Kay, R., *Models of Peano Arithmetic*. Oxford Logic Guides 15, Clarendon Press Oxford 1991.
- [8] Kohlenbach, U., Pointwise hereditary majorization and some applications. *Arch. Math. Logic* **31**, pp.227–241 (1992).
- [9] Kohlenbach, U., Analysing proofs in analysis. In: W. Hodges, M. Hyland, C. Steinhorn, J. Truss, editors, *Logic: from Foundations to Applications. European Logic Colloquium* (Keele, 1993), pp. 225–260, Oxford University Press (1996).
- [10] Kohlenbach, U., Real growth in standard parts of analysis. Habilitationsschrift, pp. xv+166, Frankfurt (1995).
- [11] Kohlenbach, U., A note on the Π_2^0 -induction rule. *Arch. Math. Logic* **34**, pp. 279-283 (1995).
- [12] Kohlenbach, U., Mathematically strong subsystems of analysis with low rate of provably recursive functionals. *Arch. Math. Logic* **36**, pp. 31–71 (1996).
- [13] Kohlenbach, U., Elimination of Skolem functions for monotone formulas. To appear in: *Archive for Mathematical Logic*.
- [14] Kohlenbach, U., The use of a logical principle of uniform boundedness in analysis. To appear in: Proc. ‘Logic in Florence 1995’.
- [15] Kohlenbach, U., Arithmetizing proofs in analysis. To appear in: Proc. Logic Colloquium ’96 (San Sebastian).
- [16] Luckhardt, H., Extensional Gödel functional interpretation. A consistency proof of classical analysis. *Springer Lecture Notes in Mathematics* **306** (1973).

- [17] Paris, J., Kirby, L., Σ_n^0 -collection schema in arithmetic. In: Logic Colloquium '77, pp. 199-209. North-Holland, Amsterdam, 1978.
- [18] Parsons, C., On a number theoretic choice schema and its relation to induction. In: Intuitionism and proof theory, pp. 459-473. North-Holland, Amsterdam (1970).
- [19] Parsons, C., On n-quantifier induction. J. Symbolic Logic **37**, pp. 466–482 (1972).
- [20] Shoenfield, J.R., Mathematical Logic, Addison-Wesley, New York, 1967.
- [21] Sieg, W., Fragments of arithmetic. Ann. Pure Appl. Logic **28**, pp. 33-71 (1985).
- [22] Simpson, S.G., Which set existence axioms are needed to prove the Cauchy/Peano theorem for ordinary differential equations. J. Symbolic Logic **49**, pp. 783–801 (1984).
- [23] Spector, C., Provably recursive functionals of analysis: a consistency proof of analysis by an extension of principles formulated in current intuitionistic mathematics. In: Recursive function theory, Proceedings of Symposia in Pure Mathematics, vol. 5 (J.C.E. Dekker (ed.)), AMS, Providence, R.I., pp. 1–27 (1962).
- [24] Troelstra, A.S. (ed.) Metamathematical investigation of intuitionistic arithmetic and analysis. Springer Lecture Notes in Mathematics **344** (1973).

Recent BRICS Report Series Publications

- RS-97-23 Ulrich Kohlenbach. *On the Arithmetical Content of Restricted Forms of Comprehension, Choice and General Uniform Boundedness*. August 1997. 35 pp.
- RS-97-22 Carsten Butz. *Syntax and Semantics of the logic $\mathcal{L}_{\omega\omega}^\lambda$* . July 1997. 14 pp.
- RS-97-21 Steve Awodey and Carsten Butz. *Topological Completeness for Higher-Order Logic*. July 1997. 19 pp.
- RS-97-20 Carsten Butz and Peter T. Johnstone. *Classifying Toposes for First Order Theories*. July 1997. 34 pp.
- RS-97-19 Andrew D. Gordon, Paul D. Hankin, and Søren B. Lassen. *Compilation and Equivalence of Imperative Objects*. July 1997. iv+64 pp. Appears also as Technical Report 429, University of Cambridge Computer Laboratory, June 1997. To appear in *Foundations of Software Technology and Theoretical Computer Science: 17th Conference, FCT&TCS '97 Proceedings, LNCS, 1997*.
- RS-97-18 Robert Pollack. *How to Believe a Machine-Checked Proof*. July 1997. 18 pp. To appear as a chapter in the book *Twenty Five Years of Constructive Type Theory*, eds. Smith and Sambin, Oxford University Press.
- RS-97-17 Peter Bro Miltersen. *Error Correcting Codes, Perfect Hashing Circuits, and Deterministic Dynamic Dictionaries*. June 1997. 10 pp.
- RS-97-16 Noga Alon, Martin Dietzfelbinger, Peter Bro Miltersen, Erez Petrank, and Gábor Tardos. *Linear Hashing*. June 1997. 22 pp. A preliminary version appeared with the title *Is Linear Hashing Good?* in *The Twenty-ninth Annual ACM Symposium on Theory of Computing, STOC '97*, pages 465–474.
- RS-97-15 Pierre-Louis Curien, Gordon Plotkin, and Glynn Winskel. *Bistructures, Bidomains and Linear Logic*. June 1997. 41 pp.
- RS-97-14 Arne Andersson, Peter Bro Miltersen, Søren Riis, and Mikkel Thorup. *Dictionaries on AC^0 RAMs: Query Time $\Theta(\sqrt{\log n / \log \log n})$ is Necessary and Sufficient*. June 1997. 18 pp. Appears in *37th Annual Symposium on Foundations of Computer Science, FOCS '96 Proceedings*, pages 441–450.