Basic Research in Computer Science

BRICS RS-00-29 L. Santocanale: The Alternation Hierarchy for the Theory of μ-lattices

The Alternation Hierarchy for the Theory of μ-lattices

Luigi Santocanale

BRICS Report Series **RS-00-29**

ISSN 0909-0878 **November 2000**

Copyright © 2000, Luigi Santocanale.
BRICS, Department of Computer Science
University of Aarhus. All rights reserved.

Reproduction of all or part of this work
is permitted for educational or research use
on condition that this copyright notice is
included in any copy.

See back inner page for a list of recent BRICS Report Series publications.
Copies may be obtained by contacting:

BRICS
Department of Computer Science
University of Aarhus
Ny Munkegade, building 540
DK–8000 Aarhus C
Denmark
Telephone: +45 8942 3360
Telefax: +45 8942 3255
Internet: BRICS@brics.dk

BRICS publications are in general accessible through the World Wide Web and anonymous FTP through these URLs:

`http://www.brics.dk`
`ftp://ftp.brics.dk`
This document in subdirectory `RS/00/29/`

The alternation hierarchy for the theory of μ-lattices

Luigi Santocanale
luigis@brics.dk

BRICS*

Abstract

The alternation hierarchy problem asks whether every μ-term, that is a term built up using also a least fixed point constructor as well as a greatest fixed point constructor, is equivalent to a μ-term where the number of nested fixed point of a different type is bounded by a fixed number.

In this paper we give a proof that the alternation hierarchy for the theory of μ-lattices is strict, meaning that such number does not exist if μ-terms are built up from the basic lattice operations and are interpreted as expected. The proof relies on the explicit characterization of free μ-lattices by means of games and strategies.

1 Introduction

The alternation hierarchy problem is at the core of the definition of categories of μ-algebras [Niw85, Niw86] which we resume as follows. For a given equational theory $\mathbb{T}$, we let T_0 be the category of its partially ordered models and order preserving morphisms. Out of T_0 we can select objects and morphisms so that all the "desired" least prefix-points

*Basic Research in Computer Science,
Centre of the Danish National Research Foundation.

exist and are preserved, this process giving rise to a category S_1. The desired least prefix-points are those needed to have models of an iteration theory [BÉ93] where the dagger operation is interpreted as the least prefix-point. If we use, as a selection criterion, the existence of greatest postfix-points and their preservation, we obtain a category P_1. If we let T_1 be the intersection of S_1 and P_1, then T_1 is a quasivariety and the category of models of a theory $\mathbb{T}_1$ which is axiomatized by equational implications. We can repeat the process out of T_1 and $\mathbb{T}_1$ and the iteration of this process leads to construct categories $\mathrm{S}_n, \mathrm{P}_n, \mathrm{T}_n$ for arbitrary positive numbers n. The category of μ-$\mathbb{T}$-algebras is defined to be the inverse limit of the corresponding diagram of inclusions; the alternation hierarchy problem asks whether this process stops after a finite number of steps, i.e. whether the category of μ-$\mathbb{T}$-algebras is equivalent to a category among $\mathrm{S}_n, \mathrm{P}_n, \mathrm{T}_n$ for some $n \geq 0$. The main contribution presented in this paper is theorem 3.6 stating that the alternation hierarchy for the theory of μ-lattices is strict, i.e. that there is no such number when $\mathbb{T}$ is the theory of lattices.

The alternation hierarchy for the propositional μ-calculus has recently been shown to be strict in several cases [Arn99, Bra98a, Len96]; together with open problems on fix-point free polynomials in free lattices [FJN95], these results have challenged us to the hierarchy problem for the theory of μ-lattices; in particular we were interested in understanding whether the explicit characterization of free μ-lattices [San00a, San00b] could be of help.

It is our opinion that μ-algebras are algebraic objects suitable to generalize the role of iteration theories in the context of the theorization of communication and interactive computation. This statement is exemplified by the consideration of free μ-lattices which have been characterized by means of games and strategies. Games for free μ-lattices model bidirectional synchronous communication channels which can be recursively constructed from a few primitives: left and right choices – the lattice operations – and left and right iterations – the least and greatest fixed point operators. If G and H are games for a free μ-lattice, we say that $G \leq H$ if there exists a winning strategy in a compound game $\langle G, H \rangle$ for a player whom we call Mediator. Such a strategy can be understood as a protocol for letting the left user of the communication channel G communicate with the right user of H in an asynchronous way. The order theoretic point of view, which we adopt here, identifies two such channels G and H if there are protocols in both directions, i.e. winning

strategies for Mediator in both games $\langle G, H\rangle$ and $\langle H, G\rangle$. The analysis of different strategies, in the spirit of categorical proof theory and of the bicompletion of categories [HJ99, Joy95a, Joy95b], is probably a more appropriate setting in which to understand communication; this study is under way and suggests a possible characterization of free bicomplete categories with enough initial algebras and terminal coalgebras of functors. However, we can still ask whether the order theoretic identification is degenerate by posing the alternation hierarchy problem; its translation in the language of communication sounds as follows: *is every channel equivalent to another one where the number of alternations between left and right iterations is bounded by a fixed positive integer?* The negative answer we provide to the order theoretic problem implies also that a categorical identification is not degenerate; moreover the answer depends on a coincidence of order theoretic ideas with categorical ideas. There are games A for which the copycat strategy, which plays the role of the identity, is the unique strategy in the game $\langle A, A\rangle$; because of that, the asynchronous communication, which is the result of a protocol mediating between the left user of the left channel and the right user of the right channel, has the same dynamic as the communication along the single channel A; therefore it happens to be synchronous and we call these games *synchronizing*. These games impose strong conditions on the structure of games H equivalent to A, we show that they hard, i.e. they are good representatives of their equivalence class as far as we are concerned with their alternation complexity.

The ideas presented here have originated from Philip Whitman's proof that free lattices are not in general complete [Whi42]. It is difficult to relate these ideas with those contained in previous works on the alternation hierarchy for the propositional μ-calculus [Arn99, Bra98a, Bra98b, Len96, Niw86]. The main reason is that the traditional models of this calculus are boolean algebras of sets with modal operators and that the alternation hierarchy for the class of distributive μ-lattices is degenerate, since every distributive lattice is a μ-lattice. In particular, the main theorem presented in this paper cannot be derived from those results, at least not in a straightforward way. Existing techniques for proving hardness of a μ-calculus formula, as summarized in [Arn99], are diagonalization arguments and rely on the presence of the boolean complement; they cannot be applied in the context of μ-lattices. However, our technique is also a sort of a diagonalization argument, but of a categorical flavor. An analogous technique for the propositional μ-calculus would go through

the explicit characterization of the order relation by means of refutations [Wal95]. Other works [AM99, Bla92, Joy97] have studied the structure of games as models of linear logic. With respect to those works, the focus is here on the algebraic structure imposed by fix-points instead of the algebraic structure imposed by the multiplicative connectives of linear logic.

The paper is structured as follows. In section 2 we present definitions of key concepts and introduce the notation we shall use. In section 3 we present μ-lattices and the alternation hierarchy problem; we sketch the general strategy used to answer the problem. In section 4 we review the structure of free μ-lattices in view of the hierarchy; for the sake of completeness, we present once more the proof that the preorder relation on games is transitive. In section 5 we define synchronizing games and prove their hardness. In section 6 we construct synchronizing games of arbitrary complexity.

2 Notation and useful definitions

2.1 Least and greatest fix-points

Let P be a partially ordered set and let $\phi : P \longrightarrow P$ be an order preserving function. The *least prefix-point* of ϕ, whenever it exists, is an element $\mu_z.\phi(z)$ of P such that $\phi(\,\mu_z.\phi(z)\,) \leq \mu_z.\phi(z)$ and such that, if $\phi(\,p\,) \leq p$, then $\mu_z.\phi(z) \leq p$. The *greatest postfix-point* of ϕ is defined dually and is denoted by $\nu_z.\phi(z)$. Least prefix-point and greatest postfix-point are Conway operators in the sense of [BÉ93]. A summary of their properties can be found in [Niw85].

2.2 Pointed graphs and trees with back edges

By a *graph* G we mean a tuple $\langle G_0, G_1, \mathrm{dom}\,, \mathrm{cod}\,\rangle$, where G_0 is a set (of vertexes or states), G_1 is a set (of directed edges or transitions) and $\mathrm{dom}\,, \mathrm{cod}\, : G_1 \longrightarrow G_0$ are functions. By a *morphism of graphs* $f : G \longrightarrow H$ we mean a pair of functions $f_i : G_i \longrightarrow H_i$, $i = 0, 1$, such that $f_0 \circ \mathrm{dom}\, = \mathrm{dom}\, \circ f_1$ and $f_0 \circ \mathrm{cod}\, = \mathrm{cod}\, \circ f_1$. We often write a graph as a pair $\langle G_0, G_1\rangle$ and leave in the background the functions $\mathrm{dom}\,, \mathrm{cod}\,$. If $G_1 \subseteq G_0 \times G_0$, we assume that dom and cod are the restrictions of the

projections to G_1.

Let $G = \langle G_0, G_1 \rangle$ be a graph, a *path* γ in G is a morphism of graphs $\gamma : \hat{n} \longrightarrow G$ where $\hat{n}$ is the graph $0 \to 1 \to \ldots \to n$. The length of γ, denoted $|\gamma|$, is defined to be n. We set $\mathrm{dom}\,\gamma = \gamma(0)$ and $\mathrm{cod}\,\gamma = \gamma(|\gamma|)$. Paths γ_1, γ_2 can be composed in the usual way, provided that $\mathrm{cod}\,\gamma_1 = \mathrm{dom}\,\gamma_2$, and we write $\gamma_1 \star \gamma_2$ for their composition; if $g \in G_0$ we write 1_g for the unique path γ such that $|\gamma| = 0$ and $\mathrm{dom}\,\gamma = g = \mathrm{cod}\,\gamma$. A category $F(G)$, free over G, is defined in this way. Let G, H be graphs and let $f : G \longrightarrow F(H)$ be a morphism of graphs, we say that f is *non-decreasing* if $|f(\tau)| \leq 1$ for every $\tau \in G_1$. The morphism of graphs f is non-decreasing if and only if its extension $f : F(G) \longrightarrow F(H)$ to a functor is *convex*, i.e. if $f(\gamma) = \delta_1 \star \delta_2$, then we can find γ_1, γ_2 such that $f(\gamma_i) = \delta_i$, $i = 1, 2$, and $\gamma = \gamma_1 \star \gamma_2$. We say that a path γ in G *visits* a vertex $g \in G_0$, or equivalently that g *lies* on γ, if there exists $i \in \{0, \ldots, |\gamma|\}$ such that $\gamma(i) = g$. We say that a path γ is *simple* if it does not visit a node twice, i.e. if γ_0 is injective as a function. We say that γ is a *cycle* if $\mathrm{dom}\,\gamma = \mathrm{cod}\,\gamma$ and that γ is *proper* if $|\gamma| > 0$.

A *pointed graph* is a tuple $\langle G_0, G_1, g_0 \rangle$ such that $\langle G_0, G_1 \rangle$ is a graph and $g_0 \in G_0$; we shall say that g_0 is the *root* of $\langle G_0, G_1, g_0 \rangle$. A *morphism of pointed graphs* $f : G \longrightarrow H$ is a morphism of graphs $f : \langle G_0, G_1 \rangle \longrightarrow \langle H_0, H_1 \rangle$ such that $f_0(g_0) = h_0$. An *infinite path* in G is a morphism of graphs $\gamma : \hat{\omega} \longrightarrow G$ where $\hat{\omega}$ is the graph $0 \to 1 \to \ldots \to n \to \ldots$. Since the pointed graph $\langle \hat{\omega}, 0 \rangle$ is the inductive limit of the pointed graphs $\langle \hat{n}, 0 \rangle$, we shall often identify an infinite path γ with the set $\{\gamma_n\}_{n \geq 0}$ of prefixes of γ of finite length. On the other hand, if $\{\gamma_n\}_{n \geq 0}$ is a set of paths such that $|\gamma_n| = n$ and $\gamma_{n+1} = \gamma_n \star \tau_{n+1}$, we shall use that same notation $\{\gamma_n\}_{n \geq 0}$ to denote the infinite path which associates to the transition $n \to n+1$ of $\hat{\omega}$ the transition τ_{n+1}. A pointed graph $\langle G_0, G_1, g_0 \rangle$ is said to be *reachable* if for every $g \in G_0$ there exists a path γ such that $\mathrm{dom}\,\gamma = g_0$ and $\mathrm{cod}\,\gamma = g$. Let G be a graph and let $g_0 \in G_0$, we denote by $\overline{G, g_0}$ the greatest subgraph H of G such that $\langle H, g_0 \rangle$ is reachable.

Definition 2.1 A *tree with back edges* is a pointed graph $\langle G_0, G_1, g_0 \rangle$ such that $G_1 \subseteq G_0 \times G_0$ and with the property that, for every vertex $g \in G_0$, there exists a unique simple path γ_g from g_0 to g. In this case, we say that an edge $\tau : g \to g'$ is a *forward edge* if $\gamma_g \star \tau = \gamma_{g'}$ and that it is a *back edge* otherwise.

To give a tree with back edges is equivalent to give the pair $\langle\langle G_0, F, g_0\rangle, B\rangle$, where F is the set of forward edges and B is the set of back edges. Then $\langle G_0, F, g_0\rangle$ is a tree, i.e. a pointed graph such that, for every vertex $g \in G_0$, there exists a unique path from g_0 to g; moreover, if $g \to g'$ is an edge from B, then g' is an ancestor of g in the tree $\langle G_0, F, g_0\rangle$. We can specify a tree with back edges by giving a pair $\langle T, B\rangle$, where $T = \langle T_0, T_1, t_0\rangle$ is a tree and $B \subseteq T_0 \times T_0$ is a set of pairs with the above property.

Let $\langle T, B\rangle$ be a finite tree with back edges. A vertex $r \in T_0$ is called a *return* if there exists a back edge $t \to r$. Observe that, for an infinite path γ in $\langle T, B\rangle$, there exists a unique return r_γ which is visited infinitely often and which is of minimal height. The height of a vertex in $\langle T, B\rangle$ is the length of the unique simple path from the root to the vertex. Similarly, for every proper cycle γ in $\langle T, B\rangle$, there exists a unique return r_γ of minimal height lying on γ. A vertex $x \in T_0$ is said to be a *leaf* if it is a leaf of T in the usual sense and there are no back edges from x; this can be summarized by saying that $\{\, g' \mid x \to g' \,\}$ is empty. There is an operation of substitution of a tree with back edges for a leaf induced by the analogous operations on trees. Let x be a leaf of $\langle T_1, B_1\rangle$, we define:

$$\langle T_1, B_1\rangle[\langle T_2, B_2\rangle/x] \;=\; \langle T_1[T_2/x], B_1 + B_2\rangle.$$

If $\langle T, B\rangle$ is a tree with back edges and $t \in T_0$, we say that t is a *complete vertex* if for every descendant t' of t and every back edge $t' \to r$, r is also a descendant of t. If t is a complete vertex, we can define trees with back edges $T^t_\downarrow$ and $T^\downarrow_t$ so that t is a leaf of $T^\downarrow_t$, t is the root of $T^t_\downarrow$, and moreover $T = T^\downarrow_t[T^t_\downarrow/t]$. Indeed, let $T_1 = \overline{T, t}$ and $T_2 = \overline{T', t_0}$, where $T'_0 = T_0$ and $T'_1 = T_1 \setminus \{\, \tau \mid \operatorname{dom}\tau = t \,\}$; define then $T^t_\downarrow$ as $\langle T_1, B|_{T_1}\rangle$ and $T^\downarrow_t$ as $\langle T_2, B|_{T_2}\rangle$. A return r is said to be *minimal* if there are no other returns on the path γ_r from the root to r. A minimal return is easily seen to be a complete vertex.

2.3 Games and strategies

Definition 2.2 A *partial game* is a tuple $G = \langle G_0, G_1, g_0, \epsilon, W_\sigma\rangle$ where $\langle G_0, G_1\rangle$ is a graph, $g_0 \in G_0$, $\epsilon : G_0 \longrightarrow \{0, \sigma, \pi\}$ is a coloring, and W_σ is a set of infinite paths in $\langle G_0, G_1\rangle$. We require that if $\epsilon(g) = 0$, then $\{\, g' \mid g \to g' \,\} = \emptyset$.

The above data must be interpreted as follows: G_0 is the set of *positions* of G, g_0 is the *initial position* and G_1 is the set of possible *moves*. For a position $g \in G_0$, if $\epsilon(g) = \sigma$, then it is player σ who must move, if $\epsilon(g) = \pi$, it is π's turn to move. A position $g \in G_0$ is final if there are no possible moves from g, i.e. if $\{ g' \mid g \to g' \} = \emptyset$. In this case, if $\epsilon(g) = \sigma$, then player σ loses, if $\epsilon(g) = \pi$, then player π loses, if $\epsilon(g) = 0$, then it is a draw and we call g a *partial final position*. We shall write X_G for the set $\{ x \in G_0 \mid \epsilon(x) = 0 \}$ of partial final positions of G. Finally, W_σ is the set of infinite plays which are *wins for player* σ. We define W_π to be the complement of W_σ; we assume that there are no infinite draws so that W_π is meant to be the set of infinite plays which are wins for player π. A *game* is a partial game G such that $X_G = \emptyset$. We shall say that a partial game is *bipartite* if $\epsilon(g) \neq \epsilon(g')$ for every move $g \to g'$.

In the definitions below, G will be a fixed a partial game as defined in 2.2.

Definition 2.3 A *winning strategy* for player σ in G is a non empty set S of paths in G satisfying the following properties:

- $\gamma \in S$ implies $\operatorname{dom}\gamma = g_0$,
- $\gamma_1 \star \gamma_2 \in S$ implies $\gamma_1 \in S$,
- if $\gamma \in S$ and $\epsilon(\operatorname{cod}\gamma) = \pi$, then $\gamma \star \tau \in S$ for every $\tau \in G_1$ such that $\operatorname{dom}\tau = \operatorname{cod}\gamma$,
- if $\gamma \in S$ and $\epsilon(\operatorname{cod}\gamma) = \sigma$, then there exists $\tau \in G_1$ such that $\operatorname{dom}\tau = \operatorname{cod}\gamma$ and $\gamma \star \tau \in S$,
- if γ is an infinite path in G such that for every $n \geq 0$ the prefix of γ of length n belongs to S, then γ belongs to W_σ.

A strategy for player σ in G is a nonempty set S of paths in G such that the first three properties hold. Let γ be a path in G, we say that γ has been played according to the strategy S if there exists a path γ_0 such that $\gamma_0 \star \gamma \in S$.

Definition 2.4 A *bounded memory winning strategy* for player σ in G is a pair (S, ψ), where $S = \langle S_0, S_1, s_0 \rangle$ is a finite reachable pointed graph and $\psi : S \longrightarrow \langle G_0, G_1, g_0 \rangle$ is a morphism of pointed graphs. The following properties hold:

- If $s \in S_0$ and $\epsilon(\psi(s)) = \pi$, then for every $\tau \in G_1$ such that $\operatorname{dom}\tau = \psi(s)$ there exists a transition $\tau' \in S_1$ such that $\operatorname{dom}\tau' = s$ and $\psi(\tau') = \tau$,
- if $s \in S$ and $\epsilon(\psi(s)) = \sigma$, then there exists $\tau' \in S_1$ such that $\operatorname{dom}\tau' = s$,
- if γ is an infinite path in S, then $\psi \circ \gamma \in W_\sigma$.

Lemma 2.5 Let (S, ψ) be a bounded memory winning strategy for player σ. Then the set of paths

$$\psi S \;=\; \{\, \psi \circ \gamma \,|\, \gamma \text{ is a path in } S \text{ and } \operatorname{dom}\gamma = s_0 \,\}$$

is a winning strategy for player σ as defined in 2.3.

3 The theory of μ-lattices and the hierarchy

We begin by defining μ-lattices and the hierarchy. We shall later give an equivalent but more combinatorial definition of μ-lattices using partial games. This approach will allow us to have a combinatorial grasp on the hierarchy.

Definition 3.1 The set of terms Λ_ω and the arity-function $a : \Lambda_\omega \longrightarrow \mathbb{N}$ are defined by induction as follows:

1. $\bigwedge_k \in \Lambda_\omega$ and $a(\bigwedge_k) = k$, for $k \geq 0$.
2. $\bigvee_k \in \Lambda_\omega$ and $a(\bigvee_k) = k$, for $k \geq 0$.
3. If $\phi \in \Lambda_\omega$, $a(\phi) = k$, and $\phi_i \in \Lambda_\omega$ for $i = 1, \ldots, k$, then $\phi \circ (\phi_1, \ldots, \phi_k) \in \Lambda_\omega$ and $a(\phi \circ (\phi_1, \ldots, \phi_k)) = \sum_{i=1,\ldots,k} a(\phi_i)$.
4. If $\phi \in \Lambda_\omega$, $a(\phi) = k + 1$, then $\mu_s.\phi \in \Lambda_\omega$ and $a(\mu_s.\phi) = k$, for $s = 1, \ldots, k + 1$.
5. If $\phi \in \Lambda_\omega$, $a(\phi) = k + 1$, then $\nu_s.\phi \in \Lambda_\omega$ and $a(\nu_s.\phi) = k$, for $s = 1, \ldots, k + 1$.

Definition 3.2 Let L be a lattice, we define a partial interpretation of terms $\phi \in \Lambda_\omega$, $a(\phi) = k$, as order preserving functions $|\phi| : L^k \longrightarrow L$.

1. $|\bigwedge_k|(l_1, \ldots, l_k) \;=\; \bigwedge_{i=1,\ldots,k} l_i$.
2. As in 1, but substituting each symbol $\bigwedge$ with the symbol $\bigvee$.
3. Let $\phi \in \Lambda_\omega$, $a(\phi) = k$, and let $\phi_i \in \Lambda_\omega$ for $i = 1, \ldots, k$. Suppose $|\phi|$ and $|\phi_i|$ are defined. In this case we define $|\phi \circ (\phi_1, \ldots, \phi_k)|$ to be:

$$\begin{aligned} &|\phi \circ (\phi_1, \ldots, \phi_k)|(l_1, \ldots, l_h) \\ &\qquad = \;\; |\phi|(\, |\phi_1|(l_{h_1^-}, \ldots, l_{h_1^+}), \ldots, |\phi_k|(l_{h_k^-}, \ldots, l_{h_k^+})\,)\,, \end{aligned}$$

   where $h_i^- = 1 + \sum_{j=1}^{i-1} a(\phi_j)$, $h_i^+ = \sum_{j=1}^{i} a(\phi_j)$ and $h = h_k^+ = \sum_{j=1}^{k} a(\phi_j)$. Otherwise $|\phi \circ (\phi_1, \ldots, \phi_k)|$ is undefined.
4. Let $\phi \in \Lambda_\omega$ be such that $a(\phi) = k + 1$. Suppose that $|\phi|$ is defined and let $s \in \{1, \ldots, k+1\}$. If for each vector $(l_1, \ldots, l_k) \in L^k$ the least prefix-point of the order preserving function $|\phi|(l_1, \ldots, l_{s-1}, z, l_s, \ldots, l_k)$ exists, then we define $|\mu_s.\phi|$ to be:

$$|\mu_s.\phi|(l_1, \ldots, l_k) \;=\; \mu_z.|\phi|(l_1, \ldots, l_{s-1}, z, l_s, \ldots, l_k)\,.$$

   Otherwise $|\mu_s.\phi|$ is undefined.
5. As in 4, but substituting each symbol μ with the symbol ν, and the word least prefix-point with the word greatest postfix-point.

Definition 3.3 A lattice L is a *μ-lattice* if the interpretation of terms $\phi \in \Lambda_\omega$ is a total function. Let L_1, L_2 be two μ-lattices. An order preserving function $f : L_1 \longrightarrow L_2$ is a *μ-lattice morphism* if the equality $|\phi| \circ f^{a(\phi)} = f \circ |\phi|$ holds for all $\phi \in \Lambda_\omega$. We shall write L_ω for the category of μ-lattices.

Definition 3.4 We define classes of terms $\Sigma_n, \Pi_n, \Lambda_n \subseteq \Lambda_\omega$, for $n \geq 0$. We set $\Sigma_0 = \Pi_0 = \Lambda_0$, where Λ_0 is the least class which contains $\bigvee_k$ and $\bigwedge_k$, $k \geq 0$, and which is closed under substitution (rule 3 of definition 3.1). Suppose that Σ_n and Π_n have been defined. We define Σ_{n+1} to be the least class of terms which contains $\Sigma_n \cup \Pi_n$ and which is closed under substitution and the μ-operation (rule 4 of definition 3.1). Similarly, we define Π_{n+1} to be the least class of terms which contains

$\Sigma_n \cup \Pi_n$ and which is closed under substitution and the ν-operation (rule 5 of definition 3.1). We let $\Lambda_n = \Pi_{n+1} \cap \Lambda_{n+1}$ and observe that $\Lambda_\omega = \bigcup_{n\geq 0} \Sigma_n = \bigcup_{n\geq 0} \Pi_n = \bigcup_{n\geq 0} \Lambda_n$.

Definition 3.5 We say that a lattice is a Σ_n-*model* if for every $\phi \in \Sigma_n$ $|\phi| : L^{a(\phi)} \longrightarrow L$ is defined. Let L_1, L_2 be two Σ_n-models, an order preserving function $f : L_1 \longrightarrow L_2$ is a *morphism of* Σ_n*-models* if for every $\phi \in \Sigma_n$ the equality $f \circ |\phi| = |\phi| \circ f^{a(\phi)}$ holds. We let S_n be the category of Σ_n-models and morphisms of Σ_n-models. We define in a similar way a Π_n-model, a morphism of Π_n-models and the category P_n, a Λ_n-model, a morphism of Λ_n-models and the category L_n.

Clearly L_0 is the category of lattices and we have inclusion of categories



The **alternation hierarchy problem** for the theory of μ-lattices can be stated in the following way: *is there a number* $n \geq 0$ *and a category* C_n *among* $\mathrm{S}_n, \mathrm{P}_n, \mathrm{L}_n$ *such that the inclusion functor* $\mathrm{L}_\omega \hookrightarrow \mathrm{C}_n$ *is an equivalence of categories*? If such a C_n exists, then $\mathrm{C}_n = \mathrm{L}_\omega$, since if P is a partially ordered set which is order-isomorphic to a μ-lattice, then it is itself a μ-lattice; as a consequence for every $m > n$ all the $\mathrm{S}_m, \mathrm{P}_n$ and L_n are equal to L_ω.

Theorem 3.6 The alternation hierarchy for the theory of μ-lattices is strict, i.e. there is no positive integer n such that $\mathrm{L}_n = \mathrm{L}_\omega$.

Proof. For every $n \geq 0$, we exhibit in 4.12 a sub-Λ_n-model $\mathcal{J}_{n,P}$ of the free μ-lattice $\mathcal{J}_P$ over the partially ordered set P. The Λ_n-model $\mathcal{J}_{n,P}$ is the free Λ_n-model over the partially ordered set P, in particular it is generated by P and the inclusion $i_{n,P} : \mathcal{J}_{n,P} \hookrightarrow \mathcal{J}_P$ preserves the generators. If $\mathrm{L}_n = \mathrm{L}_\omega$, then $i_{n,P}$ has to be an isomorphism; however, we show in 6.2 that $i_{n,P}$ is a proper inclusion for every $n \geq 0$ if P contains an antichain of cardinality six. □

The alternation hierarchy problem for a class $\mathcal{K}$ of μ-lattices can be stated as follows. Let $\overline{\mathcal{K}}_\omega$ be the quasi-variety generated by the class of μ-lattices $\mathcal{K}$ in $\mathcal{L}_\omega$, by which we mean the closure of the full sub-category determined by objects in $\mathcal{K}$ under products, sub-objects and regular epis. Similarly let $\overline{\mathcal{K}}_n$ be the quasi-variety generated by the class $\mathcal{K}$ in L_n. The inclusion functors $\mathrm{L}_\omega \hookrightarrow \mathrm{L}_n$ restrict to inclusions $\overline{\mathcal{K}}_\omega \hookrightarrow \overline{\mathcal{K}}_n$ and the problem is to determine whether there exists a number $n \geq 0$ such that the above inclusion is an equivalence. The above theorem has the following consequence.

Theorem 3.7 The alternation hierarchy for the class of complete lattices is strict.

Proof. Let $\mathcal{K}$ be the class of complete lattices. Since every free μ-lattice can be embedded in a complete lattice by a morphism of μ-lattices, as proved in [San00a, San00b], then $\overline{\mathcal{K}}_\omega = \mathrm{L}_\omega$. Similarly $\overline{\mathcal{K}}_n = \mathrm{L}_n$, since $\mathcal{J}_{n,P}$ is the free Λ_n-model, so that free Λ_n-models can be embedded into complete lattices. However $\mathrm{L}_n \neq \mathrm{L}_\omega$. □

The theory of μ-lattices has an equivalent presentation by means of a class $\mathcal{J}$ of partial games, cf. 2.2, which are a sort of combinatorial terms. A partial game G in this class comes always with its set of partial final positions X_G; given a lattice L we can define the partial interpretation of games $G \in \mathcal{J}$ as order preserving functions $|G| : L^{X_G} \longrightarrow L$. A lattice L turns out to be a μ-lattice if and only if the interpretation of a partial game $G \in \mathcal{J}$ is always defined.

Definition 3.8 A partial game G is in the class $\mathcal{L}$ if and only if $\langle G_0, G_1, g_0 \rangle$ is a finite tree with back edges and moreover $\gamma \in W_\sigma$ if and only if $\epsilon(r_\gamma) = \pi$. If $G \in \mathcal{L}$, we denote by $R(G)$ the set of positions which are returns of $\langle G_0, G_1, g_0 \rangle$ and by $\chi(G)$ the number $\operatorname{card} G_0 + \operatorname{card} R(G)$.

When specifying a partial game $G \in \mathcal{L}$ we shall omit to give the set W_σ, since this is determined by the underlying tree with back edges and the coloring ϵ.

Definition 3.9 On the class $\mathcal{L}$ the following constants and operations are defined.

0. x is the partial game with just one partial final position, which we call again x.

1. Let I be a finite set, the partial game $\bigwedge_I$ has starting position $\wedge_0 \notin I$, $\epsilon(\wedge_0) = \pi$, partial final positions i and moves $\wedge_0 \to i$ for every $i \in I$.

2. Let I be a finite set, the partial game $\bigvee_I$ is defined in a similar way: it has starting position $\vee_0 \notin I$, $\epsilon(\vee_0) = \sigma$, partial final positions i and moves $\vee_0 \to i$ for every $i \in I$.

3. *Substitution.* Let G and H be partial games in $\mathcal{L}$ and let $x \in X_G$. The underlying pointed graph of the game $G[H/x]$ is obtained by substitution of the tree with back edges underlying H for x in the tree with back edges underlying G. The coloring ϵ is defined accordingly, i.e. if $g \neq x$ is a position coming from G_0, then $\epsilon(g)$ is as in G, otherwise, for a position h coming from H_0, $\epsilon(h)$ is as in H.

4. *μ-operation.* Let $G \in \mathcal{L}$ be a partial game and let $x \in X_G$. The underlying graph of the game $\mu_x.G[x]$ is the same as the underlying graph of G with one more move $x \to g_0$. The initial position of $\mu_x.G[x]$ is x and we let $\epsilon(x) = \sigma$.

5. *ν-operation.* Let $G \in \mathcal{L}$ be a partial game and let $x \in X_G$. The underlying pointed graph of $\nu_x.G[x]$ is the same as the underlying graph of $\mu_x.G[x]$, however we let $\epsilon(x) = \pi$.

It is useful to have a picture of those operations.

0. The game x is

1. 2. Let $I = \{i_1, \ldots, i_n\}$ be a finite set. The games $\bigwedge_I$ and $\bigvee_I$ are:



3. Let G, H be partial games in $\mathcal{L}$ and let $x \in X_G$. We represent those games as:



The game $G[H/x]$ can be represented as:



4. 5. Let G be a partial game in $\mathcal{L}$ and let $x \in X_G$. This game can be represented as above. We represent the games $\mu_x.G[x]$ and $\nu_x.G[x]$ as:



Definition 3.10 We let $\mathcal{J}$ be the class of games $G \in \mathcal{L}$ for which the following two conditions hold, for every $r \in R(G)$:

1. there exists a unique back edge $P(r) \to r$,
2. there exists a unique move $r \to S(r)$.

We let $\mathcal{K}$ be the class of games $G \in \mathcal{L}$ for which only the first of the two conditions above holds, for any $r \in R(G)$.

We have the inclusion of classes $\mathcal{J} \subseteq \mathcal{K} \subseteq \mathcal{L}$, but those classes are essentially the same. A more detailed account of the equivalence between $\mathcal{J}$ and $\mathcal{K}$ is given in the proof of theorem 5.2.

We shall write $G = H$ if G, H are partial games and there exists an isomorphism of structure between them. This means that there exists an isomorphism f of the underlying pointed graphs such that $\epsilon = \epsilon \circ f_0$ and such that $\gamma \in W_\sigma$ if and only if $f \circ \gamma \in W_\sigma$. Substitution satisfies the commutativity rule $(G[H/x])[K/y] = (G[K/y])[H/x]$ if $x, y \in X_G$ and $x \neq y$. Hence if $\{ H_x \}_{x \in X_G}$ is a collection of games in $\mathcal{L}$, we shall write by $G[H_x/x]_{x \in X_G}$ for any sequence of substitutions.

Remark 3.11 It is possible to show that the class $\mathcal{J}$ is the least subclass of $\mathcal{L}$ which is closed under the constants and the operations of definition 3.9. Indeed, a stronger result holds: a partial game $G \in \mathcal{J}$ has a unique form x, $\bigwedge_I [H_i/i]_{i \in I}$, $\bigvee_I [H_i/i]_{i \in I}$, $\mu_x.H[x]$, $\nu_x.H[x]$, where H or the H_i satisfy $\chi(H) < \chi(G)$ and $\chi(H_i) < \chi(G)$. Hence, in what follows, we shall be able to define by induction on the structure of partial games in $\mathcal{J}$.

In a similar way as we did before, we define a partial interpretation for games in $\mathcal{J}$.

Definition 3.12 Let L be a lattice. We define an interpretation of partial games $G \in \mathcal{J}$ as order preserving functions $|G| : L^{X_G} \longrightarrow L$. The correspondence sending G to $|G|$ is in general only a partial function, i.e. $|G|$ could sometime be undefined.

0. Let $G = x$ so that $X_G = \{x\}$. We let $|G|(\lambda) = \lambda(x)$.
1. Let $G = \bigwedge_I [H_i/i]_{i \in I}$, so that $X_G = \sum_{i \in I} X_{H_i}$. If the $\{ |H_i| \}_{i \in I}$ are defined, then we define
$$|G|(\lambda) \;=\; \bigwedge_{i \in I} |H_i|(\lambda_{H_i}) \,,$$
where λ_{H_i} is the restriction of λ to X_{H_i}. Otherwise $|G|$ is undefined.
2. As in 1, but substituting each symbol $\bigwedge$ with the symbol $\bigvee$.
4. Let $G = \mu_x.H[x]$ so that $X_H = X_G \cup \{x\}$. If $|H|$ is defined and if also for each collection $\lambda \in L^{X_G}$ there exists the least prefix-point of the unary order preserving function
$$\phi(l) \;=\; |H|(\lambda^l) \,,$$

where $\lambda^l(y) = \lambda(y)$ if $y \neq x$ and $\lambda^l(x) = l$, then we define

$$|G|(\lambda) \quad = \quad \mu_z.\phi(\, z\,)\,.$$

Otherwise $|G|$ is undefined.

5. As in 4, but substituting each symbol μ with the symbol ν, and the word least prefix-point with the word greatest postfix-point.

Proposition 3.13 A lattice L is a μ-lattice if and only if $|G|$ is defined on L for every game $G \in \mathcal{J}$. An order preserving function $f : L_1 \longrightarrow L_2$ is a μ-lattice morphism if for all $G \in \mathcal{J}$ we have $f \circ |G| = |G| \circ f^{X_G}$, i.e.

$$f(\,|G|(\lambda)\,) \quad = \quad |G|(\,f \circ \lambda\,)\,,$$

for every $\lambda \in L_1^{X_G}$.

Proof. Using the correspondence between the rules of definition 3.1 and the operations defined in 3.9, inductively define for each $\phi \in \Lambda_\omega$ a pair $\langle G_\phi, \lambda_\phi \rangle$ where $G_\phi \in \mathcal{J}$ and $\lambda_\phi : a(\phi) \longrightarrow X_{G_\phi}$ is a bijection, with the following property: for each lattice L, G_ϕ is defined on L if and only if ϕ is defined and moreover

$$|\phi|(\lambda) \quad = \quad |G_\phi|(\lambda \circ \lambda_\phi)\,.$$

This shows that if L is a lattice such that $|G|$ is defined on L for every $G \in \mathcal{J}$, then L is a μ-lattice. The above formula leads to show that if f preserves the interpretation of games, then it is a μ-lattice morphism:

$$\begin{aligned} f \circ |\phi|(\lambda) &= f \circ |G_\phi|(\lambda \circ \lambda_\phi) \\ &= |G_\phi|(f \circ \lambda \circ \lambda_\phi) \\ &= |\phi|(f \circ \lambda)\,. \end{aligned}$$

On the other hand, assign to each $G \in \mathcal{J}$ a pair $\langle \phi_G, \lambda_G \rangle$ where $\phi_G \in \Lambda_\omega$ and $\lambda_G : a(\phi_G) \longrightarrow X_G$ is a bijection, so that for each lattice L, $|\phi_G|$ is defined on L if and only if $|G|$ is defined on L; moreover

$$|\phi_G|(\lambda) \quad = \quad |G|(\lambda \circ \lambda_G)\,.$$

This is done by induction on the structure of games in $\mathcal{J}$, cf. 3.11. The game x is sent to $\bigwedge_1$, the game $\bigwedge_I [G_i/i]_{i \in I}$ is sent to $\bigwedge_k \circ (\phi_{G_{\psi(1)}}, \ldots, \phi_{G_{\psi(k)}})$,

where $\psi : k \longrightarrow I$ is a bijection, and $\mu_x.G[x]$ is sent to $\mu_s.\phi_G$ where $s = \lambda_G^{-1}(x)$. Similar definitions are given for games of the form $\bigvee_I [G_i/i]_{i\in I}$ and $\nu_x.G[x]$.

The above assignment leads to show that if L is a μ-lattice, i.e. if for each $\phi \in \Lambda_\omega$ $|\phi|$ is defined on L, then for every $G \in \mathcal{J}$ $|G|$ is defined on L too. A morphisms which preserves the interpretation of terms will also preserve the interpretation of games:

$$\begin{aligned} f \circ |G|(\lambda) &= f \circ |G|(\lambda \circ \lambda_G^{-1} \circ \lambda_G) \\ &= f \circ |\phi_G|(\lambda \circ \lambda_G^{-1}) \\ &= |\phi_G|(f \circ \lambda \circ \lambda_G^{-1}) \\ &= |G|(f \circ \lambda \circ \lambda_G^{-1} \circ \lambda_G) \\ &= |G|(f \circ \lambda)\,. \end{aligned}$$

□

Definition 3.14 We define by induction classes of partial games $\mathcal{S}_n, \mathcal{P}_n$, $\mathcal{L}_n$, for $n \geq 0$. We set $\mathcal{S}_0 = \mathcal{P}_0 = \mathcal{L}_0$, where $\mathcal{L}_0$ is the least class which contains $x, \bigvee_I$ and $\bigwedge_I$, where I is a finite set, and which is closed under substitution. Suppose that $\mathcal{S}_n$ and $\mathcal{P}_n$ have been defined. We define $\mathcal{S}_{n+1}$ to be the least class of games which contains $\mathcal{S}_n \cup \mathcal{P}_n$ and which is closed under substitution and the μ-operation. Similarly, we define $\mathcal{P}_{n+1}$ to be the least class of games which contains $\mathcal{S}_n \cup \mathcal{P}_n$ and which is closed under substitution and the ν-operation. We let $\mathcal{L}_n = \mathcal{S}_{n+1} \cap \mathcal{P}_{n+1}$ and observe that $\mathcal{J} = \bigcup_{n\geq 0} \mathcal{S}_n = \bigcup_{n\geq 0} \mathcal{P}_n = \bigcup_{n\geq 0} \mathcal{L}_n$.

Proposition 3.15 A lattice is a Σ_n-model if and only if for every $G \in \mathcal{S}_n$ $|G| : L^{X_G} \longrightarrow L$ is defined. Let L_1, L_2 be two Σ_n-models, an order preserving function $f : L_1 \longrightarrow L_2$ is a morphism of Σ_n-models if and only if for every $G \in \mathcal{S}_n$ we have $f \circ |G| = |G| \circ f^{X_G}$. Analogous results hold for the classes $\mathcal{P}_n$ and $\mathcal{L}_n$, Π_n-models, and Λ_n-models, Π_n-morphisms and Λ_n-morphisms, respectively.

Proof. The transformation of terms into partial games $\phi \longmapsto \langle G_\phi, \lambda_\phi \rangle$, which we defined in the proof of proposition 3.13, restricts to a transformation $\Sigma_n \longrightarrow \mathcal{S}_n$, so that if G is defined on L for every $G \in \mathcal{S}_n$, then L is a Σ_n-model, and a lattice morphism which preserves the interpretation of every partial game in $\mathcal{S}_n$ is a morphism of Σ_n-models.

In a similar way the transformation $G \longmapsto \langle \phi_G, \lambda_G \rangle$ carries partial games in $\mathcal{S}_n$ into terms in Σ_n. $\square$

In the rest of this section we give a combinatorial characterization of the classes $\mathcal{S}_n, \mathcal{P}_n, \mathcal{L}_n$.

Definition 3.16 Let $G \in \mathcal{L}$. A *chain* C in G is a totally ordered subset $\{r_0 < \ldots < r_k\} \subseteq R(G)$ such that:

1. $\epsilon(r_i) \neq \epsilon(r_{i+1})$, for $i = 0, \ldots, k-1$,
2. for $i = 0, \ldots, k-1$, there is a cycle γ of G such that $r_\gamma = r_i$ and r_{i+1} lies on γ.

We say that C is a σ-chain if $\epsilon(r_0) = \sigma$, otherwise we say that C is a π-chain. We shall write $C \sqsubset G$ if C is a chain in G, $C \sqsubset_\sigma G$ if C is σ-chain in G and $C \sqsubset_\pi G$ if C is a π-chain in G.

Definition 3.17 Let $G \in \mathcal{L}$, we define

$$\begin{aligned} L(G) &= \max\{\, \mathrm{card}\, C \mid C \sqsubset G \,\}, \\ L_\sigma(G) &= \max\{\, \mathrm{card}\, C \mid C \sqsubset_\sigma G \,\}, \\ L_\pi(G) &= \max\{\, \mathrm{card}\, C \mid C \sqsubset_\pi G \,\}. \end{aligned}$$

For every $n \geq 0$, we define the class $L_n \subseteq \mathcal{J}$ by saying that $G \in L_n$ if and only if $L(G) \leq n$. We let $S_0 = P_0 = L_0$. For every $n \geq 1$ we define the classes S_n, P_n by saying that

$$\begin{aligned} G \in S_n \quad &\text{if and only if} \quad L_\sigma(G) \leq n \ \text{ and } \ L_\pi(G) \leq n-1\,, \\ G \in P_n \quad &\text{if and only if} \quad L_\sigma(G) \leq n-1 \ \text{ and } \ L_\pi(G) \leq n\,. \end{aligned}$$

Proposition 3.18 We have equalities $L_n = \mathcal{L}_n$, for $n \geq 0$ and $S_n = \mathcal{S}_n$, $P_n = \mathcal{S}_n$, for $n \geq 1$.

Proof. We prove the proposition for $n = 0$. Observe that $G \in L_0$ if and only if every chain has cardinality less or equal to 0, i.e. the only chain is the empty set. This happens if and only if there are no returns in G, since a return r gives rise to a chain $\{r\}$. It is clear that $G \in \mathcal{L}_0$ if and only if $R(G) = \emptyset$.

Suppose that $\mathcal{S}_n = S_n$ and that $\mathcal{P}_n = P_n$.

We shall show first that $\mathcal{S}_n \cup \mathcal{P}_n \subseteq S_{n+1}$ and that S_{n+1} is closed under substitution and the μ-operation.

It is clear from the definition that $S_n \cup P_n \subseteq S_{n+1}$, so that $\mathcal{S}_n \cup \mathcal{P}_n \subseteq S_{n+1}$. The class S_{n+1} is closed under substitution: every chain in $G[H/x]$ is either a chain from $G[x]$ or a chain of H, since if $g \in G_0$ and $h \in H_0$, then there is no cycle γ of $G[H/x]$ on which both g and h lie. It is also closed under the μ-operation. Let $G[x]$ be in S_{n+1} and let $C = \{r_0, \ldots, r_k\}$ be a chain in $\mu_x.G[x]$. Observe first that if $r_0 \neq x$ then $\{r_0, \ldots, r_k\}$ is also a chain of $G[x]$. This is because a cycle γ_i such that $r_{\gamma_i} = r_i$ does not contain the transition $x \to S(x)$, otherwise $r_{\gamma_i} = x$. If C is a π-chain, then $r_0 \neq x$ so that C is a π-chain of $G[x]$ and $\operatorname{card} C \leq n$. If C is a σ-chain, we distinguish two cases: either $r_0 \neq x$, so that C is also a chain of $G[x]$ and $\operatorname{card} C \leq n+1$; or $r_0 = x$, then $\{r_1, \ldots, r_k\}$ is a π-chain in $G[x]$, so that $k \leq L_\pi(G[x]) \leq n$ and $\operatorname{card} C = k+1 \leq n+1$.

We shall now prove that if $\mathcal{C} \subseteq \mathcal{J}$ is a class such that $\mathcal{S}_n \cup \mathcal{P}_n \subseteq \mathcal{C}$ which is also closed under substitution and the μ-operation, then $S_{n+1} \subseteq \mathcal{C}$. If $G \in S_{n+1}$ we let $\zeta(G)$ be the number

$$\operatorname{card}\{\, C \sqsubset_\sigma G \,|\, \operatorname{card} C = n+1\},$$

and prove that $G \in \mathcal{C}$ by induction on $\zeta(G)$.

Suppose that $\zeta(G) = 0$. Then all chains of G have cardinality less than n so that $G \in L_n$.

Lemma 3.19 The class L_n is the closure under substitution of $S_n \cup P_n$.

Proof. Indeed, given $G \in L_n$, if we can pick up a return r which is a complete vertex distinct from the root, then we can write $G = G^{\downarrow}_r[G^r_{\downarrow}/r]$ and deduce that G belongs to the closure of $S_n \cup P_n$ under substitution by the inductive hypothesis that this property holds for both $G^{\downarrow}_r$ and $G^r_{\downarrow}$. If this is not possible, then every return of G lies on the same strongly connected component. Indeed it suffices to observe that if there is a return, then there is a unique minimal return; which, being a complete vertex, is then the root. For every other return, we can find a cycle on which both r and the root lie. Indeed, a return r is either the root, or else we can find a path γ from r to a return $r' \neq r$ which lies on γ_r. In this way we construct by induction a path from r to the root, and therefore a cycle on which both r and the root lie.

To prove the lemma, it is then enough to observe that if $G \in \mathcal{L}$ is such that $L(G) = n$ and every return of G lies on the same strongly connected component, then $G \in S_n \cup P_n$. Suppose we can find a σ-chain $\{r_0, \ldots, r_{n-1}\}$ as well as a π-chain $\{r'_0, \ldots, r'_{n-1}\}$ in G, both of cardinality n. Choose a cycle γ on which both r_0, r'_0 lie. If $\epsilon(r_\gamma) = \pi$, then $\{r_\gamma, r_0, \ldots, r_{n-1}\}$ is a π-chain of cardinality $n+1$. If $\epsilon(r_\gamma) = \sigma$, then $\{r_\gamma, r'_0, \ldots, r'_{n-1}\}$ is a σ-chain of cardinality $n+1$. This contradicts the hypothesis that $G \in L_n$. □

Using the lemma we conclude that if $G \in S_{n+1}$ and $\zeta(G) = 0$, then $G \in \mathcal{C}$, since $\mathcal{C}$ contains $\mathcal{S}_n \cup \mathcal{P}_n = S_n \cup P_n$ and is closed under substitution.

So suppose that $\zeta(G) \geq 1$. Consider the following order on σ-chains C such that $\operatorname{card} C = n + 1$:

$$\{r_0, \ldots, r_n\} \leq \{r'_0, \ldots, r'_n\} \quad \text{if and only if}$$
$$\{r_1, \ldots, r_n\} = \{r'_1, \ldots, r'_n\} \text{ and } r_0 \text{ is an ancestor of } r'_0.$$

Choose a σ-chain $C = \{r_0, \ldots, r_n\}$ which is minimal with respect to this order. We claim that r_0 is a complete vertex, i.e. we can represent G as $G^{\downarrow}_{r_0}[G^{r_0}_{\downarrow}/r_0]$, so that $G^{r_0}_{\downarrow} = \mu_{r_0}.H[r_0]$.

Suppose that r_0 is not a complete vertex. We can find a return r which is a proper ancestor of r_0 and a cycle γ on which r, r_0 lie and such that $r_\gamma = r$. If $\epsilon(r) = \pi$ then $\{r, r_0, \ldots, r_n\}$ is a π-chain in G of cardinality $n+2$, against the assumptions. If $\epsilon(r) = \sigma$, then $\{r, r_2, \ldots, r_{n+1}\}$ is a σ-chain which is strictly smaller than C in the previous order, contradicting again the hypothesis.

Since $\zeta(G^{\downarrow}_{r_0}[r_0]) < \zeta(G)$ and similarly $\zeta(H[r_0]) < \zeta(G)$, by the induction hypothesis, we have $G^{\downarrow}_{r_0}[r_0] \in \mathcal{C}$ and $H[r_0] \in \mathcal{C}$. Since $\mathcal{C}$ is closed under substitution and the μ-operation, we see that $G = G^{\downarrow}_{r_0}[\mu_{r_0}.H[r_0]/r_0] \in \mathcal{C}$.

A similar argument shows that $\mathcal{P}_{n+1} = P_{n+1}$.

By definition it is also clear that $L_n = S_{n+1} \cap P_{n+1}$, so that $\mathcal{L}_n = L_n$. This ends the proof of proposition 3.18. □

4 Free μ-lattices and free Λ_n-models

The goal of this section is to review the characterization of free μ-lattices. We shall then describe a canonical sub-Λ_n-model of a free μ-lattice $\mathcal{J}_P$

and argue that it is the free Λ_n-model over the partially ordered set P.

Definition 4.1 Let P be a partially ordered set. A *game over* P is a pair $\langle G, \lambda \rangle$ where G is a game in $\mathcal{K}$ and $\lambda : X_G \longrightarrow P$ is a valuation of the partial final positions in P. We write $\mathcal{K}(P)$ for the class of games over P and $\mathcal{J}(P)$ for the subclass of pairs $\langle G, \lambda \rangle$ such that $G \in \mathcal{J}$.

We can understand a game over P as a game with complete information with a payoff function taking values in the partially ordered set P. Player σ is trying to maximize his payoff, while his opponent π is trying to minimize the payoff; however, we can also adopt the opponent's view and think of G as a game over P^{op}, so player π is also trying to maximize the payoff but in the dual poset.

We shall use the simplified notation G for a game $\langle G, \lambda \rangle$ over P, leaving in the background the valuation $\lambda : X_G \longrightarrow P$. In particular, let $G \in \mathcal{K}$, let $\{\langle H_x, \lambda_x \rangle\}_{x \in X_G}$ be a collection of elements of $\mathcal{K}(P)$, and observe that the set $X_{G[H_x/x]_{x \in X_G}}$ is the disjoint union of the sets X_{H_x} for $x \in X_G$; the notation $G[H_x/x]_{x \in X_G}$ will abbreviate $\langle G[H_x/x]_{x \in X_G}, \lambda \rangle$, where $\lambda(y) = \lambda_x(y)$ whenever $y \in X_{H_x}$.

We describe now a preorder on the class $\mathcal{K}(P)$. This is done by constructing a game $\langle G, H \rangle$, where $G, H \in \mathcal{K}(P)$, and by saying that $G \leq H$ if one of the players, Mediator, has a winning strategy in this game, cf. 2.3. This game, which is essentially the same game described in [Bla92, Joy97], is played on the two boards G and H at the same time. One player, the one we call *Mediator* and denote by the letter M, is a team composed by player π on G and player σ on H; the other player, whom we call *the Opponents* and denote by the letter O, is formed out of player σ on G and player π on H. Mediator, in order to choose a move, must wait for the Opponents to have exhausted their moves on both boards. Mediator's goal is to reach a pair of positions $(x, y) \in X_G \times X_H$, such that $\lambda(x) \leq \lambda(y)$; in the case of an infinite play, his goal is to win on at least one board. We picture the game as follows:



The frame around Mediator's team is meant to suggest that Mediator can behave like a single player, like a master playing on different chess boards, where the Opponents are indeed two distinct players, since they do not get any advantage from sharing information. We formally define the game $\langle G, H\rangle$ as follows.

Definition 4.2 Let $G, H \in \mathcal{K}(P)$. The game $\langle G, H\rangle$ is defined as:

- Positions of $\langle G, H\rangle$ are pairs of positions from G and H. The initial position is (g_0, h_0).
- The coloring $\epsilon(g, h)$ is calculated as $\epsilon(g) \cdot \epsilon(h) \in \{?, M, O\}$, where the product is given by the table:

$\cdot$	π	σ	0
σ	O	O	O
π	O	M	M
0	O	M	$?$

  If $\epsilon(x) \cdot \epsilon(y) =?$, i.e. if $x \in X_G$ and $y \in X_H$, then $\epsilon(x, y) = O$ if and only if $\lambda(x) \leq \lambda(y)$: the pair (x, y) becomes a winning final position for Mediator exactly when $\lambda(x) \leq \lambda(y)$.

- Moves of $\langle G, H\rangle$ are either left moves $(g, h) \to (g', h)$, where $g \to g' \in G_1$, or right moves $(g, h) \to (g, h')$, where $h \to h' \in H_1$; however, the Opponents can play only with σ on G or with π on H. This means that the set of moves is obtained from the set $G_1 \times H_0 + G_0 \times H_1$ by removing right moves $(g, h) \to (g, h')$ whenever $(\epsilon(g), \epsilon(h)) = (\sigma, \sigma)$ and left moves $(g, h) \to (g', h)$ whenever $(\epsilon(g), \epsilon(h)) = (\pi, \pi)$.

  A morphism of graphs from (the graph underlying) $\langle G, H\rangle$ to the free category on (the graph underlying) G, is defined as follows: $(g, h)_G = g$ and if $\tau = (g, h) \to (g', h)$ is a left move, then $\tau_G = g \to g'$, if $\tau = (g, h) \to (g, h')$ is a right move, then $\tau_G = 1_g$. This morphism is extended to a convex functor from the free category on $\langle G, H\rangle$ to the free category on G and to a correspondence sending infinite paths in $\langle G, H\rangle$ to finite or infinite paths in G. We call all these three correspondences *left projection* and denote them by $(_)_L$ or $(_)_G$. The *right projection*, denoted by $(_)_R$ or $(_)_H$, is defined in an analogous way.

- An infinite play γ is a win for Mediator if and only if either its left projection γ_G is an infinite play and $\epsilon(r_{\gamma_G}) = \sigma$, or its right projection γ_H is an infinite play and $\epsilon(r_{\gamma_H}) = \pi$.

Definition 4.3 Let $G, H \in \mathcal{K}(P)$ be games over P. We declare that $G \leq H$ if and only if Mediator has a winning strategy in the game $\langle G, H \rangle$.

We remark that if Mediator has a winning strategy in the game $\langle G, H \rangle$, then he has also a bounded memory winning strategy, cf. 2.4. This follows from [San00a, §4] and from well known facts of the theory of games played on finite graphs [Tho97, Zie98].

Proposition 4.4 Let $G, H, K \in \mathcal{K}(P)$ be games over P. Then $G \leq G$ and if $G \leq H$ and $H \leq K$, then $G \leq K$.

Definition 4.5 Let $G, H \in \mathcal{J}(P)$. We write $G \equiv H$ if $G \leq H$ and $H \leq G$, so that $\equiv$ is an equivalence relation. We shall denote by $[G]$ the equivalence class of G and by $\mathcal{J}_P$ the set of those equivalence classes.

The following is the main achievement of [San00b].

Theorem 4.6 For every ordered set P, $\mathcal{J}_P$ is a μ-lattice, where if $G \in \mathcal{J}$ and $\{ [H_x] \}_{x \in X_G} \in \mathcal{J}_P^{X_G}$, then

$$|G|\{ [H_x] \}_{x \in X_G} \;=\; [G[H_x/x]_{x \in X_G}] \,.$$

The μ-lattice $\mathcal{J}_P$ is free over P, i.e. it comes with an embedding $\eta_P : P \longmapsto \mathcal{J}_P$ with the following universal property: if $f : P \longrightarrow L$ is an order preserving function from P to a μ-lattice L, then there exists a unique morphism of μ-lattices $\tilde{f}$ such that $\tilde{f} \circ \eta_P = f$.

We shall review here the concepts developed to prove proposition 4.4, since we will need them later.

We proved that $G \leq G$ by exhibiting the *copycat strategy* in $\langle G, G \rangle$. It is played as follows. From a position of the form (g, g) it is always the case that the Opponents have to move just on one board. When they stop moving, if they do, Mediator will have the opportunity to copy all

the moves played by the Opponents so far on the other board until the play again reaches a position of the form (g', g').

We proved that if $G \leq H$ and $H \leq K$ then $G \leq K$, by describing a game $\langle G, H, K\rangle$ with the two following properties: given two winning strategies R and S on $\langle G, H\rangle$ and $\langle H, K\rangle$ there exists a winning strategy $R||S$ on $\langle G, H, K\rangle$; given a winning strategy T on $\langle G, H, K\rangle$ there exists a winning strategy $T_{\backslash H}$ on $\langle G, K\rangle$. The game $\langle G, H, K\rangle$ is obtained by gluing the games $\langle G, H\rangle$ and $\langle H, K\rangle$ along the center board H. One player, whom we call *the Mediators* and denote by the letter M, is a team composed by Mediator on $\langle G, H\rangle$ and Mediator on $\langle H, K\rangle$; the other player, called *the Opponents* and denoted by the letter O, is formed out of player σ on G and player π on K. The Mediators can exchange information through the center board H. The game can be pictured as follows:



Definition 4.7 Let $G, H, K \in \mathcal{K}(P)$. The game $\langle G, H, K\rangle$ is defined as follows:

- Positions of $\langle G, H, K\rangle$ are triples of positions $(g, h, k) \in G_0 \times H_0 \times K_0$. The initial position is (g_0, h_0, k_0).
- The coloring $\epsilon(g, h, k)$ is calculated as $\epsilon(g) \cdot \epsilon(k) \in \{?, O, M\}$, where the product is given by the table of page 21. If $\epsilon(x) \cdot \epsilon(z) =?$, i.e. if $x \in X_G$ and $z \in X_K$, then if $\epsilon(h) \neq 0$, then $\epsilon(x, h, z) = M$, otherwise, if also $h = y \in X_H$, then $\epsilon(x, y, z) = O$ if and only if $\lambda(x) \leq \lambda(y) \leq \lambda(z)$.
- Moves of $\langle G, H, K\rangle$ are either left moves $(g, h, k) \to (g', h, k)$, where $g \to g' \in G_1$, or central moves $(g, h, k) \to (g, h', k)$, where $h \to h' \in H_1$, or right moves $(g, h, k) \to (g, h, k')$, where $k \to k' \in K_1$; however, the Opponents can play only with σ on G or with π on K. Several kind of projections on subsets of the three boards can be defined as in definition 4.2; they will be denoted $(_)_G$, $(_)_H$, $(_)_K$, $(_)_{\langle G,H\rangle}$, $(_)_{\langle H,K\rangle}$, $(_)_{\langle G,K\rangle}$. Observe that for every path γ, finite or infinite, $(\gamma_{\langle G,H\rangle})_G = (\gamma_{\langle G,K\rangle})_G = \gamma_G$ and similar equalities hold in the other cases.

- An infinite play γ is a win for the Mediators if and only if its left projection γ_G is an infinite play and $\epsilon(r_{\gamma_G}) = \sigma$, or its right projection γ_K is an infinite play and $\epsilon(r_{\gamma_K}) = \pi$.

Definition 4.8 Let R be a winning strategy for Mediator in $\langle G, H\rangle$ and let S be a winning strategy for Mediator in $\langle H, K\rangle$. A strategy $R||S$ for the Mediators in the game $\langle G, H, K\rangle$ is described by the formula

$$R||S \;=\; \{\, \gamma \,|\, \gamma_{\langle G,H\rangle} \in R \text{ and } \gamma_{\langle H,K\rangle} \in S \,\}\,.$$

Let T be a winning strategy for the Mediators in the game $\langle G, H, K\rangle$. A strategy $T_{\backslash H}$ for Mediator in the game $\langle G, H, K\rangle$ is defined by the formula

$$T_{\backslash H} \;=\; \{\, \gamma_{\langle G,K\rangle} \,|\, \gamma \in T \,\}\,.$$

Proposition 4.9 The strategy $R||S$ is a winning strategy for the Mediators in the game $\langle G, H, K\rangle$.

Proof. We check that all the conditions of definition 2.3 are satisfied.

If $\gamma \in R||S$ then $\gamma_{\langle G,H\rangle} \in R$ and $\gamma_{\langle H,K\rangle} \in S$, so that $\mathrm{dom}\,\gamma_{\langle G,H\rangle} = (g_0, h_0)$ and $\mathrm{dom}\,\gamma_{\langle H,K\rangle} = (h_0, k_0)$. Let $\mathrm{dom}\,\gamma = (g, h, k)$, then $(g, h) = (\mathrm{dom}\,\gamma)_{\langle G,H\rangle} = \mathrm{dom}\,\gamma_{\langle G,H\rangle} = (g_0, h_0)$, similarly $(h, k) = (h_0, k_0)$, so that $\mathrm{dom}\,\gamma = (g, h, k) = (g_0, h_0, k_0)$.

Suppose that $\gamma \star \tau \in R||S$. From $\gamma_{\langle G,H\rangle} \star \tau_{\langle G,H\rangle} = (\gamma \star \tau)_{\langle G,H\rangle} \in R$, we argue that $\gamma_{\langle G,H\rangle} \in R$; similarly $\gamma_{\langle H,K\rangle} \in S$, so that $\gamma \in R||S$.

Let γ be a path in $R||S$ and let $(g, h, k) = \mathrm{cod}\,\gamma$.

Firstly, let us assume that $\epsilon(g, h, k) = O$. If τ is a move available to the Opponents from this position, then it is either a left move or a right move. Suppose it is a left move, say $\tau = (g, h, k) \to (g', h, k)$. We deduce that $\epsilon(g) = \sigma$, which in turn implies that $\epsilon(g, h) = O$. Since $\gamma_{\langle G,H\rangle} \in R$ and $\mathrm{cod}\,\gamma_{\langle G,H\rangle} = (g, h)$, then the path $\gamma_{\langle G,H\rangle} \star \tau_{\langle G,H\rangle} \in R$. We deduce that $\gamma \star \tau \in R||S$, since $(\,\gamma \star \tau\,)_{\langle G,H\rangle} = \gamma_{\langle G,H\rangle} \star \tau_{\langle G,H\rangle} \in R$ and $(\,\gamma \star \tau\,)_{\langle H,K\rangle} = \gamma_{\langle H,K\rangle} \in S$. If τ is a right move, then a similar argument shows that $\gamma \star \tau \in R||S$.

Suppose now that $\epsilon(g, h, k) = M$. If $\epsilon(h) = 0$ then either $\epsilon(g) = \pi$ or $\epsilon(k) = \sigma$, suppose $\epsilon(g) = \pi$. In this case $\epsilon(g, h) = M$, and we can find a transition τ of $\langle G, H\rangle$ such that $\gamma_{\langle G,H\rangle} \star \tau \in R$. Since $\tau = (g, h) \to (g', h)$

is a left move, we can lift it to a left move $\tau' = (g, h, k) \to (g', h, k)$, so that $\gamma \star \tau' \in R||S$ since $(\gamma \star \tau')_{\langle G,H\rangle} = \gamma_{\langle G,H\rangle} \star \tau \in R$ and $(\gamma \star \tau')_{\langle H,K\rangle} = \gamma_{\langle H,K\rangle} \in S$. We can reason similarly if $\epsilon(k) = \sigma$.

Suppose that $\epsilon(h) \in \{\sigma, \pi\}$, say $\epsilon(h) = \sigma$. In this case $\epsilon(g, h) = M$, and we can find a transition τ of $\langle G, H\rangle$ such that $\gamma_{\langle G,H\rangle} \star \tau \in R$. If τ is a left move, then we lift it to τ' as before and conclude that $\gamma \star \tau' \in R||S$. Hence suppose that $\tau = (g, h) \to (g, h')$ is a right move. Since $\epsilon(h) = \sigma$, we deduce that $\epsilon(h, k) = O$, so that $\gamma_{\langle H,K\rangle} \star \tilde{\tau} \in S$, where $\tilde{\tau} = (h, k) \to (h', k)$. If we set $\tau' = (g, h, k) \to (g, h', k)$, we deduce that $\gamma \star \tau' \in S||R$, since $(\gamma \star \tau')_{\langle G,H\rangle} = \gamma_{\langle G,H\rangle} \star \tau \in R$ and $(\gamma \star \tau')_{\langle H,K\rangle} = \gamma_{\langle H,K\rangle} \star \tilde{\tau} \in S$. We can reason similarly if $\epsilon(h) = \pi$.

Suppose now that $(g, h, k) = (x, y, z)$, $x \in X_G, y \in X_H, z \in X_K$. Since $\operatorname{cod} \gamma_{\langle G,H\rangle} = (x, y)$, we deduce that $\epsilon(x, y) = O$, so that $\lambda(x) \leq \lambda(y)$. Since $\operatorname{cod} \gamma_{\langle H,K\rangle} = (y, z)$, we deduce that $\epsilon(y, z) = O$, so that $\lambda(y) \leq \lambda(z)$.

Consider an infinite play γ in $\langle G, H, K\rangle$ which is the result of playing in this way. Either $\gamma_{\langle G,H\rangle}$ is an infinite play, or $\gamma_{\langle H,K\rangle}$ is an infinite play; we suppose the first. If $\epsilon(\gamma_G)$ is not an infinite winning play for player π in G, then γ_H is an infinite winning play for σ on H: indeed, the pair (γ_G, γ_H) is the left and right projection of the infinite play $\gamma_{\langle G,H\rangle}$, which has been played according to the winning strategy R. We can then argue that $\gamma_{\langle H,K\rangle}$ is also an infinite play, moreover it has been played according to the winning strategy S. Since γ_H is not an infinite winning play for π on H, and the pair (γ_H, γ_K) is the left and right projection of $\gamma_{\langle H,K\rangle}$, it follows that γ_K is an infinite winning play for σ on K. A similar argument is used if $\gamma_{\langle H,K\rangle}$ is an infinite play. □

Proposition 4.10 The strategy $T_{\backslash H}$ is a winning strategy for Mediator in the game $\langle G, K\rangle$.

Proof. Let $\gamma \in T$, then $\operatorname{dom} \gamma = (g_0, h_0, k_0)$, so that $\operatorname{dom} \gamma_{\langle G,K\rangle} = (\operatorname{dom} \gamma)_{\langle G,K\rangle} = (g_0, k_0)$. Similarly, if $\gamma_{\langle G,K\rangle} = \gamma_1 \star \tau$, then we can find a factorization $\gamma = \gamma'_1 \star \gamma'_2$ such that $(\gamma'_1)_{\langle G,K\rangle} = \gamma_1$ and $(\gamma'_2)_{\langle G,K\rangle} = \tau$. It follows that $\gamma_1 \in T_{\backslash H}$.

Choose a play $\gamma \in T$ and suppose that $\operatorname{cod} \gamma = (g, h, k)$, so that $\operatorname{cod} \gamma_{\langle G,K\rangle} = (g, k)$.

Suppose that $\epsilon(g, k) = O$ and let τ be a move available to the Opponents from (g, k). If $\tau = (g, h) \to (g', h)$ is a left move, then $\epsilon(g) = \sigma$, and

therefore $\epsilon(g,h,k) = O$. The transition $\tau' = (g,h,k) \to (g',h,k)$ is a move of $\langle G,H,K\rangle$, hence $\gamma \star \tau' \in T$ and $\gamma_{\langle G,K\rangle} \star \tau = \gamma_{\langle G,K\rangle} \star \tau'_{\langle G,K\rangle} = (\gamma \star \tau')_{\langle G,K\rangle} \in T_{\backslash H}$.

Suppose that $\epsilon(g,k) = M$, and observe that $\epsilon(g,h,k) = M$; we must find a transition τ such that $\gamma_{\langle G,K\rangle} \star \tau \in T_{\backslash H}$. Suppose also that we have constructed paths $\{\gamma_i\}_{i=0,\ldots,n}$ with the following properties: $|\gamma_j| = j$, γ_i is a prefix of γ_j if $i \leq j$, $\gamma \star \gamma_i \in T$ and $(\gamma \star \gamma_i)_{\langle G,K\rangle} = \gamma_{\langle G,K\rangle}$. Since $\epsilon(\operatorname{cod}\gamma \star \gamma_n) = M$, there exists a transition τ such that $\gamma \star \gamma_n \star \tau \in T$. If τ is a left or right transition, then $\tau_{\langle G,K\rangle}$ is a transition of $\langle G,K\rangle$, so that $\gamma_{\langle G,K\rangle} \star \tau_{\langle G,K\rangle} \in T_{\backslash H}$, and we are done. If τ is a central transition, then we extend the above collection by letting $\gamma_{n+1} = \gamma_n \star \tau$. Since we cannot build an infinite collection $\{\gamma_n\}_{n\geq 0}$ with the above properties – the infinite path $\{\gamma \star \gamma_n\}_{n\geq 0}$ is played according to the winning strategy T, but it is not a win for the Mediators in $\langle G,H,K\rangle$ – we shall eventually find $n \geq 0$ and a right or left transition τ such that $\gamma \star \gamma_n \star \tau \in T$.

Finally, consider an infinite path $\gamma = \{\gamma_n\}_{n\geq 0}$ played according to the strategy $T_{\backslash H}$. Consider the set of paths

$$T(\gamma) \;=\; \{\, \gamma' \in T \mid \gamma'_{\langle G,K\rangle} \in \{\gamma_n\}_{n\geq 0} \,\}\,.$$

The set $T(\gamma)$ is closed under prefixes and it is infinite. Hence, it has the structure of a finitely branching infinite tree and we can find an infinite path $\gamma' = \{\gamma'_k\}$ on this tree, which is a subtree of T, such that $\gamma'_{\langle G,K\rangle} = \gamma$. Since T is a winning strategy, we have that $\epsilon(r_{\gamma_G}) = \epsilon(r_{\gamma'_G}) = \sigma$ or $\epsilon(r_{\gamma_K}) = \epsilon(r_{\gamma'_K}) = \sigma$.

This concludes the proof of proposition 4.4. □

Definition 4.11 For every $[G] \in \mathcal{J}_P$, we define $L[G]$ to be the number

$$\min\{\, n \mid L(H) = n\,, H \in [G]\}$$

and let

$$\mathcal{J}_{n,P} \;=\; \{\, [G] \mid L[G] \leq n \,\}\,.$$

Proposition 4.12 The set $\mathcal{J}_{n,P}$ is a sub-Λ_n-model of $\mathcal{J}_P$ and the embedding $\eta_P : P \longmapsto \mathcal{J}_P$ restricts to an embedding $\eta_{n,P} : P \longmapsto \mathcal{J}_{n,P}$. With this structure $\mathcal{J}_{n,P}$ is free over of P, i.e. the above embedding has the usual universal property with respect to order preserving functions with codomain a Λ_n-model.

Proof. Let G be a game in $\mathcal{L}_n$ and $\{\,[H_x]\,\}_{x\in X_G}$ be a collection of elements in $\mathcal{J}_{n,P}$; we must prove that $|G|\{\,[H_x]\,\}_{x\in X_G} \in \mathcal{J}_{n,P}$. For each $x \in X_G$ choose $H'_x \in [H_x]$ such that $L(H'_x) \leq n$, so that $[H_x] = [H'_x]$. The equalities

$$\begin{aligned} |G|\{\,[H_x]\,\}_{x\in X_G} &= |G|\{\,[H'_x]\,\}_{x\in X_G} \\ &= [G[H'_x/x]_{x\in X_G}] \end{aligned}$$

shows that $|G|\{\,[H_x]\,\}_{x\in X_G} \in \mathcal{J}_{n,P}$, since $G \in \mathcal{L}_n$, $H'_x \in \mathcal{L}_n$ and $\mathcal{L}_n$ is closed under substitution. Since $\eta_P(p) = [x, \lambda^p]$, where $\lambda^p(x) = p$, it is clear that $\eta_P(p) \in \mathcal{J}_{n,P}$.

As a Λ_n-model, $\mathcal{J}_{n,P}$ is isomorphic to $\mathcal{L}_{n,P}$, the antisymmetric quotient of the preordered class $\mathcal{L}_n(P)$ of pairs $\langle G, \lambda\rangle$ with $G \in \mathcal{L}_n$, and we shall prove freeness of $\mathcal{L}_{n,P}$. Observe first that the correspondence $\langle G, \lambda\rangle \longmapsto \langle G, f \circ \lambda\rangle$, induced by an ordered preserving function $f : P \longrightarrow Q$, induces an order preserving correspondence $\mathcal{L}_n(f) : \mathcal{L}_n(P) \longrightarrow \mathcal{L}_n(Q)$, and a morphism of Λ_n-models $\mathcal{L}_{n,f} : \mathcal{L}_{n,P} \longrightarrow \mathcal{L}_{n,Q}$. This makes up a functor $\mathcal{L}_n$ and η_n is then a natural transformation in the obvious sense.

On the other hand, if L is a Λ_n-model, then the correspondence $EV_n(L) : \mathcal{L}_n(L) \longrightarrow L$, defined by $EV_n(L)\langle G, \lambda\rangle = |G|(\lambda)$, preserves also the order, so that it induces an morphism of Λ_n-models $EV_L : \mathcal{L}_{n,L} \longrightarrow L$ such that $EV_{n,L} \circ \eta_{n,L} = Id_L$. To prove this, the same argument as in [San00a, §5.15] is used.

If $f : P \longrightarrow L$ is an order preserving function with codomain a Λ_n-model, then $EV_{n,L} \circ \mathcal{L}_{n,f}$ is the desired unique extension of f to a morphism of Λ_n-models from $\mathcal{L}_{n,P}$ to L. □

We shall denote by $i_{n,P} : \mathcal{J}_{n,P} \hookrightarrow \mathcal{J}_P$ the inclusion, so that $i_{n,P}$ is a morphism of Λ_n-models and the equality $\eta_P = i_{n,P} \circ \eta_{n,P}$ holds.

5 Synchronizing games

The goal of this section is to give a general criterion by which to prove that the inclusion $i_{n,P} : \mathcal{J}_{n,P} \hookrightarrow \mathcal{J}_P$ is proper.

Definition 5.1 Let $A \in \mathcal{K}(P)$. We say that A is *synchronizing* if it is bipartite and the only winning strategy for Mediator in the game $\langle A, A\rangle$ is the copycat strategy.

The intuitions which have induced us to call these games synchronizing are explained as follows: if Mediator is playing according to a winning strategy in the game $\langle G, H\rangle$, then it is impossible for both the Opponents to win, so that at least one must lose. We can imagine therefore that there is a sort of asynchronous game going on between player σ on G and player π on H, the asynchrony being induced by the mediating choices of Mediator. However, if $G = H = A$ and the only winning strategy for Mediator in $\langle A, A\rangle$ is the copycat strategy, then there are very few mediating choices. If moreover the game A is bipartite, then the resulting game between player σ on the left and player π on the right is easily recognized to be equivalent with the game A itself, in which the two players act on a synchronous base.

Since a strategy in the game $\langle A, A\rangle$ is morally an endomorphism of A, and the copycat strategy plays the role of the identity, we may consider a synchronizing game to be a particular kind of asymmetric object. We remark that if $A \in \mathcal{J}(P)$ is such that $L(A) = 0$, then A can be identified with a term for the the free lattice over P. In this case, A is synchronizing if and only if it is in normal form as a free lattice term [FJN95, Whi41]. It is an open problem whether this notion of synchronizing game leads to a normal form for μ-lattice terms. Examples of synchronizing games are given in section 6. Their remarkable property is stated in the following proposition.

Theorem 5.2 Let $A \in \mathcal{K}(P)$ be a synchronizing game such that $L(A) = n$. Then we can construct a game $A_\bullet \in \mathcal{J}(P)$ such that $L[A_\bullet] = n$.

Corollary 5.3 In order to show that the inclusion $i_{n,P} : \mathcal{J}_{n,P} \hookrightarrow \mathcal{J}_P$ is proper, it is enough to find a synchronizing game A in $\mathcal{K}(P)$ such that $L(A) > n$.

Proof. In the following let $A = \langle A_0, A_1, a_0, \epsilon, \lambda\rangle$ be such a synchronizing game.

The game $A_\bullet$ is obtained from A by forcing property 2 in definition 3.10 to hold. If we let $A_\bullet$ be $\langle A_{\bullet 0}, A_{\bullet 1}, a_{\bullet 0}, \epsilon_\bullet, \lambda_\bullet\rangle$, this game is formally defined as follows.

- The set of positions is

$$A_{\bullet 0} \;=\; R(A) \times \{\, 0 \,\} \;\cup\; A_0 \times \{\, 1 \,\}\,.$$

The initial position $a_{\bullet 0}$ is $(a_0, 0)$ if a_0 is in $R(A)$, otherwise the initial position is $(a_0, 1)$.

- The set of moves is

$$\begin{aligned} A_{\bullet 1} \;=\; & \{\, (g,1) \to (g',1) \,|\, g \to g' \in A_1,\ g' \notin R(A) \,\} \\ & \cup\ \{\, (g,1) \to (r,0) \,|\, g \to r \in A_1,\ r \in R(A) \,\} \\ & \cup\ \{\, (r,0) \to (r,1) \,|\, r \in R(A) \,\}\,. \end{aligned}$$

- We let $\epsilon_\bullet(g,i) = \epsilon(g)$, and observe that if $\epsilon_\bullet(g,i) = 0$, then $i = 1$ and $\epsilon(g) = 0$; hence, we define $\lambda_\bullet(x,i) = \lambda(x)$ if $\epsilon_\bullet(x,i) = 0$.

Observe that this construction preserves the essential structure of cycles and the color of the returns, from which we deduce that $L(A) = L(A_\bullet)$. It is easily seen that $A_\bullet \in \mathcal{J}(P)$, so that $L[A_\bullet] \leq L(A_\bullet) = L(A) = n$.

In order to argue that $L[A_\bullet] \geq n$, choose an arbitrary $H \in [A_\bullet]$. A copycat-like strategy can be used by Mediator to win in both the games $\langle A, A_\bullet \rangle$ and $\langle A_\bullet, A \rangle$, so that, by transitivity, we obtain $A \leq H$ and $H \leq A$. By the analysis of possible plays in the game $\langle A, H, A \rangle$, we shall construct a chain C in H such that $\operatorname{card} C = n$; it will follow that $L(H) \geq n$ and $L[A_\bullet] \geq n$. If θ is a path in $\langle A, H, A \rangle$ or $\langle A, A \rangle$, we shall use θ_L and θ_R for the left and right projections, since the notation θ_A would be ambiguous.

The lemma 5.4 below has the following interpretation: *if the Opponents know that the Mediators are playing according to a winning strategy, then they can choose a path γ of A and force the Mediators to play on $\langle A, H, A \rangle$ so that the chosen path is played on the left board as well as on the right board.* In this case, we informally say that *the Opponents play along the path γ*. We let T be any winning strategy for the Mediators in the game $\langle A, H, A \rangle$.

Lemma 5.4 Let γ be a path of A and let $a = \operatorname{dom} \gamma$. Suppose that there is a position h of H such that the position (a, h, a) of $\langle A, H, A \rangle$ has been reached using T. We can lift γ to a play θ_γ of $\langle A, H, A \rangle$ with the following properties:

- $\operatorname{dom} \theta_\gamma = (a, h, a)$,
- the left projection $(\,\theta_\gamma\,)_L$ as well as the right projection $(\,\theta_\gamma\,)_R$ are both equal to γ,

- the play θ_γ has been played according to the strategy T by the Mediators.

Proof. Firstly, we prove the lemma in case $\gamma = a \to a'$ is a transition of A; we also suppose that $\epsilon(a) = \sigma$; if $\epsilon(a) = \pi$ we can reason by duality. From position (a, h, a) it is the Opponents' turn to move, on the left, so that they can choose the move τ on the left.

Since A is bipartite, we have $\epsilon(a') \neq \epsilon(a)$, and in position (a', h, a) it is the Mediators' turn to move. From this position, the strategy T will suggest to play a finite path on H $(a', h, a) \to^* (a', h', a)$, possibly of zero length, and then it will suggest to play on an external board. An infinite path played only on H cannot arise, since T is a winning strategy, and such an infinite path would not be a win for the Mediators. T cannot suggest a move on the left board - otherwise the strategy $T_{\backslash H}$ would not be the copycat strategy - hence it will suggest a move on the right board. Since $T_{\backslash H}$ is the copycat strategy, the only suggested move will be $(a', h', a) \to (a', h', a')$.

The generalization of the statement to paths is obtained by induction on the length. If $|\gamma| = 0$, then we lift γ to $1_{(a,h,a)}$. If $\gamma = \gamma' \star \tau$, then we define θ_γ to be $\theta_{\gamma'} \star \theta_\tau$, where $\theta_{\gamma'}$ is obtained by the induction hypothesis and θ_τ is obtained as in the previous paragraph from position $\operatorname{cod} \theta_{\gamma'}$, observing that $(\operatorname{cod} \theta_{\gamma'})_L = \operatorname{dom} \tau = (\operatorname{cod} \theta_{\gamma'})_R$. □

Lemma 5.5 Let θ be a path of $\langle A, H, A \rangle$, which has been played according to the winning strategy T, such that $\theta_L = \theta_R$. If $\gamma = \theta_L = \theta_R$ has a factorization $\gamma = \gamma_1 \star \gamma_2$, then there exists a factorization $\theta = \theta_1 \star \theta_2$ such that $(\theta_1)_L = (\theta_1)_R = \gamma_1$ and $(\theta_2)_L = (\theta_2)_R = \gamma_2$.

Proof. We shall prove that if δ is a path of $\langle A, A \rangle$, which has been played according to the winning strategy $T_{\backslash H}$, i.e according to the copycat strategy, for which the equalities $\delta_L = \delta_R = \gamma$ hold, then we can lift a factorization $\gamma = \gamma_1 \star \gamma_2$ to a factorization $\delta = \delta_1 \star \delta_2$ in $\langle A, A \rangle$ such that $(\delta_i)_L = (\delta_i)_R = \gamma_i$, for $i = 1, 2$. To obtain the statement of the lemma, it will be enough to let $\delta = \theta_{\langle A,A \rangle}$, and then lift the factorization $\theta_{\langle A,A \rangle} = \delta_1 \star \delta_2$ to a factorization $\theta = \theta_1 \star \theta_2$, which is possible since the functor $(_)_{\langle A,A \rangle}$ is convex.

The statement is proved by induction on the length of γ_2. If $|\gamma_2| = 0$, the result is obvious; suppose therefore that $|\gamma_2| > 0$. Since $|\delta| = |\delta_L| + |\delta_R| =$

$2|\gamma| \geq 2$, we can write $\delta = \delta' \star \tau'$ where $|\tau'| = 2$. Since $T_{\backslash H}$ is the copycat strategy and A is bipartite, we deduce that $\delta'_L = \delta'_R = \gamma'$, $\tau'_L = \tau'_R = \tau$ so that $|\tau| = 1$, and write $\gamma = \gamma' \star \tau$. Let $\gamma_2 = \gamma'_2 \star \tau$ and $\gamma' = \gamma_1 \star \gamma'_2$. Since $|\gamma'_2| < |\gamma_2|$, we can use the induction hypothesis and let $\delta' = \delta_1 \star \delta'_2$ be such that $(\delta_1)_L = (\delta_1)_R = \gamma_1$ and $(\delta'_2)_L = (\delta'_2)_R = \gamma'_2$. Then $\delta = \delta'_1 \star (\delta'_2 \star \tau')$ is the desired factorization. □

The lifting property of the previous lemma can be generalized: if $\theta_L = \theta_R = \gamma_1 \star \ldots \star \gamma_n$, where $n \geq 0$, then we can find a factorization $\theta = \theta_1 \star \ldots \star \theta_n$ such that $(\theta_i)_L = (\theta_i)_R = \gamma_i$ for $i = 1, \ldots, n$.

We have observed that $A \leq H$ and $H \leq A$, hence we shall fix two winning bounded memory strategies for Mediator in the games $\langle A, H \rangle$ and $\langle H, A \rangle$, say (R, ψ) and (S, ϕ), respectively. We let

$$K \quad = \quad \max(\operatorname{card} R_0, \operatorname{card} S_0)\,,$$

and consider the strategy $\psi R || \phi S$ in the game $\langle A, H, A \rangle$, the definition of which is found in 2.5 and 4.8.

The following lemma can be interpreted as follows: *if the Opponents play enough time along a cycle γ of A, then they can force the Mediators to play in a cycle of H of the same color as γ.*

Lemma 5.6 Let γ be a proper cycle of A and let θ be a path of $\langle A, H, A \rangle$, played according to the strategy $\psi R || \phi S$, such that $\theta_L = \theta_R = \gamma^K$. It is possible to find a factorization

$$\theta \quad = \quad \Theta_0 \star \Theta \star \Theta_1$$

such that Θ_L, Θ_H and Θ_R are all proper cycles. Moreover, $\Theta_L = \Theta_R = \gamma^k$, with $1 \leq k \leq K$, and $\epsilon(r_{\Theta_H}) = \epsilon(r_\gamma)$.

Proof. Let θ be a path of $\langle A, H, A \rangle$, played according to the strategy $\psi R || \phi S$, such that $\theta_L = \theta_R = \gamma^K$, where γ is proper cycle of A such that $\epsilon(r_\gamma) = \pi$; if $\epsilon(r_\gamma) = \sigma$, a dual argument – with the strategy ϕS instead of ψR – can be used. According to lemma 5.5, we can factor θ as

$$\theta \quad = \quad \theta_1 \star \ldots \star \theta_K\,,$$

so that, for each $i = 1, \ldots, K$, the relations $(\theta_i)_L = (\theta_i)_R = \gamma$ hold.

By the definition of the strategies ψR and $\psi R||\phi S$, there exists a path ρ in R such that $\psi \circ \rho = \theta_{\langle A,H\rangle}$. Consider the factorization in $\langle A, H\rangle$

$$\begin{aligned}\psi \circ \rho &= \theta_{\langle A,H\rangle} \\ &= (\theta_1 \star \ldots \star \theta_K)_{\langle A,H\rangle} \\ &= (\theta_1)_{\langle A,H\rangle} \star \ldots \star (\theta_K)_{\langle A,H\rangle}\,.\end{aligned}$$

Since the functor $\psi \circ _$ is convex, we can lift this factorization to a factorization in R

$$\rho \;=\; \rho_1 \star \ldots \star \rho_K$$

such that $\psi \circ \rho_i = (\theta_i)_{\langle A,H\rangle}$, for $i = 1, \ldots, K$. Consider also the set

$$\{\, \mathrm{dom}\,\rho_i\,,\ \mathrm{cod}\,\rho_i \,|\, i = 1, \ldots, K \,\}\,,$$

and observe that there exist $i_0, i_1 \in \{1, \ldots, K\}$ such that $i_0 \leq i_1$ and $\mathrm{dom}\,\rho_{i_0} = \mathrm{cod}\,\rho_{i_1}$, since $\mathrm{card}\,R_0 \leq K$. Let

$$\Upsilon \;=\; \rho_{i_0} \star \ldots \star \rho_{i_1},$$

then Υ is a cycle, and if we let $k = i_1 - i_0 + 1$, then

$$\begin{aligned}(\psi \circ \Upsilon)_L &= (\psi \circ (\,\rho_{i_0} \star \ldots \star \rho_{i_1}\,))_L \\ &= (\,(\psi \circ \rho_{i_0}) \star \ldots \star (\psi \circ \rho_{i_1})\,)_L \\ &= (\,(\theta_{i_0})_{\langle A,H\rangle} \star \ldots \star (\theta_{i_1})_{\langle A,H\rangle}\,)_L \\ &= ((\theta_{i_0})_{\langle A,H\rangle})_L \star \ldots \star ((\theta_{i_1})_{\langle A,H\rangle})_L \\ &= (\theta_{i_0})_L \star \ldots \star (\theta_{i_1})_L \\ &= \gamma^k\,.\end{aligned}$$

Observe also that $r_{(\psi\circ\Upsilon)_L} = r_{\gamma^k} = r_\gamma$ and that $\epsilon(r_\gamma) = \pi$. Since ψR is a winning strategy, we argue that $(\,\psi \circ \Upsilon\,)_H$ is a proper cycle and that $\epsilon(r_{(\,\psi\circ\Upsilon)_H}) = \pi$. Otherwise Υ would give rise, by infinite iteration, to the infinite path Υ^ω in the graph R such that $\psi \circ \Upsilon^\omega = (\,\psi \circ \Upsilon\,)^\omega$ is not a win for Mediator in the game $\langle A, H\rangle$.

In order to conclude the argument, let

$$\Theta \;=\; \theta_{i_0} \star \ldots \star \theta_{i_1}\,,$$

and find Θ_0, Θ_1 so that the relation $\theta = \Theta_0 \star \Theta \star \Theta_1$ is satisfied. Then $\Theta_L = \Theta_R = \gamma^k$ with $k \geq 1$ by construction. We have also

$$\begin{aligned}
\Theta_H &= (\Theta_{\langle A,H\rangle})_H \\
&= (\,(\theta_{i_0})_{\langle A,H\rangle} \star \ldots \star (\theta_{i_1})_{\langle A,H\rangle}\,)_H \\
&= (\,(\psi \circ \rho_{i_0}) \star \ldots \star (\psi \circ \rho_{i_1})\,)_H \\
&= (\psi \circ \Upsilon)_H\,,
\end{aligned}$$

so that we can conclude that Θ_H is a proper cycle such that $\epsilon(r_{\Theta_H}) = \pi = \epsilon(r_\gamma)$. □

The following lemma can be interpreted as follows: *if the Opponents play enough time along a chain C of A, then they can force the Mediators to play along a chain of H of the same length and color as C.*

Lemma 5.7 Let $C = \{a_0, \ldots, a_{n-1}\}$ be a chain in A. For $j = 0, \ldots, n-2$, let γ_j be proper cycles such that $r_{\gamma_j} = a_j$ and a_{j+1} lie on γ_j. Factor γ_j as $\gamma_j = \gamma_j^{\downarrow} \star \gamma_j^{\uparrow}$, where $\operatorname{dom}\gamma_j^{\downarrow} = a_j$ and $\operatorname{cod}\gamma_j^{\downarrow} = a_{j+1}$. Similarly, let γ_{n-1} be a proper cycle of A such that $r_{\gamma_{n-1}} = a_{n-1} = \operatorname{dom}\gamma_{n-1}$. For $j = n-1, \ldots, 0$, define cycles Γ_j in A as follows:

$$\begin{aligned}
\Gamma_{n-1} &= \gamma_{n-1}^K\,, \\
\Gamma_{j-1} &= (\gamma_{j-1}^{\downarrow} \star \Gamma_j \star \gamma_{j-1}^{\uparrow})^K\,.
\end{aligned}$$

Let θ be a path in $\langle A, H, A\rangle$, played according to the strategy $\psi R || \phi S$, such that $\theta_L = \theta_R = \Gamma_j$. We can find a factorization

$$\theta = \Theta_0 \star \Theta \star \Theta_1,$$

with the property that Θ_H is a proper cycle visiting a chain $\{r_{\Theta_H} = r_j, \ldots, r_{n-1}\}$. Moreover $\epsilon(r_j) = \epsilon(a_j)$.

Proof. We prove the proposition by induction on $j = n-1, \ldots, 0$.

If $j = n-1$, apply lemma 5.6 to θ and γ_{n-1}: it is possible to find a factorization

$$\theta = \Theta_0 \star \Theta \star \Theta_1$$

such that Θ_H is a proper cycle and $\epsilon(r_{\Theta_H}) = \epsilon(r_{\gamma_{n-1}}) = \epsilon(a_{n-1})$.

Suppose that we have proven the assertion for j; we prove it for $j-1$. Let θ be a path such that $\theta_L = \theta_R = (\gamma_{j-1}^{\downarrow} \star \Gamma_j \star \gamma_{j-1}^{\uparrow})^K$. Apply lemma 5.6 to θ and $\gamma_{j-1}^{\downarrow} \star \Gamma_j \star \gamma_{j-1}^{\uparrow}$ and find a factorization

$$\theta \;=\; \Theta_0 \star \Theta \star \Theta_1$$

such that Θ_H is a proper cycle and $\epsilon(r_{\Theta_H}) = \epsilon(a_{j-1})$, since $r_{(\gamma_{j-1}^{\downarrow} \star \Gamma_j \star \gamma_{j-1}^{\uparrow})} = r_{\gamma_{j-1}} = a_{j-1}$. Moreover, there exists $k \geq 1$ such that

$$\Theta_L = \Theta_R \;=\; (\,\gamma^{k-1} \star \gamma_{j-1}^{\downarrow}\,) \star \Gamma_j \star \gamma_{j-1}^{\uparrow}\,,$$

where $\gamma = \gamma_{j-1}^{\downarrow} \star \Gamma_j \star \gamma_{j-1}^{\uparrow}$. By lemma 5.5, we can lift the above factorization to a factorization

$$\Theta \;=\; \delta_0 \star \delta \star \delta_1\,,$$

so that $\delta_L = \delta_R = \Gamma_j$. Using the induction hypothesis, there is a factorization

$$\delta \;=\; \Delta_0 \star \Delta \star \Delta_1\,,$$

such that Δ_H is a proper cycle visiting a chain $\{\, r_{\Delta_H} = r_j, \ldots, r_{n-1} \,\}$; moreover $\epsilon(r_j) = \epsilon(a_j)$. Since

$$\Theta_H \;=\; (\delta_0 \star \Delta_0)_H \star \Delta_H \star (\Delta_1 \star \delta_1)_H\,,$$

we deduce that Θ_H visits $r_j, \ldots, r_{n-1}$. If we let $r_{j-1} = r_{\Theta_H}$, then the desired chain visited by Θ_H is $\{r_{j-1}, r_j, \ldots, r_{n-1}\}$, since $\epsilon(r_j) = \epsilon(a_j) \neq \epsilon(a_{j-1}) = \epsilon(r_{j-1})$. This concludes the proof of lemma 5.7. □

In order to prove theorem 5.2, choose a chain C of A, the cardinality of which is maximal, i.e. it is n. Define Γ_0 as it has been done in lemma 5.7, and let γ be the unique simple path from the initial position a_0 to $\operatorname{dom}\Gamma_0$. Using lemma 5.4, we can lift the path $\gamma \star \Gamma_0$ of A to a path $\theta_{\gamma\star\Gamma_0}$ of $\langle A, H, A\rangle$, played according to the strategy $\psi R || \phi S$ by the Mediators, such that both $(\theta_{\gamma\star\Gamma_0})_L$ and $(\theta_{\gamma\star\Gamma_0})_R$ are equal to $\gamma\star\Gamma_0$. Using lemma 5.5, we can also lift the given factorization to a factorization $\theta_{\gamma\star\Gamma_0} = \theta_\gamma \star \theta_{\Gamma_0}$ so that in particular $(\theta_{\Gamma_0})_L = (\theta_{\Gamma_0})_R = \Gamma_0$. Using lemma 5.7, the center projection $(\theta_{\Gamma_0})_H$ visits a chain in H of cardinality n which is, moreover, of the same color of the given chain C of A. This concludes the proof of theorem 5.2. □

6 Generalized Whitman polynomials

In this section we let $X = \{a_0, a_1, a_2, b_0, b_1, b_2\}$ be a set of six generators and define a bunch of synchronizing games $W^n \in \mathcal{K}(X)$, $n \geq 1$, such that $L(W^n) = n$. We can argue that $[W^n_\bullet] \notin \mathcal{J}_{m,X}$ if $n > m$, as in 5.3.

Theorem 6.1 For every $n \geq 1$, the inclusion $i_{n,X} : \mathcal{J}_{n,X} \hookrightarrow \mathcal{J}_X$ is proper, where X is a set of six generators.

Corollary 6.2 If P contains an antichain $\{a_0, a_1, a_2, b_0, b_1, b_2\}$, then the inclusion $i_{m,P} : \mathcal{J}_{m,P} \hookrightarrow \mathcal{J}_P$ is proper.

The strictness of the alternation hierarchy will follow as explained in 3.6. The construction of the games W^n has been suggested by the Whitman polynomial

$$p(x) \quad = \quad a \vee (b \wedge (c \vee (a \wedge (b \vee (c \wedge x))))) \, .$$

Earlier in Birkhoff's paper [Bir35], a partition lattice and an infinite chain of the form $p^n(a)$ are exhibited, so that the free lattice on three generators is shown to be infinite. A more functional interpretation to $p(x)$ is given in [Whi42]. P. Whitman proved that the free lattice on three generators is not complete by showing that free lattices are continuous – so that if the join of the infinite chain $p^n(a)$ exists, then it has to be a fixed point of $p(x)$ – and by proving that this polynomial has no fixed point. Later, Crawley and Dean [CD59] characterized free lattices with infinite operations and used the above polynomial to give lower bounds on the cardinality of those lattices, and a similar technique was used by Hales [Hal64] to show that the free complete lattices do not exist in general. Philip Whitman's result, also documented in the monography [FJN95], can be used to show that the inclusion $i_{0,X} : \mathcal{J}_{0,X} \hookrightarrow \mathcal{J}_X$ is proper when $\operatorname{card} X \geq 3$, where we recall that $\mathcal{J}_{0,X}$ coincides with the free lattice on the set X.

Definition 6.3 The game W^n is defined as follows:

- The set of position is

  $$\{ g_j, w_j \mid j = 0, \ldots, 6n-1 \}$$

  and the initial position is g_0.

- The set of forward edges is

$$\{\, g_j \to g_{j+1} \,|\, j = 0, \ldots, 6n-2 \,\} \quad \cup \quad \{\, g_j \to w_j \,|\, j = 0, \ldots, 6n-1 \,\}.$$

- The set of back edges is

$$\{\, g_{3(2n-k)-1} \to g_{3k} \,|\, k = 0, \ldots, n-1 \,\}.$$

- $\epsilon(w_j) = 0$, for $i = 1, \ldots, 6n-1$ and $\epsilon(g_j) = Q_{j \bmod 2}$, where $Q_0 = \sigma$ and $Q_1 = \pi$.
- Eventually, $\lambda(w_j) = a_{j \bmod 3}$, if $j < 3n$ and $\lambda(w_j) = b_{j \bmod 3}$, if $j \geq 3n$.

For example, the games W^1 and W^2 are pictured as follows:





We define $W(a_0, a_1, a_2)(x)$ to be

$$W(a_0, a_1, a_2)(x) \quad = \quad a_0 \wedge (a_1 \vee (a_2 \wedge x))\,,$$

and draw this equality by diagrams as



Similarly we let $W^*(a_0, a_1, a_2)(x)$ be the dual of $W(a_0, a_1, a_2)(x)$, i.e.

$$W^*(a_0, a_1, a_2)(x) \quad = \quad a_0 \vee (a_1 \wedge (a_2 \vee x))\,,$$

or by diagrams



Using these conventions, the game $W^n \in \mathcal{K}(\{a_0, a_1, a_2, b_0, b_1, b_2\})$ can be pictured as follows:



where we have supposed that n is an odd number. If n is even, W^n is pictured in a similar way.

Theorem 6.4 For each $n \geq 1$, the game $W^n \in \mathcal{K}(X)$ is synchronizing and $L(\,W^n\,) = n$.

It is easy to see that the game W^n is bipartite. Moreover:

Proposition 6.5 The unique strategy in the game $\langle W^n, W^n \rangle$ is the copycat strategy.

Proof. In the proof, we shall use the following notation: if g is a position of W^n such that $\epsilon(g) \neq 0$, we shall write w_g for the unique g' such that $g \to g'$ and $\epsilon(g') = 0$. We let $S^n(g)$ be the set of elements g' such that $\epsilon(g') \neq 0$ for which there exists a path of length n from g to g'. We shall use the notation S^n_g for any element $g' \in S^n(g)$, S_g will stand for S^1_g and we shall have $g = S^0_g$; with the above notation we must take care that identities like $S^n_g = S^n_g$ do not hold. The following lemma will prove to be useful.

Lemma 6.6 If $n \not\equiv m \bmod 3$, then $\lambda(w_{S^n_g}) \neq \lambda(w_{S^m_g})$.

Proof. It is enough to observe that if $g_j \to g_{j'}$ is an edge, then $j' \equiv j + 1 \bmod 3$. Hence, if $g = g_i$, $g' \in S^n(g)$ and $g'' \in S^m(g)$, then $\lambda(w_{g'}) \in \{a_{i+n \bmod 3}, b_{i+n \bmod 3}\}$ and $\lambda(w_{g''}) \in \{a_{i+m \bmod 3}, b_{i+m \bmod 3}\}$. However

$$\{a_{i+n \bmod 3}, b_{i+n \bmod 3}\} \cap \{a_{i+m \bmod 3}, b_{i+m \bmod 3}\} = \emptyset\,,$$

since $i + n \not\equiv i + m \bmod 3$. □

In order to prove proposition 6.5, we shall suppose that the game has reached a position of the form (g, g), with $\epsilon(g) \neq 0$. We shall suppose that $\epsilon(g) = \sigma$, and use a dual argument if $\epsilon(g) = \pi$. Depending on the Opponents' choice, we shall analyse the moves available to Mediator and show that the only possible reply is the one suggested by the copycat strategy. We shall draw trees to represent possible moves as well as winning strategies for the Opponents. Positions are labeled on the left by the player who must move. Dotted transitions are used for Mediator's moves leading to winning positions for the Opponents.

From (g, g), the Opponents have the following two types of moves:



From position (w_g, g), Mediator can play as follows:



Position (w_g, S_g) is winning for the Opponents, since they can move $(w_g, S_g) \to (w_g, w_{S_g})$, were they win because of lemma 6.6.

From position (S_g, g), Mediator can play as follows:



We exhibit a winning strategy for the Opponents from position (w_{S_g}, g):



We exhibit a winning strategy for the Opponents from position (S_g^2, g):

$$\begin{array}{ll} O & (S_g^2, g) \\ & \downarrow \\ M & (w_{S_g^2}, g) \longrightarrow (w_{S_g^2}, w_g) \\ & \downarrow \\ O & (w_{S_g^2}, S_g) \\ & \downarrow \\ M & (w_{S_g^2}, w_{S_g}) \end{array}$$

We exhibit a winning strategy for the Opponents from position (S_g, w_g):

$$\begin{array}{lll} M & (w_{S_g}, w_g) \longleftarrow & (S_g, w_g) \\ & & \downarrow \\ O & & (S_g^2, w_g) \\ & & \downarrow \\ M & & (w_{S_g^2}, w_g) \end{array}$$

To complete the argument, we must show that if (S_g, S_g) is a position such that $S_g \neq S_g$, i.e. it is of the form $(g_{3k}, g_{3(2n-k)})$ or $(g_{3(2n-k)}, g_{3k})$ with $k \in \{1, \ldots, n-1\}$, then the Opponents have a winning strategy. Suppose that $(S_g, S_g) = (g_{3k}, g_{3(2n-k)})$, so that $\lambda(w_{3k}) = a_0$ and $\lambda(w_{3(2n-k)}) = b_0$; the Opponents have the following strategy:

$$\begin{array}{lll} O & & (g_{3k}, g_{3(2n-k)}) \\ & & \downarrow \\ M & (w_{3k}, w_{3(2n-k)}) \longleftarrow & (g_{3k}, w_{3(2n-k)}) \\ & & \downarrow \\ O & & (S_g^2, w_{S_g}) \\ & & \downarrow \\ M & & (w_{S_g^2}, w_{S_g}) \end{array}$$

Similarly, the Opponents have a winning strategy from position $(g_{3(2n-k)}, g_{3k})$. This concludes the proof of proposition 6.5 $\square$

Proposition 6.7 The game W^n contains a σ-chain of cardinality n. Since card $R(W^n) = n$, we deduce that $L(W^n) = n$.

Proof. Firstly, we observe that $R(W^n) = \{\, g_{3k} \,|\, k = 0, \ldots, n-1 \,\}$, and then that $R(W^n)$ is itself a chain. For if $0 \leq k < k' \leq n-1$, then $3k' < 3(2n-k') - 1 < 3(2n-k) - 1$, so that $g_{3k'}$ lies on the cycle going from g_{3k} down to $g_{3(2n-k)-1}$ and back to g_{3k}. □

References

[AM99] S. Abramsky and P.-A. Melliès. Concurrent games and full completeness. In *Proceedings of the 14th. annual IEEE Symposium on Logic in Computer Science*, pages 431–442, 1999.

[Arn99] A. Arnold. The μ-calculus alternation hierarchy is strict on binary trees. *Theoretical Informatics and Applications*, (33):329–339, 1999.

[BÉ93] S. L. Bloom and Z. Ésik. *Iteration theories.* Springer-Verlag, Berlin, 1993.

[Bir35] G. Birkhoff. On the structure of abstract algebras. *Proc. Camb. Philos. Soc.*, 31:433–454, 1935.

[Bla92] A. Blass. A game semantics for linear logic. *Ann. Pure Appl. Logic*, 56(1-3):183–220, 1992.

[Bra98a] J. C. Bradfield. The modal mu-calculus alternation hierarchy is strict. *Theoret. Comput. Sci.*, 195(2):133–153, 1998.

[Bra98b] J. C. Bradfield. Simplifying the modal mu-calculus alternation hierarchy. In *STACS 98*, volume 1373 of *Lecture Notes in Comput. Sci.*, pages 39–49, 1998.

[CD59] P. Crawley and R. A. Dean. Free lattices with infinite operations. *Trans. Amer. Math. Soc.*, 92:35–47, 1959.

[FJN95] R. Freese, J. Ježek, and J. B. Nation. *Free lattices.* American Mathematical Society, Providence, RI, 1995.

[Hal64] A. W. Hales. On the non-existence of free complete Boolean algebras. *Fund. Math.*, 54:45–66, 1964.

[HJ99] H. Hu and A. Joyal. Coherence completions of categories. *Theoret. Comput. Sci.*, 227(1-2):153–184, 1999.

[Joy95a] A. Joyal. Free bicomplete categories. *C. R. Math. Rep. Acad. Sci. Canada*, 17(5):219–224, 1995.

[Joy95b] A. Joyal. Free bicompletion of enriched categories. *C. R. Math. Rep. Acad. Sci. Canada*, 17(5):213–218, 1995.

[Joy97] A. Joyal. Free lattices, communication, and money games. In *Logic and scientific methods*, pages 29–68, 1997. Proceedings of the 10th international congress of logic, methodology and philosophy of science, Florence, Italy, August 19–25, 1995.

[Len96] G. Lenzi. A hierarchy theorem for the μ-calculus. In *Automata, languages and programming*, volume 1099 of *Lecture Notes in Comput. Sci.*, pages 87–97, 1996.

[Niw85] D. Niwiński. Equational μ-calculus. In *Computation theory*, volume 208 of *Lecture Notes in Comput. Sci.*, pages 169–176, 1985.

[Niw86] D. Niwinski. On fixed-point clones. In *Automata, languages and programming*, volume 226 of *Lecture Notes in Comput. Sci.*, pages 464–473, 1986.

[San00a] L. Santocanale. Free μ-lattices. 2000. Accepted for the proceedings of Category Theory 99.

[San00b] L. Santocanale. *Sur les μ-treillis libres.* PhD thesis, Université du Québec à Montréal, July 2000. Available: http://www.brics.dk/∼luigis.

[Tho97] W. Thomas. Languages, automata, and logic. In G.Rozenberg and A. Salomaa, editors, *Handbook of formal languages. Vol. 3*, pages 389–455. Springer-Verlag, Berlin, 1997.

[Wal95] I. Walukiewicz. Completeness of Kozen's axiomatizations of the propositional μ-calculus. In *Proc. of the 10th IEEE Symp. on Logic in Computer Science*, 1995.

[Whi41] P. M. Whitman. Free lattices. *Ann. of Math. (2)*, 42:325–330, 1941.

[Whi42] P. M. Whitman. Free lattices. II. *Ann. of Math. (2)*, 43:104–115, 1942.

[Zie98] W. Zielonka. Infinite games on finitely coloured graphs with applications to automata on infinite trees. *Theoret. Comput. Sci.*, 200(1-2):135–183, 1998.

Recent BRICS Report Series Publications

RS-00-29 **Luigi Santocanale.** *The Alternation Hierarchy for the Theory of μ-lattices*. **November 2000. 44 pp. Extended abstract appears in** *Abstracts from the International Summer Conference in Category Theory*, **CT2000, Como, Italy, July 16–22, 2000.**

RS-00-28 **Luigi Santocanale.** *Free μ-lattices*. **November 2000. 51 pp. Short abstract appeared in** *Proceedings of Category Theory 99*, **Coimbra, Portugal, July 19-24, 1999. Full version to appear in a special conference issue of the** *Journal of Pure and Applied Algebra*.

RS-00-27 **Zoltán Ésik and Werner Kuich.** *Inductive -Semirings*. **October 2000. 34 pp.**

RS-00-26 **František Čapkovič.** *Modelling and Control of Discrete Event Dynamic Systems*. **October 2000. 58 pp.**

RS-00-25 **Zoltán Ésik.** *Continuous Additive Algebras and Injective Simulations of Synchronization Trees*. **September 2000. 41 pp.**

RS-00-24 **Claus Brabrand and Michael I. Schwartzbach.** *Growing Languages with Metamorphic Syntax Macros*. **September 2000.**

RS-00-23 **Luca Aceto, Anna Ingólfsdóttir, Mikkel Lykke Pedersen, and Jan Poulsen.** *Characteristic Formulae for Timed Automata*. **September 2000. 23 pp.**

RS-00-22 **Thomas S. Hune and Anders B. Sandholm.** *Using Automata in Control Synthesis — A Case Study*. **September 2000. 20 pp. Appears in Maibaum, editor,** *Fundamental Approaches to Software Engineering: First International Conference*, **FASE '00 Proceedings, LNCS 1783, 2000, pages 349–362.**

RS-00-21 **M. Oliver Möller and Rajeev Alur.** *Heuristics for Hierarchical Partitioning with Application to Model Checking*. **August 2000. 30 pp.**

RS-00-20 **Luca Aceto, Willem Jan Fokkink, and Anna Ingólfsdóttir.** *2-Nested Simulation is not Finitely Equationally Axiomatizable*. **August 2000. 13 pp.**

RS-00-19 **Vinodchandran N. Variyam.** *A Note on* $\mathrm{NP} \cap \mathrm{coNP/poly}$. **August 2000. 7 pp.**